

# HOMOTOPY ASSOCIATIVITY OF SPHERE EXTENSIONS

To the memory of Professor J. F. Adams

by N. IWASE

(Received 20th June 1988)

Throughout this paper, we work in the category of ( $p$ -localized) spaces having the homotopy type of connected CW-complexes of finite type with base point. We consider a principal bundle

$$G_{n-1} \rightarrow X \rightarrow S^{2dn-1}, \quad (0.1)$$

where  $G_n = SU(n)$ ,  $U(n)$  or  $Sp(n)$  and  $d = 1, 1$  or  $2$  respectively. In this case, the bundle is obtained as an induced bundle by a mapping  $f$  of base space  $S^{2dn-1}$  from the classical group extension as follows:

$$\begin{array}{ccc} G_{n-1} & \xlongequal{\quad} & G_{n-1} \\ \cap & & \cap \\ X & \xrightarrow{\quad \bar{f} \quad} & G_n \\ \downarrow & & \downarrow \\ S^{2dn-1} & \xrightarrow{\quad f \quad} & S^{2dn-1}. \end{array}$$

We denote  $X$  by  $M(n, \lambda)$  following Zabrodsky [19] when  $\deg(f) = \lambda$ . The problem is to describe, in terms of  $d$ ,  $n$  and  $\lambda$ , the condition when  $M(n, \lambda)$  becomes a homotopy associative  $H$ -space or more generally an  $A_m$ -space for  $m \geq 3$  (see Stasheff [17]). The case  $m = 2$  was studied by many authors (see Hilton–Roitberg [8], Stasheff [18], Curtis–Mislin [4], Sigrist–Suter [16] and Zabrodsky [19, 20, 21, 22]) and solved completely by 1972 as the following form.

**Fact 1.**  $M(n, \lambda)$  is an  $H$ -space if and only if one of the following three conditions is valid.

- (a)  $\lambda$  is odd
- (b)  $dn \leq 2$
- (c)  $\lambda = 0 \pmod{2d}$  and  $dn = 4$ .

To avoid a confusion with an integer mod  $p$ , we adopt the notation “a property  $P$  at

$p$ ”, rather than “a mod  $p$  property  $P$ ”, for a localized property  $P$  at  $p$ . Now let us turn our attention to homotopy associativity (or an  $A_3$ -structure) of sphere extensions. At first, Sigrist–Suter show in [15] that  $M(n, \lambda)$  is not  $A_3$  in case  $\lambda = 0 \pmod{4}$ ,  $d = 2$  and  $n = 2$  of Fact 1. By Hemmi [7], the result of Gonçalves [5] implies that  $M(n, \lambda)$  is not  $A_3$  at the prime 2 in case (c) of Fact 1. We summarize the above results.

**Fact 2.** *In case (c) of Fact 1,  $M(n, \lambda)$  is not  $A_3$  at the prime 2. In cases (a) and (b) of Fact 1,  $M(n, \lambda)$  is an  $A_\infty$ -space at the prime 2.*

Moreover Hemmi gives the necessary condition in [7] for  $p = 3$ , that is,  $\lambda$  is prime to 6, when  $dn = r \cdot 3^*$ ,  $(r, 3) = 1$  and  $r > 3$ , where we denote by  $3^*$  a power of 3. On the other hand, the sufficiency condition is considered by M. Mimura and the author in Section 6 of [14] more generally as the construction of new (higher) homotopy associative  $H$ -spaces. The purpose of this paper is to describe the condition in terms of  $d$ ,  $n$  and  $\lambda$ , working with a concept slightly stronger than homotopy associativity (or  $A_m$ -structure). Let  $Y$  be an  $A_m$ -space. Hopf’s theorem implies that  $Y$  is rationally equivalent to a product of Eilenberg–MacLane spaces  $\Pi_i K(Q, 2n_i - 1)$  which is a loop space. The space  $Y$  is defined to be  $A_m$ -primitive, following [14], if the rational equivalence preserves the  $A_m$ -structures, that is, it is an  $A_m$ -mapping.

**Theorem A.** *The following three conditions are equivalent for  $3 \leq m \leq \infty$ .*

- (1)  $M(n, \lambda)$  has an  $A_m$ -structure extending that of  $G_{n-1}$ .
- (2)  $M(n, \lambda)$  is an  $A_m$ -primitive  $A_m$ -space.
- (3) For every prime  $p \leq m$ , one of the following two is valid.
  - (a)  $\lambda$  is prime to  $p$
  - (b)  $p \geq dn$ .

**Remark 1.** If  $m$  is not a prime, the primitivity condition in (2) is omissible. And if  $dn \leq 2p$ ,  $A_p$ -structure supports  $A_p$ -primitivity for dimensional reasons.

**Theorem B.** *Let  $p$  be an odd prime. Then the following three conditions are equivalent.*

- (1)  $M(n, \lambda)$  is an  $A_p$ -primitive  $A_p$ -space at  $p$ .
- (2)  $M(n, \lambda)$  is an  $A_\infty$ -space (loop space or monoid) at  $p$ .
- (3)  $\lambda$  is prime to  $p$  or  $p \geq dn$ .

**Remark 2.** It is sufficient to prove for  $G_n = U(n)$  and  $Sp(n)$ , because  $U(n)$  has the homotopy type of  $S^1 \times SU(n)$ . So, we may consider only for the cases  $G_n = U(n)$  and  $Sp(n)$ .

**Remark 3.** (2) implies clearly (1). (3) implies that  $M(n, \lambda)$  is homotopy equivalent to  $G_n$  at  $p$ . Hence (3) implies (2).

We will show in Section 1 that Theorem B implies Theorem A. So, we shall show that

(1) implies (3) to prove Theorem B in cases  $G_n=U(n)$  and  $Sp(n)$ . To show this, we calculate that  $p$ -divisibility of Hubbuck operations (see [9, 10, 11]) on the projective space of  $M(n, \lambda)$ . Although the divisibility is not determined naturally and depends on the choice of a splitting of  $K$ -theory, the calculations on  $BU(n)$  can be applied on the suspension space of  $M(n, \lambda)$ .

The author thanks the Department of Mathematics of University of Aberdeen for its hospitality. He also thanks John Hubbuck for discussions about  $K$ -theory operations without which work on this paper would not have begun and also Michael Crabb and many other persons in University of Aberdeen for conversations which helped to organize my thoughts.

**1. Proof of Theorem A from Theorem B**

Let  $\Pi$  be the set of all primes,  $\mathbb{P}_1$  the set of primes  $p$  with  $p \geq dn$ ,  $\mathbb{P}_2$  the set of primes  $p$  with  $dn > p > m$  and  $\mathbb{P}_3 = \Pi - \mathbb{P}_1 - \mathbb{P}_2$ . Then  $G_n$  has the homotopy type of a product of spheres at  $\mathbb{P}_1$ . In particular, the bundle (0.1) is trivial. Hence, the pull-back  $M(n, \lambda)$  is also trivial and homotopy equivalent to  $G_n$  at  $\mathbb{P}_1$ . Hence  $M(n, \lambda)$  has an  $A_\infty$ -structure extending that of  $G_{n-1}$  at  $\mathbb{P}_1$ . Secondly, we may write  $\lambda = \lambda_1 \cdot \lambda_2$  where  $(\lambda_1, \mathbb{P}_2) = 1$  and  $(\lambda_2, \Pi - \mathbb{P}_2) = 1$ . Let  $q = \text{Min } \mathbb{P}_2$  where we regard  $\text{Min } \phi = \infty$ . Then  $M(n, \lambda)$  is homotopy equivalent to  $M(n, \lambda_2)$  at  $\mathbb{P}_2$  which has an  $A_{q-1}$ -structure extending that of  $G_{n-1}$ , by Theorem 6.5 of [14]. Therefore,  $M(n, \lambda)$  has an  $A_{q-1}$ -structure extending that of  $G_n$ , if and only if it has at  $\mathbb{P}_3$ , by the property (P7) of [14].

Firstly we assume (3). Then  $f$  is a homotopy equivalence between  $M(n, \lambda)$  and  $G_n$  at  $\mathbb{P}_3$  and  $G_n$  has an  $A_\infty$ -structure extending that of  $G_{n-1}$ . This implies (1). Secondly we assume (1). Then by [13], it follows that the generators of  $H^*(M(n, \lambda); \mathbb{Q})$  are all  $A_{m-1}$ -primitive and therefore represented by  $A_{m-1}$ -mappings, by the property (P9) of [14]. By the proof of the Corollary in [13], the obstruction to be  $A_m$ -primitive is in  $H^{2i}(M(n, \lambda) * \dots * M(n, \lambda); \mathbb{Q}) = H^{2i}(G_{n-1} * \dots * G_{n-1}; \mathbb{Q})$ ,  $i \leq n$ . (1) implies that the inclusion mapping  $G_{n-1} \rightarrow M(n, \lambda)$  induces a homomorphism of spectral sequences of Stasheff's type (see [17, 13]). For dimensional reasons, the obstructions are mapped to 0 by the injective homomorphism induced from the inclusion. Hence the generators are represented by  $A_m$ -mappings and  $M(n, \lambda)$  is  $A_m$ -primitive. This implies (2). Thirdly, we assume (2). Then by Fact 1 and Fact 2, it follows that  $\lambda$  is odd or  $dn \leq 2$ , since  $m$  is greater than or equal to 3. For an odd prime  $p \leq m$ , (2) implies that  $M(n, \lambda)$  is an  $A_p$ -primitive space at  $p$ . Then by Theorem B, we obtain (3). This completes the proof of Theorem A.

**2. Decomposition of  $BU$  and  $\tilde{K}(BT^n)$  at an odd prime**

Let  $R$  be the ring of localized integers at an odd prime  $p$ ,  $BU_{(p)}$  the localization of  $BU$  and  $\tilde{K}(X) = \tilde{K}(X; R) = [X, BU_{(p)}]$ . By Adams [3],  $BU_{(p)}$  is decomposable to  $p-1$  factors such as  $BU_{(p)} \simeq BU^{(1)} \times \dots \times BU^{(p-1)}$  and the Chern character is also decomposable to  $p-1$  factors

$$ch^{(i)}: BU^{(i)} \rightarrow \prod_{j \geq 0} K(\mathbb{Q}, i + j(p-1)).$$

We denote by  $\tilde{K}(X)^{(i)}$  the factor  $[X, BU_{(p)}^{(i)}]$  of  $\tilde{K}(X)$ . Then it follows that  $\tilde{K}(X) \simeq \tilde{K}(X)^{(1)} \oplus \cdots \oplus \tilde{K}(X)^{(p-1)}$ . For  $X = BT$ , we have

$$\begin{aligned} \tilde{K}(BT)^{(i)} &= R[[x^{p-1}]] \cdot x^i, \\ x \in \tilde{K}(BT)^{(1)}, ch^{(1)}(x) &= \sum_{j \geq 0} \frac{\bar{\alpha}_j}{(1+j(p-1))!} y^{1+j(p-1)}, \end{aligned} \quad (2.1)$$

with  $\bar{\alpha}_0 = 1$  and  $\bar{\alpha}_j \in \mathbb{Z}$ .

Then by (2.1), it follows that, for  $X = BT^n$ ,

$$\tilde{K}(BT^n)^{(i)} = R\{X_{a_1}^{i_1 + j_1(p-1)} \times \cdots \times X_{a_m}^{i_m + j_m(p-1)}\}$$

where  $j_i \geq 0$ ,  $i_1 + \cdots + i_m = i$ ,  $1 \leq a_1 < \cdots < a_m \leq n$ ,  $m \leq n$  and  $x_a \in \tilde{K}(BT^n)^{(1)}$  corresponds to the generator of the  $a$ th factor of  $BT^n$ .

Using  $x_a$  as above, we write  $K$ -algebras  $K(BU(n))$  and  $K(BSp(n))$  as follows:

$$K(BU(n)) = R[[c_1^K, \dots, c_n^K]] \cong R[[x_1, \dots, x_n]]^{\Sigma_n},$$

$$K(BSp(n)) = R[[c_2^K, \dots, c_{2n}^K]],$$

where  $c_i^K \in K(BU(n))^{(i)}$  is mapped to  $\sigma_i(x_1, \dots, x_n)$  by the monomorphism  $K(BU(n)) \rightarrow K(BT^n)$ ,  $\sigma_i$  is the  $i$ th elementary symmetric polynomial and  $\sigma_n$  is the symmetric group on  $n$  letters.

**Remark 2.1.**  $c_i^K$  is the class obtained by modifying the  $\gamma$ -class so that  $ch(c_i^K)$  lies in  $\Pi_{j \geq 0} H^{2i+2j(p-1)}(BU; \mathbb{R})$ . Hence  $c_{2i+1}^K$  is mapped to 0 in  $K(BSp(n))$ .

### 3. Hubbuck operations in $K(BU(n))$

Let  $E$  be the fake  $R \times BU_{(p)}$  such as  $E = \Pi_{j \geq 0} K(R, 2j)$  and  $E(X) = [X, E]$ . Then  $E(BU(n)) \simeq R[[c_1, \dots, c_n]]$  and  $E(BT^n) \simeq R[[y_1, \dots, y_n]]$ , where  $c_1$  is the  $i$ th Chern class and is mapped to  $\sigma_i(y_1, \dots, y_n)$  by the ring monomorphism  $E(BU(n)) \rightarrow E(BT^n)$ .

To define Hubbuck operation, we need a splitting. Let us define the ring isomorphisms  $J: E(BU(n)) \rightarrow K(BU(n))$  and  $J_0: E(BT^n) \rightarrow K(BT^n)$  as follows:

$$J(c_i) = c_i^K,$$

$$J_0(y_i) = x_i.$$

We regard the algebras  $K(BU(n))$  and  $E(BU(n))$  as the subalgebra of  $K(BT^n)$  and  $E(BT^n)$ , respectively. Then it follows that  $J$  can be regarded as the restriction of  $J_0$  to  $E(BU(n))$  and so we often denote  $J_0$  by  $J$ . We wish to know the manner of Hubbuck operations on  $c_i^K$  in  $K$ -theory.

Let us recall the Chern character on  $K(BT)$ . By (2.1), it follows that

$$ch(x) = \sum_{j \geq 0} \frac{\alpha_j}{p^j} y^{j(p-1)+1},$$

where  $\alpha_j = \bar{\alpha}_j \cdot p^j / (1 + j(p-1))!$  is in  $R$ , because  $(q+1)!$  divides  $m(q) = \prod_{\text{all primes}} p^{[q/(p-1)]}$  (see Adams [1]). To simplify notation, we introduce some functions in  $\mathbb{Q}[[t]]$  where  $t$  is transcendental:

$$e(t) = \sum_j \frac{\alpha_j}{p^j} t^{j(p-1)+1}.$$

Since  $d/dt(e(t))|_{t=0} = \alpha_0 = \bar{\alpha}_0 = 1$ ,  $e(t)$  has the inversion  $\ell(t)$  in  $\mathbb{Q}[[t]]$ . We choose local integers  $\beta_j$  in  $R$  such that

$$\ell(t) = \sum_j \frac{\beta_j}{p^j} t^{j(p-1)+1} \quad \text{with } \beta_0 = 1.$$

Then it follows that

$$ch(x) = e(y) \quad \text{and} \quad ch(\ell(x)) = \ell(e(y)) = y. \tag{3.1}$$

We will describe Adams operations by using  $e$  and  $\ell$ .

Firstly we will define a fake Adams operation  $\Psi^k$  on the fake  $K$ -theory  $E$  (see Hubbuck [9, 10, 11]) and reserve the symbol  $\psi^k$  for the genuine Adams operation.

**Definition 3.1.** The fake Adams operation  $\Psi^k$  on  $E(-)$  is defined by the following formula:

$$\Psi^k(x_n) = k^n \cdot x_n \quad \text{for } x_n \in H^{2n}(X; R).$$

Then the Chern character commutes with (fake) Adams operations  $\psi^k$  and  $\Psi^k$ . Therefore the Adams operation preserves the mod  $p$  decomposition of (fake)  $K$ -theories. So, we may write for the generator  $x$  of  $K(BT)$ ,

$$\psi^k(x) = \sum_{j \geq 0} \bar{r}_j(k) \cdot x^{j(p-1)+1}.$$

where  $\bar{r}_j(k)$  is a local integer in  $R$ ,  $\bar{r}_0(k) = k$  and  $\bar{r}_1(p) = 1 \pmod p$ . On the other hand, we can compute the Adams operation by using  $e$  and  $\ell$  as follows:

$$\psi^k(x) = \psi^k(e(\ell(x))) = e(\psi^k(\ell(x)))$$

and

$$ch(\psi^k(\ell(x))) = \Psi^k(ch(\ell(x))) = \Psi^k(y) = k \cdot y = k \cdot ch(\ell(x)) = ch(k \cdot \ell(x)).$$

Therefore, we obtain  $\psi^k(\ell(x)) = k \cdot \ell(x)$  and then it follows that

$$\begin{aligned}\psi^k(x) &= e(k \cdot \ell(x)) \\ &= \sum_{j \geq 0} \frac{\alpha_j}{p^j} \cdot k^{j(p-1)+1} \cdot \ell(x)^{j(p-1)+1} \\ &= \sum_{q \geq 0} \frac{k}{p^q} \left( \sum_{q=j+1} \alpha_j \cdot k^{j(p-1)} \cdot \beta_{j,i} \right) \cdot x^{q(p-1)+1},\end{aligned}$$

where  $\beta_{j,i} \in R$  is given by the formula

$$\ell(x)^{j(p-1)+1} = \sum_{j \geq 0} \frac{\beta_{j,i}}{p^i} \cdot x^{(i+j)(p-1)+1}.$$

Using  $\alpha_j$  and  $\beta_{j,i}$ , we can define more “stabilized” decomposition of the Adams operation  $\psi^k$  by the following formula

$$\psi^k(x) = k \cdot r(k; x)$$

where

$$\begin{aligned}r(k; t) &= \sum_{j \geq 0} \frac{r_j(k)}{p^j} \cdot t^{j(p-1)+1}, \\ r_q(k) &= \sum_{q=j+i} \alpha_j \cdot k^{j(p-1)} \cdot \beta_{j,i}.\end{aligned}$$

We remark that  $\bar{r}_j(k)$  and  $r_j(k)$  has the following relation

$$\bar{r}_j(k) = \frac{k \cdot r_j(k)}{p^j}. \quad (3.2)$$

We have prepared to describe the Hubbuck operations on  $K(BU(n))$ . Let us define  $Q_i$ ,  $S_i$  and  $R(k)_i$  in the ring  $Q[[t_1, \dots, t_n]]$  as follows:

$$\begin{aligned}Q_i(t_1, \dots, t_n) &= \sigma_i(e(t_1), \dots, e(t_n)) \\ &= \sum_{j \geq 0} \frac{1}{p^j} \cdot Q_i^j(t_1, \dots, t_n), \\ S_i(t_1, \dots, t_n) &= \sigma_i(\ell(t_1), \dots, \ell(t_n)) \\ &= \sum_{j \geq 0} \frac{1}{p^j} \cdot S_i^j(t_1, \dots, t_n),\end{aligned} \quad (3.3)$$

$$\begin{aligned}
 R(k)_i(t_1, \dots, t_n) &= \sigma_i(r(k; t_1), \dots, r(k; t_n)) \\
 &= \sum_{j \geq 0} \bar{R}^j(k)_i(t_1, \dots, t_n)
 \end{aligned}$$

where  $Q_i^j$ ,  $S_i^j$  and  $\bar{R}^j(k)_i$  are in  $R[t_1, \dots, t_n]^{\Sigma^n}$  and are written as polynomials of the elementary symmetric polynomials  $\sigma_1, \dots, \sigma_n$  of  $t_1, \dots, t_n$ , if  $(k, p) = 1$ . The following equations can be easily checked:

$$ch(c_i^k) = Q_i(y_1, \dots, y_n),$$

$$ch(S_i(x_1, \dots, x_n)) = c_i$$

$$\psi^k(c_i^k) = k^i \cdot R(k)_i(x_1, \dots, x_n), \quad \text{when } (k, p) = 1.$$

Next we define  $R$ -endomorphisms of  $K(BU(n))$  extending the following relation by the Cartan formula (see Hubbuck [9, 10, 11]):

$$Q^j(c_i^k) = Q_i^j(x_1, \dots, x_n),$$

$$S^j(c_i^k) = S_i^j(x_1, \dots, x_n),$$

$$\bar{R}^j(k)(c_i^k) = \bar{R}^j(k)_i(x_1, \dots, x_n), \quad \text{when } (k, p) = 1.$$

Using them, we define  $Q(t)$ ,  $S(t)$  and  $R(k; t)$  in  $\text{End}_R(K(BU(n))) \otimes_R \mathbb{Q}[[t]]$ , by the following formula

$$Q(t) = \sum_j \frac{1}{p^j} Q^j \cdot t^{j(p-1)},$$

$$S(t) = \sum_j \frac{1}{p^j} S^j \cdot t^{j(p-1)},$$

$$R(k; t) = \sum_j \bar{R}^j(k) \cdot t^{j(p-1)}.$$

Then by the definition, we obtain

**Proposition 3.1.** *The following four equations are valid:*

- (1)  $Q = J \circ ch$  and  $ch \circ S = J^{-1}$ ,
- (2)  $S \circ Q = Q \circ S = \text{Identity}$  and  $\psi^k \circ S \circ J(w_i) = k^i \cdot S \circ J(w_i)$ ,
- (3)  $\psi^k \circ J(w_i) = k^i \cdot R(k) \circ J(w_i)$ ,
- (4)  $\bar{R}^j(k) = 1/p^j \sum_{i=0}^j k^{i(p-1)} \cdot S^{j-i} \circ Q^i$ ,

where  $w_i$  is in  $H^{2i}(BU(n); R)$ .

**Proof.** (1) and (3) are obtained directly by the definitions of the Hubbuck operations on  $c_i^k$  together with the Cartan formulae. Firstly we show (2). By (1),  $J \circ ch \circ S = \text{Identity}$ . This implies that  $Q \circ S = \text{Identity}$  and therefore,  $S \circ Q = Q \circ S = \text{Identity}$ . Similarly, we have  $ch \circ S \circ J = \text{Identity}$ . This implies that  $ch \circ \psi^k \circ S \circ J = \Psi^k \circ ch \circ S \circ J = \Psi^k$  and  $ch \circ \psi^k \circ S \circ J(w_i) = \psi^k(w_i) = k^i \cdot w_i = k^i \cdot ch \circ S \circ J(w_i)$ . Hence  $\psi^k \circ S \circ J(w_i) = k^i \cdot S \circ J(w_i)$ . To show (4), it suffices to show

$$R(k) \circ J(w_i) = \sum_q (1/p^q) \cdot \sum_{q \geq j \geq 0} k^{j(p-1)} \cdot S^{q-j} \circ Q^j \circ J(w_i),$$

because both  $R^q(k)$  and  $S^{q-j} \circ Q^j$  increase the same weight  $q(p-1)$ . By (3), we obtain that

$$\begin{aligned} k^i \cdot R(k) \circ J(w_i) &= \psi^k \circ J(w_i) \\ &= \psi^k \circ S \circ Q \circ J(w_i) \\ &= \sum_j \frac{1}{p^j} \cdot \psi^k \circ S \circ Q^j \circ J(w_i). \end{aligned}$$

Here,  $Q^j \circ J(w_i)$  has the weight  $i + j(p-1)$  and then by (3), we proceed as follows:

$$\begin{aligned} k^i \cdot R(k) \circ J(w_i) &= \sum_j \frac{k^{i+j(p-1)}}{p^j} \cdot S \circ Q^j \circ J(w_i) \\ &= k^i \cdot \sum_q (1/p^q) \cdot (\sum_{q \geq j \geq 0} k^{j(p-1)} \cdot S^{q-j} \circ Q^j) \circ J(w_i). \end{aligned}$$

This completes the proof of Proposition 3.1.

#### 4. $p$ -divisibility

Before starting to prove Theorem B, we will show the key lemma of this paper. We denote by  $v_p$  the valuation of the ring of  $p$ -localized integers  $R$ , that is,  $v_p(m)$  is the largest power of  $p$  dividing  $m$ .

From now we assume that  $k = p-1$ . Then by Adams [2] or Hubbuck [10, Lemma 4.3], it follows that

$$v_p(k^{j(p-1)} - 1) = v_p(j) + 1. \quad (4.1)$$

Firstly we show the  $p$ -divisibility of Hubbuck operations in  $K(BU(n))$ .

\*If we take  $k=2$ , this equality fails for  $p=1093$  (see [6]).



**Lemma 4.1** *Let  $i \cdot n = m + j(p-1)$ ,  $i \geq 1$ ,  $n \geq m \geq 1$ . If  $v_p(m) \leq v_p(n)$ , then the coefficient of  $(c_n^K)^i$  in  $\bar{R}^j(k)(c_m^K)$  is divisible by  $n/m$  in  $R$ .*

**Proof.** We write  $\bar{R}^j(k)_m(t_1, \dots, t_n) = P_m^j(\sigma_1, \dots, \sigma_n)$  in  $R[\sigma_1, \dots, \sigma_n]$ , where  $\sigma_i$  is the  $i$ th symmetric polynomial of  $t_1, \dots, t_n$ . Then the desired coefficient is given by  $P_m^j(0, \dots, 0, 1) = \bar{R}^j(k)_m(\zeta, \zeta^2, \dots, \zeta^n)$ , where  $\zeta$  is the primitive  $n$ th root of unity in the complex number field. Using the definition (3.3), we write

$$\bar{R}^j(k)_m(t_1, \dots, t_n) = \sum_{j_1, \dots, j_m} \beta_{j_1} \cdots \beta_{j_m} \cdot L_m^{j_1, \dots, j_m}(t_1, \dots, t_n),$$

where  $j_1, \dots, j_m$  run over all integers such that  $0 \leq j_1 \leq \dots \leq j_m \leq j = j_1 + \dots + j_m$  and the polynomial  $L_m^{\dots}$  is given by

$$L_m^{j_1, \dots, j_m}(t_1, \dots, t_n) = \sum_{a_1}^{j_1(p-1)+1} \dots \sum_{a_m}^{j_m(p-1)+1},$$

where  $j'_1, \dots, j'_m$  and  $a_1, \dots, a_m$  run over the set  $\Delta$  given by  $\{(a_1, \dots, a_m; j'_1, \dots, j'_m) \mid 1 \leq a_1 < \dots < a_m \leq n, (j'_1, \dots, j'_m) = (j_1, \dots, j_m) \text{ if we ignore the ordering}\}$ , by the following calculation:

$$\begin{aligned} \sigma_m(\ell(t_1), \dots, \ell(t_n)) &= \sum_{1 \leq a_1 < \dots < a_m \leq n} \ell(t_{a_1}) \cdots \ell(t_{a_m}) \\ &= \sum_j \frac{1}{p^j} \sum_{j=j_1+\dots+j_m} \beta_{j_1} \cdots \beta_{j_m} \cdot \sum_{1 \leq a_1 < \dots < a_m} t_{a_1}^{j_1(p-1)+1} \cdots t_{a_m}^{j_m(p-1)+1} \\ &= \sum_j \frac{1}{p^j} \sum_{j_1, \dots, j_m} \beta_{j_1} \cdots \beta_{j_m} \cdot L_m^{j_1, \dots, j_m}(t_1, \dots, t_n). \end{aligned}$$

Hence we obtain that

$$P_m^j(0, \dots, 0, 1) = \sum_{j_1, \dots, j_m} \beta_{j_1} \cdots \beta_{j_m} \cdot L_m^{j_1, \dots, j_m}(\zeta, \dots, \zeta^n).$$

$$L_m^{j_1, \dots, j_m}(\zeta, \dots, \zeta^n) = \sum_{\delta \in \Delta} \zeta^{g(\delta)},$$

where  $j_1, \dots, j_m$  run over  $0 \leq j_1 \leq \dots \leq j_m \leq j = j_1 + \dots + j_m$ ,  $g(\delta) = a_1(j'_1(p-1)+1) + \dots + a_m(j'_m(p-1)+1)$  in the cyclic group of order  $n$  and  $\delta = (a_1, \dots, a_m; j'_1, \dots, j'_m)$ . We remark here that  $L_m^{\dots}$  is a localized integer. So, we are left to show that the localized integer  $L_m^{\dots}$  is divisible by  $n/m$  if  $v_p(n) \geq v_p(m)$ .

Let  $\tau$  be the element of  $\Sigma_n$  such that  $\tau(a) = a+1 \pmod n$  and  $\sigma$  the element of  $\Sigma_m$  such that  $1 \leq \tau(a_{\sigma(1)}) < \dots < \tau(a_{\sigma(m)}) \leq n$ . Then  $\sigma = \text{Identity}$  or  $\sigma(i) = i-1 \pmod m$  for all  $i$ . We remark that  $\sigma$  depends on both  $\tau$  and  $(a_1, \dots, a_m)$ . Let  $\delta = (a_1, \dots, a_m; j'_1, \dots, j'_m)$  and  $\tau \cdot \delta = (a_{\sigma(1)}, \dots, a_{\sigma(m)}; j'_{\sigma(1)}, \dots, j'_{\sigma(m)})$ . Then we obtain that

$$g(\tau \cdot \delta) = g(\delta) \text{ in } Z/nZ,$$




FIGURE 1

because  $i \cdot n = j(p-1) + m = (j_1(p-1) + 1) + \dots + (j_m(p-1) + 1)$ . Therefore we obtain the following equation

$$\sum_{\delta \in \Delta} \zeta^{g(\delta)} = \sum_{[\delta] \in \Delta'} n(\tau, \delta) \cdot \zeta^{g(\delta)},$$

where  $\Delta'$  is the quotient set of  $\Delta$  by the action of  $Z/nZ$  and  $n(\tau, \delta)$  is the cardinality of the set  $\{\delta, \tau\delta, \dots, \tau^{n-1}\delta\}$ . Hence,  $n(\tau, \delta)$  divides  $n$ .

On the other hand, the equation  $\xi^{n(\tau, \delta)} \delta = \delta$  implies  $a_{\sigma'(i)} + n(\tau, \delta) = a_i \pmod n$  and  $\sigma'(i) = i - m(\tau, \delta) \pmod m$  for some  $\sigma' \in \Sigma_m$  and  $1 \leq m(\tau, \delta) \leq m$ . Therefore, we obtain the equation

$$\begin{aligned} m(\tau, \delta) &= \#(\{a_1, \dots, a_m\} \cap [1, n(\tau, \delta)]) \\ &= \#(\{a_1, \dots, a_m\} \cap [n(\tau, \delta) + 1, 2n(\tau, \delta)]) \\ &\quad \vdots \\ &= \#(\{a_1, \dots, a_m\} \cap [n - n(\tau, \delta) + 1, n]). \end{aligned}$$

This implies  $m(\tau, \delta)$  divides  $m$  and  $m/m(\tau, \delta) = n/n(\tau, \delta)$ . Hence  $n(\tau, \delta) = n \cdot m(\tau, \delta)/m$  and

$$L_m^{j_1, \dots, j_m}(\xi, \dots, \xi^n) = (n/m) \cdot \sum_{[\delta] \in \Delta'} m(\tau, \delta) \cdot \zeta^{g(\delta)}$$

in the ring  $R$  if  $v_p(n) \geq v_p(m)$ . This implies the lemma.

### 5. Proof of Theorem B

We assume (1). To construct a Hubbuck operation on the projective spaces  $P(m)$  and  $\bar{P}(m)$  for  $G_n$  and  $M(n, \lambda)$ , we need a splitting from the fake  $K$ -theory  $E^*$  to  $K$ -theory  $K^*$ . Let us recall that  $K^*(P(m)) = M \oplus S_m$  and  $K^*(\bar{P}(m)) = \bar{M} \oplus \bar{S}_m$  where  $M$  and  $\bar{M}$  are polynomial algebras truncated at height  $m+1$  and  $S_m$  and  $\bar{S}_m$  are ideals (see [13]). By the definition of  $S_m$  and  $\bar{S}_m$ , it follows that  $\psi^k(S_m) \subset S_m$  and  $\psi^k(\bar{S}_m) \subset \bar{S}_m$ . In the proof given in [13], it is required that the  $K$ -theory of  $H$ -spaces has no torsion and that  $H$ -spaces are  $A_m$ -primitive. No other assumption is required. So, we obtain the following isomorphisms similarly to [13]:  $E^*(P(m)) = N \oplus T_m$  and  $E^*(\bar{P}(m)) = \bar{N} \oplus \bar{T}_m$  where  $N$  and  $\bar{N}$  are polynomial algebras truncated at height  $m+1$  and  $T_m$  and  $\bar{T}_m$  are ideals.

Let  $\eta_n$  be the canonical  $n$ -bundle over  $BG_n$  (complex or quaternionic). Then  $M$  is generated by  $c_{di}^K(\eta_n)$  for  $i \leq n$  and  $N$  is generated by  $c_{di}(\eta_n)$  for  $i \leq n$ . Let  $QM$  and  $Q\bar{M}$  be the indecomposable quotients of  $M$  and  $\bar{M}$ , respectively. Then by [13], it follows that  $QM \simeq QK^*(G_n)$  and  $Q\bar{M} \simeq QK^*(M(n, \lambda))$  whose generators are corresponding to each other by the homomorphism induced by  $\bar{f}$  except for the generators in exact filtration

degree  $2dn - 1$ . In the filtration degree  $2dn - 1$ , the generators are spherical and  $\bar{f}^1$  times  $\lambda$  on the generators. Hence we obtain

**Proposition 5.1.**

- (1)  $M$  is generated by  $u_i = c_{di}^K(\eta_n)$  for  $i \leq n$ ,
- (2)  $\bar{M}$  is generated by  $\bar{u}_i$  for  $i \leq n$ ,
- (3)  $\bar{u}_i = \Sigma \bar{f}^1(u_i)$  in  $Q\bar{M}$  for  $i < n$  and
- (4)  $\lambda \cdot \bar{u}_n = \Sigma \bar{f}^1(u_n)$  in  $Q\bar{M}$ .

**Proposition 5.2.**

- (1)  $N$  is generated by  $v_i = c_{di}(\eta_n)$  for  $i < n$ ,
- (2)  $\bar{N}$  is generated by  $\bar{v}_i$  for  $i \leq n$ ,
- (3)  $\bar{v}_i = \Sigma \bar{f}^1(v_i)$  in  $Q\bar{N}$  for  $i < n$  and
- (4)  $\lambda \cdot \bar{v}_n = \Sigma \bar{f}^1(v_n)$  in  $Q\bar{N}$ .

Then we define the splittings  $J$  and  $\bar{J}$  by the following equations:

$$J(v_i) = u_i, \quad i \leq n \quad \text{and}$$

$$\bar{J}(\bar{v}_i) = \bar{u}_i, \quad i \leq n.$$

The mapping  $\bar{f}$  induces the following homomorphism  $\phi$ :

$$\begin{aligned} \phi(u_i) &= \bar{u}_i, \quad i < n, \quad \text{and} \\ \phi(u_n) &= \lambda \cdot \bar{u}_n \quad \text{in } Q\bar{M}. \end{aligned} \tag{5.1}$$

**Remark 5.3.** If one extends  $\phi$  as a ring homomorphism, then  $\phi$  does not commute with Adams operations, even if  $\bar{f}$  is an  $A_m$ -mapping. Also  $\bar{f}$  induces the following homomorphism  $\phi_0$ :

$$\begin{aligned} \phi_0(v_i) &= \bar{v}_i, \quad i < n, \quad \text{and} \\ \phi_0(v_n) &= \lambda \cdot \bar{v}_n, \quad \text{in } Q\bar{N}. \end{aligned} \tag{5.2}$$

By Hubbuck [9, 10], these splitting  $J$  and  $\bar{J}$  determine  $K$ -theory operations  $S_J^h, S_{\bar{J}}^h, Q_J^h, Q_{\bar{J}}^h, \bar{R}_J^h$  and  $\bar{R}_{\bar{J}}^h$ , which now satisfy

$$\bar{R}_{\bar{J}}^h \circ \phi_0 = \phi_0 \circ \bar{R}_J^h \quad \text{in } QN, \tag{5.3}$$

since  $\phi \circ J = \bar{J} \circ \phi_0$  by (5.1) and (5.2).

We will write the Hubbuck operations by  $S^h, Q^h, R^h$  and  $\bar{R}^h$  when the formula is valid

independently of the choice of a splitting. The following formulae are due to Hubbuck (see [9, 10]):

**Proposition 5.4.**

- (1)  $\bar{R}^h$  is an integral operation,
- (2)  $\bar{R}^h$  satisfies the Cartan formula  $\bar{R}^h(x \cdot y) = \sum_{i+j=h} \bar{R}^i(x) \cdot \bar{R}^j(y)$ ,
- (3)  $S^{di}(v_i) = v_i^p \pmod{p}$ ,
- (4)  $S^{di}(\bar{v}_i) = \bar{v}_i^p \pmod{p}$ ,
- (5)  $(1 - k^{q(p-1)}) \cdot S^q = \sum_{h=1}^{q-1} k^{(q-h)(p-1)} \cdot p^h \cdot \bar{R}^h \circ S^{q-h} + p^q \cdot \bar{R}^q$ , where  $k = p - 1$ .

**Remark 5.5.** By the definition of  $J$ ,  $S^h$ ,  $Q_J^h$  and  $\bar{R}_J^h$  coincide with the restriction to  $P(m)$  of  $S^h$ ,  $Q^h$  and  $\bar{R}^h(k)$  respectively given in Section 4, if we identify  $K$ -theory with fake  $K$ -theory by the splitting  $J$  above.

Assuming that  $\lambda = 0 \pmod{p}$  and  $dn > p$ , we are led to a contradiction.

Let  $a = v_p(dn)$ . Then by a simple computation,  $(a+1)(p-1) < dn$ . By Proposition 5.4, we obtain the following proposition similarly to Hubbuck–Mimura [12].

**Proposition 5.6.** *The following two statements are valid in  $QM$ :*

- (1)  $p^{a+1} \cdot v_n \in p^h \cdot \bar{R}_J^h(QN^{dm}) \pmod{I + (p^{a+2})}$
- (2)  $p^{a+1} \cdot \bar{v}_n \in p^{h'} \cdot \bar{R}_J^{h'}(Q\bar{N}^{dm}) \pmod{\bar{I} + (p^{a+2})}$

for some  $1 \leq h, h' \leq a+1$ , where  $m = n - h(p-1)/d$ ,  $m' = n' - h'(p-1)/d$ ,  $I = (v_1, \dots, v_{n-1})$  and  $\bar{I} = (\bar{v}_1, \dots, \bar{v}_{n-1})$ .

**Proof.** The formulae given in (5.4) imply that

$$(1 - k^{dn(p-1)}) \cdot v_n^i \in p^h \cdot \bar{R}_J^h(QN^{dm}) \pmod{I + (p^{a+2})}$$

$$(1 - k^{dn(p-1)}) \cdot \bar{v}_n^{i'} \in p^{h'} \cdot \bar{R}_J^{h'}(Q\bar{N}^{dm'}) \pmod{\bar{I} + (p^{a+2})}$$

for some  $1 \leq h, h' \leq a+1$  such that  $in = m + h(p-1)/d$ ,  $i'n = m' + h'(p-1)/d$  and  $1 \leq i, i' \leq p$ . For dimensional reasons, in the formula above, we obtain that  $i = i' = 1$ . Then by (4.1), Proposition 5.6 follows.

By Lemma 4.1 and Proposition 5.6, it follows that

$$p^{a+1} \cdot v_n \in p \cdot \bar{R}_J^1(QN^{dn-(p-1)}) \pmod{p^{a+2}} \quad \text{in } QN^{dn}.$$

Also by (5.3) together with Lemma 4.1 and Proposition 5.6, it follows that

$$p^{a+1} \cdot \bar{v}_n \in p \cdot \bar{R}_J^1(Q\bar{N}^{dn-(p-1)}) \pmod{p^{a+2}} \quad \text{in } Q\bar{N}^{dn}.$$

However, if  $(\lambda, p) \neq 1$ , then by (5.3) together with Lemma 4.1 and Remark 4.2, it follows that  $\bar{R}_J^1(Q\bar{N}^{dn-(p-1)}) = \lambda \cdot \phi_0 \bar{R}_J^1(QN^{dn-(p-1)}) = 0 \pmod{p^{a+1}}$  and hence,  $p^{a+1} \cdot \bar{v}_n = 0 \pmod{p^{a+2}}$ . It is a contradiction and this completes the proof of Theorem B.

## REFERENCES

1. J. F. ADAMS, On Chern characters and the structure of the unitary group, *Proc. Cambridge Philos. Soc.* **57** (1961), 189–199.
2. J. F. ADAMS, On the groups  $J(X)$ . II, *Topology* **3** (1965), 137–172.
3. J. F. ADAMS, Lectures on generalized cohomology, Lecture 4, *Category Theory, Homology Theory and their Applications III* (Lecture Notes in Math. **99**, Springer, Berlin 1969), 77–113.
4. M. CURTIS and G. MISLIN, Two new  $H$ -spaces, *Bull. Amer. Math. Soc.* **4** (1970), 851–852.
5. D. L. GONÇALVES, *Mod 2 Homotopy-associative H-spaces, Geometric Application of Homotopy Theory I* (Lecture Notes in Math. **657** Springer, Berlin, 1978), 198–216.
6. G. H. HARDY and E. M. WRIGHT, *An Introduction to the Theory of Numbers* (Oxford University Press, London, 1954).
7. Y. HEMMI, Homotopy associative  $H$ -spaces and sphere extensions of classical groups, preprint.
8. P. HILTON and J. R. ROITBERG, On principal  $S^3$ -bundles over spheres, *Ann. of Math.* **90** (1969), 91–107.
9. J. R. HUBBUCK, Generalized cohomology operations and  $H$ -spaces of low rank, *Trans. Amer. Math. Soc.* **141** (1969), 335–360.
10. J. R. HUBBUCK, Primitivity in torsion free cohomology Hopf algebras, *Comment. Math. Helv.* **46** (1971), 13–43.
11. J. R. HUBBUCK, Some Pontrjagin rings, I, *Proc. Roy. Soc. Edinburgh Sect. A* **90** (1981), 237–256.
12. J. R. HUBBUCK and M. MIMURA, Certain  $p$ -regular  $H$ -spaces, *Arch. Math.* **49** (1987), 79–82.
13. N. IWASE, On the  $K$ -ring structure of  $X$ -projective  $n$ -space, *Mem. Fac. Sci. Kyushu Univ. Ser. A* **38** (1984), 285–297.
14. N. IWASE and M. MIMURA, Higher homotopy associativity, to appear in Arcata Proceedings.
15. F. SIGRIST and U. SUTER, Sur l'associativité homotopie des  $H$ -espaces de rang 2, *C. R. Acad. Sci. Paris* **238** (1971), 890–892.
16. F. SIGRIST and U. SUTER, Eine Anwendung der  $K$ -theorie in der  $H$ -Räume, *Comment. Math. Helv.* **47** (1972), 36–52.
17. J. D. STASHEFF, Homotopy associativity of  $H$ -spaces I and II, *Trans. Amer. Math. Soc.* **108** (1963), 275–292 and 293–312.
18. J. D. STASHEFF, Manifolds of the homotopy type of (non Lie) groups, *Bull. Amer. Math. Soc.* **75** (1969), 998–1000.
19. A. ZABRODSKY, On sphere extensions of classical Lie groups, *Transactions of the Summer Conference on Algebraic Topology* (Madison, Wisconsin, 1970).
20. A. ZABRODSKY, On spherical classes in the cohomology of  $H$ -spaces, *H-spaces Neuchâtel (Suisse) Aout 1970* (Lecture Notes in Math. **196** Springer, Berlin, 1971), 25–33.

21. A. ZABRODSKY, The classification of simply connected  $H$ -spaces with three cells I, *Math. Scand.* **30** (1972), 193–210.
22. A. ZABRODSKY, On the construction of new finite  $CW$ - $H$ -spaces, *Invent. Math.* **16** (1972), 260–266.

## CURRENT ADDRESS:

DEPARTMENT OF MATHEMATICS  
UNIVERSITY OF ABERDEEN  
THE EDWARD WRIGHT BUILDING  
ABERDEEN AB9 2TY  
U.K.

## PERMANENT ADDRESS:

DEPARTMENT OF MATHEMATICS  
OKAYAMA UNIVERSITY  
TSUSHIMA-NAKA OKAYAMA 700  
JAPAN