

ON THE CELLULAR DECOMPOSITION AND THE LUSTERNIK-SCHNIRELMANN CATEGORY OF $Spin(7)$

NORIO IWASE, MAMORU MIMURA AND TETSU NISHIMOTO

ABSTRACT. We give a cellular decomposition of the compact connected Lie group $Spin(7)$. We also determine the L-S categories of $Spin(7)$ and $Spin(8)$.

1. INTRODUCTION

In this paper, we assume that a space has the homotopy type of a CW-complex. The Lusternik-Schnirelmann category $\text{cat } X$ of a space X is the least integer n such that X is the union of $(n + 1)$ open subsets, each of which is contractible in X . G. Whitehead [15] showed that $\text{cat } X \leq n$ if and only if the diagonal map $\Delta_{n+1} : X \rightarrow \prod^{n+1} X$ is homotopic to some composition map

$$X \longrightarrow T^{n+1}(X) \longrightarrow \prod^{n+1} X,$$

where $T^{n+1}(X)$ is the fat wedge and $T^{n+1}(X) \rightarrow \prod^{n+1} X$ is the inclusion map.

The weak Lusternik-Schnirelmann category $\text{wcat } X$ is the least integer n such that the reduced diagonal map $\bar{\Delta}_{n+1} : X \rightarrow \wedge^{n+1} X$ is trivial. Then it is easy to see that $\text{wcat } X \leq \text{cat } X$, since $\wedge^{n+1} X = \prod^{n+1} X / T^{n+1}(X)$.

The strong Lusternik-Schnirelmann category $\text{Cat } X$ is the least integer n such that there exist a space X' which is homotopy equivalent to X and is covered by $(n + 1)$ open subsets contractible in themselves. $\text{Cat } X$ is closely related with $\text{cat } X$, and Ganea and Takens [14] showed that

$$\text{cat } X \leq \text{Cat } X \leq \text{cat } X + 1.$$

Ganea [3] showed that $\text{Cat } X$ is equal to the invariant which is the least integer n such that there is a cofibre sequence

$$A_i \longrightarrow X_{i-1} \longrightarrow X_i$$

where X_0 is a point and X_n is homotopy equivalent to X .

The Lusternik-Schnirelmann category for some Lie groups are determined, such as $\text{cat}(U(n)) = n$ and $\text{cat}(SU(n)) = n - 1$ by Singhof [11], $\text{cat}(Sp(2)) = 3$ by Schweitzer [10], $\text{cat}(Sp(3)) = 5$ by Fernández-Suárez, Gómez-Tato, Strom and Tanré [2], and Iwase and Mimura [6], $\text{cat}(SO(2)) = 1$, $\text{cat}(SO(3)) = 3$, $\text{cat}(SO(4)) =$

2000 *Mathematics Subject Classification.* Primary 55M30, Secondary 22E20, 57N60.

Key words and phrases. Lusternik-Schnirelmann category, Lie group, cellular decomposition.

4, $\text{cat}(SO(5)) = 8$ by James and Singhof [7]. Some general argument about the Lusternik-Schnirelmann category implies that $\text{cat}(G_2) = 4$ (see for example [6]).

As is well-known, we have the following isomorphisms:

$$\text{Spin}(3) \cong S^3, \quad \text{Spin}(4) \cong S^3 \times S^3, \quad \text{Spin}(5) \cong Sp(2), \quad \text{Spin}(6) \cong SU(4).$$

Thus $\text{Spin}(7)$ is the first non-trivial case in determining the cellular decomposition and the Lusternik-Schnirelmann category as well; it is our purpose in this paper.

Theorem 1.1. *We have $w\text{cat}(\text{Spin}(7)) = \text{cat}(\text{Spin}(7)) = \text{Cat}(\text{Spin}(7)) = 5$.*

Since $\text{Spin}(8)$ is homeomorphic to $\text{Spin}(7) \times S^7$, we obtain the following corollary.

Corollary 1.2. *We have $w\text{cat}(\text{Spin}(8)) = \text{cat}(\text{Spin}(8)) = \text{Cat}(\text{Spin}(8)) = 6$.*

The paper is organized as follows. In Section 2 we give a cellular decomposition of $\text{Spin}(7)$ such that $\text{Spin}(7)$ contains a subgroup $SU(4)$, which turns out to be useful for determining the Lusternik-Schnirelmann category of $\text{Spin}(7)$. In Section 3 we give a cone-decomposition of $SU(4)$, which gives rise to the Lusternik-Schnirelmann category of $\text{Spin}(7)$ in Section 4.

2. THE CELLULAR DECOMPOSITION OF $\text{Spin}(7)$

In this section, we use the notation in [9]. Let \mathfrak{C} be the Cayley algebra. $SO(8)$ acts on \mathfrak{C} naturally since $\mathfrak{C} \cong \mathbb{R}^8$ as \mathbb{R} -module. We regard $SO(7)$ as the subgroup of $SO(8)$ fixing e_0 , the unit of \mathfrak{C} . As is well known, the exceptional Lie group G_2 is defined by

$$G_2 = \{g \in SO(7) \mid g(x)g(y) = g(xy), x, y \in \mathfrak{C}\} = \text{Aut}(\mathfrak{C}).$$

According to [19], the group $\text{Spin}(7)$ is the set of the elements $\tilde{g} \in SO(8)$ such that $g(x)\tilde{g}(y) = \tilde{g}(xy)$ for any $x, y \in \mathfrak{C}$, where $g \in SO(7)$ is uniquely determined by \tilde{g} :

$$\text{Spin}(7) = \{\tilde{g} \in SO(8) \mid g(x)\tilde{g}(y) = \tilde{g}(xy), g \in SO(7), x, y \in \mathfrak{C}\}.$$

It is easy to see that G_2 is the subgroup of $\text{Spin}(7)$. Observe that the algebra generated by e_1 in \mathfrak{C} is isomorphic to \mathbb{C} . $SU(4)$ acts on \mathfrak{C} naturally, since as \mathbb{C} -module $\mathfrak{C} \cong \mathbb{C}^4$ whose basis is $\{e_0, e_2, e_4, e_6\}$. We regard $SU(3)$ as the subgroup of $SU(4)$ fixing e_0 and also as the subgroup of G_2 fixing e_1 .

Let D^i be the i -dimensional disc. We define four maps:

$$A : D^3 = \{(x_1, x_2, x_3) \in \mathbb{R}^3 \mid x_1^2 + x_2^2 + x_3^2 \leq 1\} \longrightarrow SO(8),$$

$$B : D^2 = \{(y_1, y_2) \in \mathbb{R}^2 \mid y_1^2 + y_2^2 \leq 1\} \longrightarrow SO(8),$$

$$C : D^1 = \{z_1 \in \mathbb{R} \mid z_1^2 \leq 1\} \longrightarrow SO(8),$$

$$D : D^2 = \{(w_1, w_2) \in \mathbb{R}^2 \mid w_1^2 + w_2^2 \leq 1\} \longrightarrow SO(8),$$

Proof. We express the map $(p_0\varphi_7)|_{V^7\setminus\partial V^7}$ as follows:

$$\begin{pmatrix} a_0 \\ a_1 \\ a_2 \\ a_3 \\ a_4 \\ a_5 \\ a_6 \\ a_7 \end{pmatrix} = D(\mathbf{w})B(\mathbf{y})A(\mathbf{x})B(\mathbf{y})^{-1}D(\mathbf{w})^{-1}e_0 = \begin{pmatrix} 1 - 2X^2Y^2W^2 \\ 2x_1XY^2W^2 \\ 2(w_1X - x_1w_2)XY^2W \\ -2(w_2X + x_1w_1)XY^2W \\ 2(-y_1X + x_1y_2)XYW \\ 2(y_2X + x_1y_1)XYW \\ 2x_2XYW \\ 2x_3XYW \end{pmatrix}$$

and hence we have

$$\begin{pmatrix} 1 - a_0 \\ a_1 \\ a_2 \\ a_3 \\ a_4 \\ a_5 \\ a_6 \\ a_7 \end{pmatrix} = 2XYW \begin{pmatrix} XYW \\ x_1YW \\ (w_1X - x_1w_2)Y \\ -(w_2X + x_1w_1)Y \\ -y_1X + x_1y_2 \\ y_2X + x_1y_1 \\ x_2 \\ x_3 \end{pmatrix}.$$

Since $X > 0$, $Y > 0$, $W > 0$ and $1 - a_0 > 0$, an easy calculation as for the first component in the above equation gives the following equation:

$$(2.1) \quad XYW = \frac{\sqrt{1 - a_2}}{\sqrt{2}},$$

from which we easily obtain

$$(2.2) \quad x_2 = \frac{a_6}{\sqrt{2(1 - a_2)}}, \quad x_3 = \frac{a_7}{\sqrt{2(1 - a_2)}}.$$

Further we obtain three more equalities from the above equalities:

$$\begin{aligned} (1 - a_0)^2 + a_1^2 &= 4X^2Y^4W^4(x_1^2 + X^2), \\ a_2^2 + a_3^2 &= 4X^2Y^4W^2(w_1^2 + w_2^2)(x_1^2 + X^2) = 4X^2Y^4W^2(1 - W^2)(x_1^2 + X^2), \\ a_4^2 + a_5^2 &= 4X^2Y^2W^2(y_1^2 + y_2^2)(x_1^2 + X^2) = 4X^2Y^2W^2(1 - Y^2)(x_1^2 + X^2). \end{aligned}$$

Using these three equalities, we obtain

$$(2.3) \quad Y^2 = \frac{(1 - a_0)^2 + a_1^2 + a_2^2 + a_3^2}{(1 - a_0)^2 + a_1^2 + a_2^2 + a_3^2 + a_4^2 + a_5^2},$$

$$(2.4) \quad W^2 = \frac{(1 - a_0)^2 + a_1^2}{(1 - a_0)^2 + a_1^2 + a_2^2 + a_3^2}.$$

It follows from (2.1), (2.3) and (2.4) that

$$(2.5) \quad X^2 = \frac{(1 - a_0)((1 - a_0)^2 + a_1^2 + a_2^2 + a_3^2 + a_4^2 + a_5^2)}{2((1 - a_0)^2 + a_1^2)}.$$

It follows also from (2.2) and (2.5) that

$$(2.6) \quad x_1^2 = \frac{a_1^2((1-a_0)^2 + a_1^2 + a_2^2 + a_3^2 + a_4^2 + a_5^2)}{2(1-a_0)((1-a_0)^2 + a_1^2)}.$$

Since Y, W, X are positive, (2.3), (2.4), (2.5) imply respectively

$$(2.7) \quad Y = \frac{\sqrt{(1-a_0)^2 + a_1^2 + a_2^2 + a_3^2}}{\sqrt{(1-a_0)^2 + a_1^2 + a_2^2 + a_3^2 + a_4^2 + a_5^2}},$$

$$(2.8) \quad W = \frac{\sqrt{(1-a_0)^2 + a_1^2}}{\sqrt{(1-a_0)^2 + a_1^2 + a_2^2 + a_3^2}},$$

$$(2.9) \quad X = \frac{\sqrt{(1-a_0)((1-a_0)^2 + a_1^2 + a_2^2 + a_3^2 + a_4^2 + a_5^2)}}{\sqrt{2((1-a_0)^2 + a_1^2)}}.$$

Since the signs of x_1 and a_1 are the same, (2.6) implies that

$$(2.10) \quad x_1 = \frac{a_1 \sqrt{(1-a_0)^2 + a_1^2 + a_2^2 + a_3^2 + a_4^2 + a_5^2}}{\sqrt{2(1-a_0)((1-a_0)^2 + a_1^2)}}.$$

Now we determine y_1 ; we have

$$-a_4 X + a_5 x_1 = 2XYW(x_1^2 + X^2)y_2.$$

Substituting the equations (2.1), (2.9) and (2.10) in the above equation, we obtain

$$(2.11) \quad y_1 = \frac{a_1 a_5 - (1-a_0)a_4}{\sqrt{((1-a_0)^2 + a_1^2)((1-a_0)^2 + a_1^2 + a_2^2 + a_3^2 + a_4^2 + a_5^2)}}.$$

We determine y_2 ; we have

$$a_4 x_1 + a_5 X = 2XYW(x_1^2 + X^2)y_2.$$

Substituting the equations (2.1), (2.9) and (2.10) in the above equation, we obtain

$$(2.12) \quad y_2 = \frac{a_1 a_4 + (1-a_0)a_5}{\sqrt{((1-a_0)^2 + a_1^2)((1-a_0)^2 + a_1^2 + a_2^2 + a_3^2 + a_4^2 + a_5^2)}}.$$

We determine w_1 ; we have

$$a_2 X - a_3 x_1 = 2XY^2W(x_1^2 + X^2)w_1.$$

Substituting the equations (2.1), (2.7), (2.9) and (2.10) in the above equation, we obtain

$$(2.13) \quad w_1 = \frac{(1-a_0)a_2 - a_1 a_3}{\sqrt{((1-a_0)^2 + a_1^2)((1-a_0)^2 + a_1^2 + a_2^2 + a_3^2)}}.$$

Finally we determine w_2 ; we have

$$-a_2 x_1 - a_3 X = 2XY^2W(x_1^2 + X^2)w_2.$$

Substituting the equations (2.1), (2.7), (2.9) and (2.10) in the above equation, we obtain

$$(2.14) \quad w_2 = \frac{-a_1 a_2 - (1-a_0)a_3}{\sqrt{((1-a_0)^2 + a_1^2)((1-a_0)^2 + a_1^2 + a_2^2 + a_3^2)}}.$$

Thus we have expressed $x_1, x_2, x_3, y_1, y_2, w_1, w_2$ in terms of a_0, \dots, a_7 , that is, the inverse map has been constructed, which completes the proof. \square

In a similar way to that of Section 3 of [9], we can obtain the following theorem, which is essentially the same as Yokota's decomposition [17].

Proposition 2.3. $e^0 \cup e^3 \cup e^5 \cup e^7 \cup e^8 \cup e^{10} \cup e^{12} \cup e^{15}$ thus obtained is a cellular decomposition of $SU(4)$.

Proof. First we show that $\dot{e}^i \cap \dot{e}^j = \emptyset$ if $i \neq j$. We consider the following three cases:

(1) For the case where $i, j \in \{0, 3, 5, 8\}$; both cells e^i and e^j are in $SU(3)$ and $e^0 \cup e^3 \cup e^5 \cup e^8$ is a cellular decomposition of $SU(3)$. Then we have $\dot{e}^i \cap \dot{e}^j = \emptyset$ if $i \neq j$.

(2) For the case where $i \in \{0, 3, 5, 8\}$ and $j \in \{7, 10, 12, 15\}$; we have $p_0(\dot{e}^i) = \{e_0\}$ and $p_0(\dot{e}^j) = S^7 \setminus \{e_0\}$. Then we have $\dot{e}^i \cap \dot{e}^j = \emptyset$.

(3) For the case where $i, j \in \{7, 10, 12, 15\}$; suppose that $A \in \dot{e}^i \cap \dot{e}^j$. Since $\dot{e}^i = \dot{e}^7 \dot{e}^{i-7}$ and $\dot{e}^j = \dot{e}^7 \dot{e}^{j-7}$, we can put $A = A_1 A_2 = A'_1 A'_2$ where $A_1, A'_1 \in \dot{e}^7$, $A_2 \in \dot{e}^{i-7}$ and $A'_2 \in \dot{e}^{j-7}$. We have $A_1 = A'_1$, since $p_0(A_1) = p_0(A_1 A_2) = p_0(A'_1 A'_2) = p_0(A'_1)$ and $p_0|_{\dot{e}^7}$ is monic. Then we have $A_2 = A'_2$ and the first case shows that $i - 7 = j - 7$, that is, $i = j$. Thus $\dot{e}^i \cap \dot{e}^j = \emptyset$ if $i \neq j$.

Next, we will check that the boundaries of the cells are included in the lower dimensional cells. In the proof of Proposition 3.2 [9], it is proved that the boundaries \dot{e}^3, \dot{e}^5 and \dot{e}^8 are included in the lower dimensional cells. Observe that the boundary \dot{e}^7 is the union of the following three sets:

$$\begin{aligned} & \{DBAB^{-1}D^{-1} \mid A \in A(\dot{D}^3), B \in B(D^2), D \in D(D^2)\}, \\ & \{DBAB^{-1}D^{-1} \mid A \in A(D^3), B \in B(\dot{D}^2), D \in D(D^2)\}, \\ & \{DBAB^{-1}D^{-1} \mid A \in A(D^3), B \in B(D^2), D \in D(\dot{D}^2)\}. \end{aligned}$$

The first set contains only the identity element, since A is the identity element. It is easy to see that the second set is contained in e^3 and that the third set is contained in e^5 . We have $\dot{e}^{10} = e^7 \dot{e}^3 \cup \dot{e}^7 e^3 \subset e^7 e^0 \cup e^5 e^3 = e^7 \cup e^8$. We also have $\dot{e}^{12} = \dot{e}^7 e^5 \cup e^7 \dot{e}^5 \subset e^5 e^5 \cup e^7 e^3 = e^8 \cup e^{10}$, and $\dot{e}^{15} = \dot{e}^7 e^5 e^3 \cup e^7 \dot{e}^5 e^3 \cup e^7 e^5 \dot{e}^3 \subset e^5 e^5 e^3 \cup e^7 e^3 e^3 \cup e^7 e^5 = e^8 \cup e^{10} \cup e^{12}$.

Finally, we will show that the inclusion map $e^0 \cup e^3 \cup e^5 \cup e^7 \cup e^8 \cup e^{10} \cup e^{12} \cup e^{15} \rightarrow SU(4)$ is epic. Let $g \in SU(4)$. If $p_0(g) = e_0$, then g is contained in $SU(3) = e^0 \cup e^3 \cup e^5 \cup e^8$. Suppose that $p_0(g) \neq e_0$. There is an element $h \in e^7$ such that $p_0(h) = p_0(g)$. Thus we have $h^{-1}g \in SU(3) = e^0 \cup e^3 \cup e^5 \cup e^8$, since $p_0(h^{-1}g) = e_0$. Therefore we have $g \in h(e^0 \cup e^3 \cup e^5 \cup e^8) \subset e^0 \cup e^3 \cup e^5 \cup e^7 \cup e^8 \cup e^{10} \cup e^{12} \cup e^{15}$. \square

Remark 2.4. (1) We regard $SO(6)$ as the subgroup of $SO(7)$ fixing e_1 . Let $\pi : Spin(6) \rightarrow SO(6)$ be the double covering. Then, according to the Proof of Lemma 2.1, $\pi(SU(4)) \subset SO(6)$ so that $\pi|_{SU(4)} : SU(4) \rightarrow SO(6)$ is the double covering.

(2) For $1 \leq n \leq 3$, the subcomplex $e^0 \cup e^3 \cup \dots \cup e^{2n+1}$ is homeomorphic to ΣCP^n , which consists of the elements

$$A \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & & & e^{2i\theta} \end{pmatrix} A^{-1} \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & 1 & \\ & & & e^{-2i\theta} \end{pmatrix}$$

for any elements A in $SU(n+1)$. Moreover, according to Proposition 2.6 of Chapter IV of [13], we have $e^{2i+1}e^{2j+1} \subset e^{2j+1}e^{2i+1}$ for $i < j$; in fact we have $e^{2i+1}e^{2j+1} = e^{2j+1}e^{2i+1}$ (see [19]).

Let S^6 be the unit sphere of \mathbb{R}^7 whose basis $\{e_i \mid 1 \leq i \leq 7\}$. We consider the following diagram

$$\begin{array}{ccccc} SU(3) & \longrightarrow & G_2 & \longrightarrow & S^6 \\ \downarrow & & \downarrow & & \parallel \\ SU(4) & \longrightarrow & Spin(7) & \xrightarrow{p} & S^6 \\ \downarrow & & \pi \downarrow & & \parallel \\ SO(6) & \longrightarrow & SO(7) & \longrightarrow & S^6 \end{array}$$

where the horizontal lines are principal fibre bundles and $p(g) = \pi(g)e_1$.

Lemma 4.1 of [9] implies the following lemma immediately.

Lemma 2.5. *Put $V^6 = D^3 \times D^2 \times D^1$. Then the composite map $p\varphi_6 : (V^6, \partial V^6) \rightarrow (S^6, \{e_1\})$ is a relative homeomorphism.*

Now we can state one of our main results.

Theorem 2.6. *The cell complex $e^0 \cup e^3 \cup e^5 \cup e^6 \cup e^7 \cup e^8 \cup e^9 \cup e^{10} \cup e^{11} \cup e^{12} \cup e^{13} \cup e^{14} \cup e^{15} \cup e^{16} \cup e^{18} \cup e^{21}$ gives a cellular decomposition of $Spin(7)$.*

Proof. First we show that $\tilde{e}^i \cap \tilde{e}^j = \emptyset$ if $i \neq j$. We consider the following three cases:

(1) For the case where $i, j \in \{0, 3, 5, 7, 8, 10, 12, 15\}$; both cells e^i and e^j are in $SU(4)$ and $e^0 \cup e^3 \cup e^5 \cup e^7 \cup e^8 \cup e^{10} \cup e^{12} \cup e^{15}$ is a cellular decomposition of $SU(4)$, whence we have $\tilde{e}^i \cap \tilde{e}^j = \emptyset$ if $i \neq j$.

(2) For the case where $i \in \{0, 3, 5, 7, 8, 10, 12, 15\}$ and $j \in \{6, 9, 11, 13, 14, 16, 18, 21\}$; we have $p(\tilde{e}^i) = \{e_1\}$ and $p(\tilde{e}^j) = S^6 \setminus \{e_1\}$, whence we have $\tilde{e}^i \cap \tilde{e}^j = \emptyset$.

(3) For the case where $i, j \in \{6, 9, 11, 13, 14, 16, 18, 21\}$, suppose that $A \in \tilde{e}^i \cap \tilde{e}^j$. Since $\tilde{e}^i = \tilde{e}^6 \tilde{e}^{i-6}$ and $\tilde{e}^j = \tilde{e}^6 \tilde{e}^{j-6}$, we can put $A = A_1 A_2 = A'_1 A'_2$, where $A_1, A'_1 \in \tilde{e}^6$, $A_2 \in \tilde{e}^{i-6}$ and $A'_2 \in \tilde{e}^{j-6}$. We have $A_1 = A'_1$, since $p(A_1) = p(A_1 A_2) = p(A'_1 A'_2) = p(A'_1)$ and $p|_{\tilde{e}^6}$ is monic. Then we have $A_2 = A'_2$ and the first case shows that $i - 6 = j - 6$, that is, $i = j$. Thus $\tilde{e}^i \cap \tilde{e}^j = \emptyset$ if $i \neq j$.

Next, we will check that the boundaries of the cells are included in the lower dimensional cells. In Proposition 2.3, it is proved that the boundaries of the cells of $SU(4)$ are included in the lower dimensional cells. In Proof of Theorem 4.2 in [9], we showed that $\dot{e}^6 \subset e^3 \cup e^5$, $\dot{e}^9 \subset e^6 \cup e^8$, $\dot{e}^{11} \subset e^5 \cup e^9$ and $\dot{e}^{14} \subset e^8 \cup e^9 \cup e^{11}$. By using (2) of Remark 2.4, we also obtain

$$\begin{aligned} \dot{e}^{13} &= e^6 \dot{e}^7 \cup \dot{e}^6 e^7 \subset e^{11} \cup e^{12}, \\ \dot{e}^{16} &= e^6 e^7 \dot{e}^3 \cup e^6 \dot{e}^7 e^3 \cup \dot{e}^6 e^7 e^3 \subset e^{13} \cup e^{14} \cup e^{15}, \\ \dot{e}^{18} &= e^6 e^7 \dot{e}^5 \cup e^6 \dot{e}^7 e^5 \cup \dot{e}^6 e^7 e^5 \subset e^{16} \cup e^{14} \cup e^{15}, \\ \dot{e}^{21} &= e^6 e^7 e^5 \dot{e}^3 \cup e^6 e^7 \dot{e}^5 e^3 \cup e^6 \dot{e}^7 e^5 e^3 \cup \dot{e}^6 e^7 e^5 e^3 \subset e^{18} \cup e^{16} \cup e^{14} \cup e^{15}. \end{aligned}$$

Finally, we will show that the inclusion map $e^0 \cup e^3 \cup e^5 \cup e^6 \cup e^7 \cup e^8 \cup e^9 \cup e^{10} \cup e^{11} \cup e^{12} \cup e^{13} \cup e^{14} \cup e^{15} \cup e^{16} \cup e^{18} \cup e^{21} \rightarrow Spin(7)$ is epic. Let $g \in Spin(7)$. If $p(g) = e_1$, then g is contained in $SU(4) = e^0 \cup e^3 \cup e^5 \cup e^7 \cup e^8 \cup e^{10} \cup e^{12} \cup e^{15}$. Suppose that $p(g) \neq e_1$. There is an element $h \in e^6$ such that $p(h) = p(g)$. Thus we have $h^{-1}g \in SU(4)$ since $p(h^{-1}g) = e_1$. Therefore we have $g \in h(e^0 \cup e^3 \cup e^5 \cup e^7 \cup e^8 \cup e^{10} \cup e^{12} \cup e^{15}) \subset e^0 \cup e^3 \cup e^5 \cup e^6 \cup e^7 \cup e^8 \cup e^9 \cup e^{10} \cup e^{11} \cup e^{12} \cup e^{13} \cup e^{14} \cup e^{15} \cup e^{16} \cup e^{18} \cup e^{21}$. \square

Remark 2.7. Araki [1] also gave a cellular decomposition of $Spin(n)$, but the one we have given here is a cellular decomposition with the minimum number of cells, satisfying the Yokota principle ([17], [18], [19]). As will be seen later, it is effectively used to determine the Lusternik-Schnirelmann category.

It is easy to give a cellular decomposition of $Spin(8)$ using a homeomorphism $Spin(8) \rightarrow Spin(7) \times S^7$.

3. THE CONE-DECOMPOSITION OF $SU(4)$

Obviously there is a filtration $F'_0 = * \subset F'_1 = SU(4)^{(7)} \subset F'_2 = SU(4)^{(12)} \subset F'_3 = SU(4)$. It is well-known that $F'_1 = \Sigma CP^3 = S^3 \cup e^5 \cup e^7$ and $F'_2 = F'_1 \cup e^8 \cup e^{10} \cup e^{12}$. Thus the integral cohomology $H^n(F'_2; \mathbb{Z})$ is given by

$$H^n(F'_2; \mathbb{Z}) \cong \begin{cases} \mathbb{Z}\langle 1 \rangle & (n = 0) \\ \mathbb{Z}\langle y_n \rangle & (n = 3, 5, 7, 8, 10, 12) \\ 0 & (\text{otherwise}). \end{cases}$$

The action of the squaring operation Sq^2 is given as follows:

$$Sq^2 y_n = \begin{cases} y_{n+2} & \text{for } n = 3, 10, \\ 0 & \text{for } n = 5, 7, 8, 12 \end{cases}$$

where y_n is regarded as an element of the mod 2 cohomology. To give the cone decomposition of $SU(4)$, we use the following homotopy fibration:

$$(3.1) \quad F \xrightarrow{\Psi} F'_1 \xrightarrow{\iota} F'_2.$$

Without loss of generality, we may regard this as a Hurewicz fibration over F'_2 .

Firstly we consider the Serre spectral sequence $(E_r^{*,*}, d_r)$ associated with the above fibration, where the generators of $E_2^{*,0}$ for $* \leq 7$ are permanent cycles and survive to E_∞ -terms. Hence F is 6-connected and the transgression $\tau : H^7(F; \mathbb{Z}) \rightarrow H^8(F'_2; \mathbb{Z})$ is an isomorphism to $H^8(F'_2; \mathbb{Z}) \cong \mathbb{Z}\langle y_8 \rangle$. Thus $H^7(F; \mathbb{Z}) \cong \mathbb{Z}\langle x_7 \rangle$ for some $x_7 \in H^7(F; \mathbb{Z})$. Similarly, the generators in $E_2^{3,7} \cong \mathbb{Z}\langle y_3 \otimes x_7 \rangle$ and $E_2^{10,0} \cong H^{10}(F'_2; \mathbb{Z}) \cong \mathbb{Z}\langle y_{10} \rangle$ must lie in the image of differentials d_3 and $d_{10} = \tau : H^9(F; \mathbb{Z}) \rightarrow H^{10}(F'_2; \mathbb{Z})$ respectively, and we have that $H^8(F; \mathbb{Z}) = 0$ and $H^9(F; \mathbb{Z}) \cong \mathbb{Z}\langle x_9 \rangle \oplus \mathbb{Z}\langle x'_9 \rangle$, where the elements x_9 and x'_9 in $H^9(F; \mathbb{Z})$ are corresponding to x_{10} and $y_3 \otimes x_7$ by the transgression τ and d_3 respectively. We remark that the choice of the generator x'_9 is not unique. Continuing this process, we have that $H^{10}(F; \mathbb{Z}) = 0$ and $H^{11}(F; \mathbb{Z}) \cong \mathbb{Z}\langle x_{11} \rangle \oplus \mathbb{Z}\langle x'_{11} \rangle \oplus \mathbb{Z}\langle x''_{11} \rangle \oplus \mathbb{Z}\langle x'''_{11} \rangle$ whose generators are corresponding to x_{12} , $y_3 \otimes x_9$, $y_3 \otimes x'_9$ and $y_5 \otimes x_7$ respectively by the transgression τ and differentials d_3 , d_3 and d_5 .

Thus the integral cohomology $H^n(F; \mathbb{Z})$ for $0 \leq n \leq 11$ is given by

$$H^n(F; \mathbb{Z}) \cong \begin{cases} \mathbb{Z}\langle 1 \rangle & (n = 0) \\ \mathbb{Z}\langle x_7 \rangle & (n = 7) \\ \mathbb{Z}\langle x_9 \rangle \oplus \mathbb{Z}\langle x'_9 \rangle & (n = 9) \\ \mathbb{Z}\langle x_{11} \rangle \oplus \mathbb{Z}\langle x'_{11} \rangle \oplus \mathbb{Z}\langle x''_{11} \rangle \oplus \mathbb{Z}\langle x'''_{11} \rangle & (n = 11) \\ 0 & (\text{otherwise}) \end{cases}$$

where x_7 , x_9 and x_{11} are transgressive generators in $H^*(F; \mathbb{Z})$. Hence F has, up to homotopy, a cellular decomposition $e^0 \cup e^7 \cup_{\varphi_1} e^9 \cup_{\varphi'_1} e^9 \cup_{\varphi_2} e^{11} \cup$ (cells in dimensions ≥ 11), where the cells e^7 , e^9 and e^{11} correspond to x_7 , x_9 and x_{11} respectively. Then we obtain a subcomplex $A' = e^0 \cup e^7 \cup_{\varphi_1} e^9 \cup_{\varphi'_1} e^9 \cup_{\varphi_2} e^{11}$ of F .

Secondly, we determine the attaching maps φ_1 and φ'_1 : Let us recall that $\pi_8(S^7) \cong \mathbb{Z}/2\langle \eta_7 \rangle$ whose generator η_7 can be detected by Sq^2 , the mod 2 Steenrod operation. Since the action of mod 2 Steenrod operation commutes with the cohomology transgression (see [8, Proposition 6.5]), we see that $Sq^2 x_7$ is transgressive, and hence is cx_9 for some $c \in \mathbb{Z}/2$. We know that $\tau x_9 = y_{10} \neq 0$ and $\tau Sq^2 x_7 = Sq^2 \tau x_7 = Sq^2 y_8 = 0$, and hence $Sq^2 x_7$ must be trivial. Thus the attaching maps φ_1 and φ'_1 are both null homotopic and A' is homotopy equivalent to $(S^7 \vee S^9 \vee S^9_1) \cup_{\varphi_2} e^{11}$.

Thirdly we check the composition of projections with the attaching map $\varphi_2 : S^{10} \rightarrow S^7 \vee S^9 \vee S^9_1$ to S^9 and S^9_1 , which can also be detected by Sq^2 . Again by the commutativity of the action of mod 2 Steenrod operation with the transgression, we see that the composition map $\text{pr}_{S^9} \circ \varphi_2 : S^{10} \xrightarrow{\varphi_2} S^7 \vee S^9 \vee S^9_1 \rightarrow S^9$ represents a generator of $\pi_{10}(S^9) \cong \mathbb{Z}/2\langle \eta_9 \rangle$, since $Sq^2 : H^8(F'_2; \mathbb{Z}/2) \rightarrow H^{10}(F'_2; \mathbb{Z}/2)$ is non-trivial. If the composition map $\phi_1 = \text{pr}_{S^9_1} \circ \varphi_2 : S^{10} \xrightarrow{\varphi_2} S^7 \vee S^9 \vee S^9_1 \rightarrow S^9_1$ is non-trivial, we replace φ_2 by the composition of φ_2 and the homotopy equivalence

$\xi : S^7 \vee S^9 \vee S_1^9 \rightarrow S^7 \vee S^9 \vee S_1^9$ where $\xi|_{S^7}$ and $\xi|_{S_1^9}$ are the identity maps and $\xi|_{S^9}$ is the unique co-H-structure map $\phi : S^9 \rightarrow S^9 \vee S_1^9$; then we obtain that ϕ_1 is trivial, since $2\eta_9 = 0$. Then A' is homotopy equivalent to $((S^7 \vee S^9) \cup_{\varphi_2} e^{11}) \vee S_1^9$. Let A denote the subcomplex $(S^7 \vee S^9) \cup_{\varphi_2} e^{11}$ of A' and $\psi = \Psi|_A : A \rightarrow F'_1$.

Lemma 3.1. F'_2 is homotopy equivalent to $F'_1 \cup_{\psi} CA$.

Proof. The image of $H^*(A; \mathbb{Z})$ in $H^*(F; \mathbb{Z})$ under the induced map of the inclusion coincides with the module of transgressive elements with respect to the fibration (3.1) (see [8, Chapter 6]). Thus we may regard that $H^{n-1}(A; \mathbb{Z}) = \delta^{-1}(\iota^*(H^n(F'_2, *))) \subset H^{n-1}(F; \mathbb{Z})$:

$$\begin{array}{ccccc} H^{n-1}(F; \mathbb{Z}) & \xrightarrow{\delta_F} & H^n(F'_1, F; \mathbb{Z}) & \xleftarrow{\iota_F^*} & H^n(F'_2, *; \mathbb{Z}) \\ \downarrow & & \downarrow & & \parallel \downarrow \\ H^{n-1}(A; \mathbb{Z}) & \xrightarrow{\delta_A} & H^n(F'_1, A; \mathbb{Z}) & \xleftarrow{\iota_A^*} & H^n(F'_2, *; \mathbb{Z}), \end{array}$$

where ι_F and ι_A are given by ι , and δ_F and δ_A denote the connecting homomorphisms of the long exact sequences for the pairs (F'_1, F) and (F'_1, A) , respectively. Thus the image of δ_A is contained in the image of ι_A^* and we also have

$$H^n(F'_1, A; \mathbb{Z}) \cong H^n(F'_1 \cup_{\psi} CA, CA; \mathbb{Z}) \cong H^n(F'_1 \cup_{\psi} CA, *; \mathbb{Z}).$$

Since the composition map $A \xrightarrow{\psi} F'_1 \xrightarrow{\iota} F'_2$ is trivial, we can define a map

$$f : F'_1 \cup_{\psi} CA \longrightarrow F'_2,$$

by $f|_{F'_1} = \iota : F'_1 \rightarrow F'_2$ and $f|_{CA} = *$.

In order to prove the lemma, we show that $f^* : H^n(F'_2; \mathbb{Z}) \cong \mathbb{Z} \rightarrow H^n(F'_1 \cup_{\psi} CA; \mathbb{Z}) \cong \mathbb{Z}$ is an isomorphism for $n = 3, 5, 7, 8, 10, 12$. We have a commutative diagram

$$\begin{array}{ccccc} H^n(F'_2; \mathbb{Z}) & & \xrightarrow{\iota^*} & & H^n(F'_1; \mathbb{Z}) \\ f^* \downarrow & & & & \parallel \downarrow \\ H^n(F'_1 \cup CA, F'_1; \mathbb{Z}) & \xrightarrow{j^*} & H^n(F'_1 \cup CA; \mathbb{Z}) & \xrightarrow{i^*} & H^n(F'_1; \mathbb{Z}), \end{array}$$

where the bottom row is a part of the exact sequence for the pair $(F'_1 \cup CA, F'_1)$. The induced map i^* is an isomorphism for $n \leq 7$, since $H^n(F'_1 \cup CA, F'_1; \mathbb{Z}) = 0$ for $n \leq 7$ and since ι^* is an isomorphism for $n \leq 7$. Then we obtain that f^* is an isomorphism for $n \leq 7$. Moreover we can show that $j^* : H^n(F'_1 \cup CA, F'_1; \mathbb{Z}) \rightarrow H^n(F'_1 \cup CA; \mathbb{Z})$ is an isomorphism for $n \geq 8$, by considering the exact sequence for the pair $(F'_1 \cup CA, F'_1)$, since we have $H^n(F'_1) = 0$ for $n \geq 8$. To perform the other cases for $n = 8, 10, 12$, it is sufficient to show that f^* is surjective. In fact, we have a commutative diagram

$$\begin{array}{ccccc}
H^{n-1}(A; \mathbb{Z}) & \xrightarrow{\delta_A} & H^n(F'_1, A; \mathbb{Z}) & \xleftarrow{\iota_A^*} & H^n(F'_2, *, \mathbb{Z}) \\
\downarrow \Sigma \cong & & \uparrow & \searrow \cong & \downarrow f^* \\
H^n(\Sigma A, *, \mathbb{Z}) & \xrightarrow{\cong} & H^n(F'_1 \cup CA, F'_1; \mathbb{Z}) & \xrightarrow{j^*} & H^n(F'_1 \cup CA, *, \mathbb{Z}),
\end{array}$$

where Σ is the suspension isomorphism. Since j^* is an isomorphism for $n \geq 8$, we obtain that δ_A is an isomorphism for $n \geq 8$. Since the image of δ_A is contained in the image of ι_A^* , we see that f^* is surjective for $n \geq 8$, and hence f is a homotopy equivalence. \square

Proposition 3.2. *We have $wcat(F'_i) = cat(F'_i) = Cat(F'_i) = i$.*

Proof. The cohomology of F'_i implies that $wcat(F'_i) \geq i$. The cone-decomposition

$$F'_1 = \Sigma \mathbb{C}P^3, \quad F'_2 \simeq F'_1 \cup CA, \quad F'_3 = F'_2 \cup CS^{14}$$

implies that $Cat(F'_i) \leq i$, which completes the proof. \square

4. PROOF OF THEOREM 1.1

We define a filtration $F_0 = * \subset F_1 \subset F_2 \subset F_3 \subset F_4 \subset F_5 = Spin(7)$ by

$$\begin{aligned}
F_1 &= SU(4)^{(7)}, & F_2 &= SU(4)^{(12)} \cup e^6, \\
F_3 &= SU(4) \cup e^6 \cup e^9 \cup e^{11} \cup e^{13}, & F_4 &= Spin(7)^{(18)}.
\end{aligned}$$

We need the following lemma to prove Theorem 4.2.

Lemma 4.1. *We have a homeomorphism of pairs*

$$(CA_1, A_1) \times (CA_2, A_2) = (C(A_1 * A_2), A_1 * A_2).$$

(The proof can be found in p.482-483 of [16].)

Now Theorem 1.1 follows from the following theorem.

Theorem 4.2. *We have $wcat(F_i) = cat(F_i) = Cat(F_i) = i$.*

Proof. The mod 2 cohomology of F_i implies that $wcat(F_i) \geq i$. Then it is sufficient to show that $Cat(F_i) \leq i$. Obviously we have a homeomorphism $F_1 = \Sigma \mathbb{C}P^3$. Since the cell e^6 is attached to F_1 , we obtain that $F_2 \simeq F_1 \cup C(S^5 \vee A)$ using Lemma 3.1. Since we have $e^9 \cup e^{11} \cup e^{13} = e^6(e^3 \cup e^5 \cup e^7)$, the composition map

$$(CS^5, S^5) \times (C\mathbb{C}P^3, \mathbb{C}P^3) \longrightarrow (CS^5, S^5) \times (\Sigma \mathbb{C}P^3, *) \longrightarrow (F_2 \cup e^9 \cup e^{11} \cup e^{13}, F_2)$$

is a relative homeomorphism. Then we obtain $F_2 \cup e^9 \cup e^{11} \cup e^{13} = F_2 \cup C(S^5 * \mathbb{C}P^3)$ using Lemma 4.1. The cell e^{15} is the highest dimensional cell of $SU(4)$ and is attached to F_2 . Then we obtain $F_3 \simeq F_2 \cup C(S^{14} \vee (S^5 * \mathbb{C}P^3))$. Now we consider the following composition map:

$$(C(S^5 * A), S^5 * A) = (CS^5, S^5) \times (CA, A) \longrightarrow (CS^5, S^5) \times (F'_2, F'_1) \longrightarrow (F_4, F_3).$$

Since we have $e^{14} \cup e^{16} \cup e^{18} = e^6(e^8 \cup e^{10} \cup e^{12})$, the right map is a relative homeomorphism. The left map induces an isomorphism of homologies of pairs so that the map $H_*(F_3 \cup C(S^5 * A), F_3; \mathbb{Z}) \rightarrow H_*(F_4, F_3; \mathbb{Z})$ is an isomorphism. Thus we obtain $F_4 \simeq F_3 \cup C(S^5 * A)$. Obviously we have a homeomorphism $F_5 = F_4 \cup CS^{20}$. \square

REFERENCES

- [1] S. Araki, On the homology of spinor groups, Mem. Fac. Sci. Kyusyu Univ. Ser. A., **9**(1955), 1–35.
- [2] L. Fernández-Suárez, A. Gómez-Tato, J. Strom and D. Tanré, The Lusternik-Schnirelmann category of $Sp(3)$, (preprint).
- [3] T. Ganea, Lusternik-Schnirelmann category and strong category, Illinois J. Math., **11**(1967), 417–427.
- [4] N. Iwase, A_∞ -method in Lusternik-Schnirelmann category, Topology, **41**(2002), 695–723.
- [5] N. Iwase, Lusternik-Schnirelmann category of a sphere-bundle over a sphere, Topology, (to appear).
- [6] N. Iwase and M. Mimura, L-S categories of simply-connected compact simple Lie groups of low rank, Proc. of the Skye Conference, (to appear).
- [7] I. M. James and W. Singhof, On the category of fibre bundles, Lie groups, and Frobenius maps, Higher homotopy structures in topology and mathematical physics (Poughkeepsie, NY, 1996), Contemp. Math., **227**(1999), 177–189.
- [8] J. McCleary, A User’s Guide to Spectral Sequences, 2nd Ed., **58** Cambridge University Press, Cambridge, (2001).
- [9] M. Mimura and T. Nishimoto, On the cellular decomposition of the exceptional Lie group G_2 , Proc. Amer. Math. Soc., **130**(2002), 2451–2459.
- [10] P. A. Schweitzer, Secondary cohomology operations induced by the diagonal mapping, Topology, **3**(1965), 337–355.
- [11] W. Singhof, On the Lusternik-Schnirelmann category of Lie groups, Math. Z., **145**(1975), 111–116.
- [12] W. Singhof, On the Lusternik-Schnirelmann category of Lie groups, II, Math. Z., **151**(1976), 143–148.
- [13] N. E. Steenrod, Cohomology operations, Annals of Mathematical Studies, **50** Princeton University Press, Princeton, N.J. (1962).
- [14] F. Takens, The Lusternik-Schnirelman categories of a product space, Compositio Math., **22**(1970), 175–180.
- [15] G. W. Whitehead, The homology suspension, *Colloque de topologie algébrique, Louvain*, (1956), 89–95.
- [16] G. W. Whitehead, Elements of Homotopy Theory, Graduate Texts in Mathematics, **61** Springer-Verlag, New York-Berlin, (1978).
- [17] I. Yokota, On the cell structures of $SU(n)$ and $Sp(n)$, Proc. Japan Acad., **31**(1955), 673–677.
- [18] I. Yokota, On the cellular decompositions of unitary groups, J. Inst. Polytech. Osaka City Univ. Ser. A., **7**(1956), 39–49.
- [19] I. Yokota, Groups and Topology, Shokabo, (1971) (in Japanese).

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE, OKAYAMA UNIVERSITY, 3-1 TSUSHIMANA-NAKA, OKAYAMA 700-8530, JAPAN

E-mail address: `mimura@math.okayama-u.ac.jp`

DEPARTMENT OF WELFARE BUSINESS, KINKI WELFARE UNIVERSITY, FUKUSAKI-CHO, HYOGO 679-2217, JAPAN

E-mail address: `nishimoto@kinwu.ac.jp`