

ON ORDINARY PRIMES FOR MODULAR FORMS AND THE THETA OPERATOR

MASATAKA CHIDA AND MASANOBU KANEKO

(Communicated by Ken Ono)

ABSTRACT. We give a criterion for a prime being ordinary for a modular form, by using the theta operator of Ramanujan.

1. INTRODUCTION AND STATEMENT OF THE RESULT

A normalized Hecke eigenform is said to be ordinary at a prime p if p does not divide its p -th Fourier coefficient. In the theory of p -adic modular forms and Galois representations attached to modular forms, this notion has fundamental importance, and there is extensive literature on the subject.

In the present paper, we shall give a criterion for ordinarity in terms of certain polynomials attached to derivatives of given modular forms. Throughout the paper, the modular forms considered are those on the full modular group $\mathrm{SL}_2(\mathbb{Z})$.

For any $f = f(z) = \sum_{n=0}^{\infty} a(n)q^n$ ($q = e^{2\pi iz}$), we define

$$\theta f := q \frac{d}{dq} f = \sum_{n=0}^{\infty} n a(n) q^n.$$

This is the derivative with respect to $2\pi iz$, and is often referred to as the “theta operator” of Ramanujan. The derivative of a modular form is no longer modular but “quasimodular”, which means, in the case of $\mathrm{SL}_2(\mathbb{Z})$, that it is an isobaric element of the ring $\mathbb{C}[E_2, E_4, E_6]$. Here, $E_k = E_k(z)$ for even k is the standard Eisenstein series

$$E_k(z) = 1 - \frac{2k}{B_k} \sum_{n=1}^{\infty} \left(\sum_{d|n} d^{k-1} \right) q^n,$$

B_k being the k -th Bernoulli number. For $k \geq 4$, the function $E_k(z)$ is modular of weight k , but $E_2(z)$ is not quite modular. The operator θ preserves the ring $\mathbb{C}[E_2, E_4, E_6]$ (as is seen by Ramanujan’s formulae $\theta E_2 = (E_2^2 - E_4)/12$, $\theta E_4 = (E_2 E_4 - E_6)/3$, $\theta E_6 = (E_2 E_6 - E_4)/2$), and hence for any modular form f and non-negative integer n , $\theta^n f$ is an element in $\mathbb{C}[E_2, E_4, E_6]$.

To any $g \in \mathbb{C}[E_2, E_4, E_6]$, we attach a polynomial $F(g; X, Y, Z)$ in three variables so that

$$g(z) = F(g; E_2(z), E_4(z), E_6(z))$$

Received by the editors October 13, 2005 and, in revised form, November 15, 2005.

2000 *Mathematics Subject Classification*. Primary 11F33; Secondary 11F11.

Key words and phrases. Ordinary prime, theta operator.

The first author was supported in part by JSPS Research Fellowships for Young Scientists.

©2006 American Mathematical Society
Reverts to public domain 28 years from publication

holds. We also define its “modular part” $F^{(0)}(g; Y, Z)$ by

$$F^{(0)}(g; Y, Z) := F(g; 0, Y, Z).$$

If in particular g is modular (i.e., $g \in \mathbb{C}[E_4, E_6]$), then $F(g; X, Y, Z)$ is free from X and $F(g; X, Y, Z) = F^{(0)}(g; Y, Z)$. If g has p -integral Fourier coefficients, the polynomial F (and hence $F^{(0)}$) also has p -integral coefficients.

For a prime $p > 3$, set $H_p(Y, Z) = F^{(0)}(E_{p-1}; Y, Z) (= F(E_{p-1}; X, Y, Z))$. The polynomial $H_p(Y, Z)$ has p -integral coefficients, and $H_p(Y, Z) \bmod p$ is known as the “Hasse invariant” ([3], [4]).

Now we can state our main theorem.

Theorem 1.1. *Let $f(z) = \sum_{n=1}^{\infty} a(n)q^n$ be a normalized eigencusp form of weight k and p a prime number greater than k . Then the following conditions are equivalent:*

- (1) $a(p) \not\equiv 0 \pmod p$.
- (2) $H_p(Y, Z) \nmid F^{(0)}(\theta^{p-k+1} f; Y, Z) \pmod p$.

2. PROOF OF THE THEOREM AND A COROLLARY

In order to prove the theorem, we use the theory of filtration of modular forms modulo p developed by Swinnerton-Dyer [4], the theory of theta cycles by Tate [1], and a formula for the derivative $\theta^n f$. We first recall the definition of the filtration and then review theorems of Tate and Swinnerton-Dyer.

Let $M_k(\mathbb{Z}_{(p)})$ be the set of modular forms of weight k (on $\mathrm{SL}_2(\mathbb{Z})$) whose Fourier coefficients belong to $\mathbb{Z}_{(p)}$, the local ring of \mathbb{Q} at p . Following [4], let \widetilde{M}_k be the \mathbb{F}_p -vector space (in $\mathbb{F}_p[[q]]$) obtained from $M_k(\mathbb{Z}_{(p)})$ by reducing Fourier coefficients modulo p . We note that, since we have $E_{p-1} \equiv 1 \pmod p$ and $E_2 \equiv E_{p+1} \pmod p$ by the Kummer congruences of Bernoulli numbers, any quasimodular form having p -integral Fourier coefficients is congruent modulo p to a modular form of suitable weight.

Definition 2.1. For $f \in \widetilde{M}_k$, we define the filtration $w(f)$ of f to be the least ℓ such that f belongs to \widetilde{M}_ℓ . For a modular or quasimodular form f whose Fourier coefficients are p -integral, we shall write $w(f)$ instead of $w(f \bmod p)$.

We call an element in \widetilde{M}_k an eigenform if it is congruent modulo p to a Hecke-eigencusp form. Tate’s theory of theta cycles connects the ordinariness of an eigenform f to the filtration of the derivative of f .

Proposition 2.2 (Tate [1]). *Let $f = \sum_{n=1}^{\infty} a(n)q^n \in \widetilde{M}_k$ be an eigenform. We assume $k < p$ and $w(f) = k$. Then we have*

$$w(\theta^{p-k+1} f) = \begin{cases} 2p - k + 2 & \text{if } a(p) \not\equiv 0 \pmod p, \\ p - k + 3 & \text{if } a(p) \equiv 0 \pmod p. \end{cases}$$

(In [1] the assumption is weaker (that f is in the kernel of the “ U -operator”), but for our purpose it is enough to restrict to the case of eigenform.)

On the other hand, the filtration of a modular form g is related to the divisibility of $F^{(0)}(g; Y, Z) \bmod p$ by the Hasse invariant.

Proposition 2.3 (Swinnerton-Dyer [4, Lemma 5]). *For $g \in M_{k'}(\mathbb{Z}_{(p)})$, the following hold:*

- (1) *If $w(g) = k'$, then $H_p(Y, Z) \nmid F^{(0)}(g; Y, Z) \pmod p$.*
- (2) *If $w(g) = k' - p + 1$, then $H_p(Y, Z) \mid F^{(0)}(g; Y, Z) \pmod p$.*

Now assume that f is a normalized eigenform of weight k . The derivative $\theta^{p-k+1}f$ is quasimodular of weight $2p - k + 2$. If $\theta^{p-k+1}f$ is congruent modulo p to a (true) modular form g of weight $2p - k + 2$, then, combining Proposition 2.2 and Proposition 2.3 (with $k' = 2p - k + 2$), the condition $a(p) \not\equiv 0 \pmod p$ is equivalent to the polynomial $F^{(0)}(g; Y, Z) \pmod p$ not being divisible by $H_p(Y, Z) \pmod p$. Our theorem is therefore a consequence of the following observation that we can indeed take g to be the modular part of $\theta^{p-k+1}f$. Here, for a quasimodular form $g = \sum_{i=0}^m g_i E_2^i$, $g_i \in \mathbb{C}[E_4, E_6]$, we call g_0 its modular part.

Lemma 2.4. *Let $p > 3$ be a prime and f a modular form of weight $k < p$ with p -integral Fourier coefficients. Then we have*

$$\theta^{p-k+1}f \equiv (\theta^{p-k+1}f)_0 \pmod p.$$

This is a consequence of a general formula for $\theta^n f$ given in [5]. Recall that, if f is modular of weight k , then

$$\partial f := \theta f - \frac{k}{12} E_2 f$$

is modular of weight $k + 2$. For a modular form f of weight k , define a sequence of modular forms f_r of weight $k + 2r$ recursively by

$$f_{r+1} = \partial f_r - \frac{r(r+k-1)}{144} E_4 f_{r-1} \quad (r \geq 0)$$

with initial condition $f_0 = f$. Then the formula (37) in [5] is equivalent to the following closed formula.

Proposition 2.5. *Let f be a modular form of weight k . Then for any $n \geq 0$ we have*

$$\frac{\theta^n f}{n!} = \sum_{i=0}^n [k+n-1i] \frac{f_{n-i}}{(n-i)!} \left(\frac{E_2}{12}\right)^i.$$

When $n = p - k + 1$, the binomial coefficients $\binom{k+n-1}{i}$ are divisible by p for all $i > 0$, and hence Lemma 2.4 follows ($f_n = (\theta^n f)_0$). This completes the proof of the theorem. □

Here we give a corollary to the theorem. As in the theorem, assume that $f(z) = \sum_{n=1}^\infty a(n)q^n$ is a normalized eigenform of weight k and p is a prime number greater than k . We denote by $b(l, m, n)$ the coefficient of $X^l Y^m Z^n$ in $F(\theta^{p-k+1}f; X, Y, Z)$:

$$F(\theta^{p-k+1}f; X, Y, Z) = \sum_{2l+4m+6n=2p-k+2} b(l, m, n) X^l Y^m Z^n.$$

Corollary 2.6. (1) *Assume that $k \equiv 0 \pmod 6$ and $p \equiv 2 \pmod 3$.*

$$\text{If } b(0, 0, \frac{2p-k+2}{6}) \not\equiv 0 \pmod p, \text{ then } a(p) \not\equiv 0 \pmod p.$$

(2) *Assume that $k \equiv 0 \pmod 4$ and $p \equiv 3 \pmod 4$.*

$$\text{If } b(0, \frac{2p-k+2}{4}, 0) \not\equiv 0 \pmod p, \text{ then } a(p) \not\equiv 0 \pmod p.$$

Proof. We only prove (1), the proof of (2) being similar. Write

$$H_p(Y, Z) = \sum_{4m+6n=p-1} c(m, n)Y^m Z^n.$$

By the assumption, $p-1$ is not divisible by 6, and hence the term with $m=0$ does not occur on the right. Therefore, if $b(0, 0, \frac{2p-k+2}{6}) \not\equiv 0 \pmod{p}$, the polynomial $F(\theta^{p-k+1}f; X, Y, Z) \pmod{p}$ is not a multiple of $H_p(Y, Z) \pmod{p}$, and thus $a(p) \not\equiv 0 \pmod{p}$ by Theorem 1.1. \square

3. RELATION TO SUPERSINGULAR j -INVARIANTS OF ELLIPTIC CURVES

We may rephrase the theorem in terms of the supersingular j -polynomial.

Let f be a modular form of weight k . Write $k = 12m + 4\delta + 6\varepsilon$ with $m \geq 0$, $\delta \in \{0, 1, 2\}$, $\varepsilon \in \{0, 1\}$. Then there exists a unique polynomial $G(f; x)$ such that

$$f(z) = \Delta(z)^m E_4(z)^\delta E_6(z)^\varepsilon G(f; j(z)),$$

where $\Delta(z) = (E_4(z)^3 - E_6(z)^2)/1728$ is the discriminant function and $j(z) = E_4(z)^3/\Delta(z)$ is the modular invariant. Moreover we put

$$\tilde{G}(f; x) := x^\delta (x - 1728)^\varepsilon G(f; x).$$

For a prime number p , we define the supersingular j -polynomial $S_p(x)$ by

$$S_p(x) := \prod_{E/\overline{\mathbb{F}}_p: \text{supersingular}} (x - j(E)) \in \mathbb{F}_p[x],$$

where the product runs over the isomorphism classes of supersingular elliptic curves in characteristic p and $j(E)$ is the j -invariant of E . Assume $p > 3$. A theorem of Deligne (cf. [3], [2]) then asserts that

$$\tilde{G}(E_{p-1}; x) \equiv S_p(x) \pmod{p}.$$

By this and Theorem 1.1, we have the following.

Theorem 3.1. *The assumption being the same as in Theorem 1.1, the following conditions are equivalent:*

- (1) $a(p) \not\equiv 0 \pmod{p}$.
- (2) $S_p(x) \not\equiv \tilde{G}((\theta^{p-k+1}f)_0; x) \pmod{p}$.

ACKNOWLEDGEMENT

The first author would like to thank Ken Ono for his helpful comments on the last section.

REFERENCES

- [1] N. Jochowitz, *A study of the local components of the Hecke algebra mod l* , Trans. Amer. Math. Soc. **270** (1982), no. 1, 253–267. MR0642340 (83e:10033a)
- [2] M. Kaneko, D. Zagier, *Supersingular j -invariants, hypergeometric series, and Atkin's orthogonal polynomials*, Computational perspectives on number theory (Chicago, IL, 1995), 97–126, AMS/IP Stud. Adv. Math., **7**. MR1486833 (99b:11064)
- [3] J.-P. Serre, *Congruences et formes modulaires (d'après H. P. F. Swinnerton-Dyer)*, Séminaire Bourbaki 1971/72, **416** (Œuvres, vol. 3, 74–88). MR0466020 (57:5904a)

- [4] H. P. F. Swinnerton-Dyer, *On ℓ -adic representations and congruences for coefficients of modular forms*, Lect. Notes. in Math. **350** (1973), 1–55. MR0406931 (53:10717a)
- [5] D. Zagier, *Modular forms and differential operators*, Proc. Indian Acad. Sci. (Math. Sci.) **104** (1994), 57–75. MR1280058 (95d:11048)

INSTITUTE OF MATHEMATICS, TOHOKU UNIVERSITY, AOBA, SENDAI, 980-8578, JAPAN
E-mail address: `sa2m22@math.tohoku.ac.jp`

FACULTY OF MATHEMATICS, KYUSHU UNIVERSITY 33, FUKUOKA, 812-8581, JAPAN
E-mail address: `mkaneko@math.kyushu-u.ac.jp`