

MHF Preprint Series

Kyushu University
21st Century COE Program
Development of Dynamic Mathematics with
High Functionality

An explicit formula of the Shapley value for a cooperative game induced from the conjugate point

T. Fuchikami & H. Kawasaki

MHF 2005-33

(Received November 14, 2005)

Faculty of Mathematics
Kyushu University
Fukuoka, JAPAN

AN EXPLICIT FORMULA OF THE SHAPLEY VALUE FOR A COOPERATIVE GAME INDUCED FROM THE CONJUGATE POINT ¹

TAKEAKI FUCHIKAMI and HIDEFUMI KAWASAKI

Abstract. The conjugate point was introduced by Jacobi to derive a sufficient optimality condition for a variational problem. Recently, the conjugate point was defined for an extremal problem in \mathbb{R}^n . The key of the conjugate point is cooperation of variables. Namely, when there exists a conjugate point for a stationary solution $x \in \mathbb{R}^n$, we can improve the solution by suitably changing some of the variables. We call such a set of variables a strict conjugate set. This idea leads us to a cooperative game, which we call a conjugate-set game. The Shapley value is an important value in game theory. It evaluates player's contribution in the cooperative game. However, its calculation is usually very hard. The purpose of this paper is to give an explicit formula of the Shapley value for the conjugate-set game induced from the shortest path problem on an ellipsoid.

1. INTRODUCTION

The conjugate point was originally introduced to guarantee local optimality of a stationary solution $x(t)$ for the simplest problem in the calculus of variations

$$\begin{aligned} & \text{Minimize} && \int_0^T f(t, x(t), \dot{x}(t)) dt \\ & \text{subject to} && x(0) = A, x(T) = B \end{aligned}$$

where A and B are given points, see e.g. Gelfand and Fomin [2]. Recently, the conjugate point was defined for an extremal problem with n variables

$$(P_0) \quad \text{Minimize} \quad f(x), \quad x \in \mathbb{R}^n,$$

see Kawasaki [3][4].




FIGURE 1. the shortest polygonal path problem

¹2000 Mathematics Subject Classification. 91A12, 90C30, 49K40

Key words and phrases. Shapley value, cooperative game, conjugate-set game, conjugate point.
 This research is supported by Kyushu Univ. 21st Century COE Program (Development of Dynamic Mathematics with High Functionality) and the Grant-in Aid for General Scientific Research from the Japan Society for the Promotion of Science 14340037.

One can see the typical example of the conjugate point for (P_0) in the shortest polygonal path problem on an ellipsoid

$$\frac{x^2}{a^2} + \frac{y^2}{a^2} + \frac{z^2}{c^2} = 1. \quad (1)$$

This problem is to find the shortest polygonal path

$$A = X_0, X_1, \dots, X_n, X_{n+1} = B$$

joining two given points $A := (a, 0, 0)$ and $B := (a \cos T, a \sin T, 0)$, where each X_k can move on a longitude ℓ_i that is equally located between A and B . Since each X_k has one-dimensional freedom, this problem is formulated as (P_0) , and the equatorial polygonal path is a stationary solution for (P_0) .

Further, whether the stationary solution is minimal or not depends on T . It is minimal when $T < a\pi/c$, and not minimal when $T > a\pi/c$ and n is sufficiently large. In the latter case, we call the first number k satisfying $(k+1)T/(n+1) > a\pi/c$ a strict conjugate point, which matches the classical conjugate point, see [4].

When there exists a strict conjugate point $k \leq n$, the Hessian matrix $A := f''(x)$ is not positive semidefinite. So, according to Sylvester's criterion, A has a negative leading principal minor. Then we can improve the stationary solution $x = (x_1, \dots, x_n)$ by suitably changing some variables $\{x_i\}_{i \in I}$. We call such a subset $\{x_i\}_{i \in I}$ (or I) a strict conjugate set. Namely, when $A_I := (a_{ij})_{i,j \in I}$ has a negative principal minor, we call $\{x_i\}_{i \in I}$ (or I) a strict conjugate set, see Kawasaki [6][7]. In this paper, we consider a cooperative game based on strict conjugate sets, and we present an explicit formula of the Shapley value for this game.

2. DEFINITIONS AND NOTATIONS

In this section, we first define conjugate-set game induced from the shortest path problem on the ellipsoid (1). Next, we introduce tools $I(i; S)$ and $Ker(i; S)$ to compute the Shapley value.

Definition 1. Let $N = \{1, \dots, n\}$ be the players set and $1 \leq k \leq n$ a natural number. We call a subset

$$[j, j+k-1] := \{j, j+1, \dots, j+k-1\}$$

of N an interval of length k . For any subset S of N , we define a characteristic function $v(S)$ as the maximum number of disjoint intervals of length k contained in S . We call this cooperative game a conjugate-set game induced from the shortest path problem on an ellipsoid and denote it by $G(n, k)$.



FIGURE 2. When $k = 3$ and S consists of circles, $v(S) = 3$.

Throughout this paper, we put

$$n = pk + r \quad (0 \leq r \leq k-1). \quad (2)$$

We denote by $\phi_i(n, k)$ or ϕ_i the Shapley value of $G(n, k)$. That is,

$$\phi_i(n, k) = \sum_{i \in S \subset N} \frac{(s-1)!(n-s)!}{n!} \{v(S) - v(S - \{i\})\}, \quad (3)$$

where $s := \#S$. The following expression is well-known, see e.g. Aumann et al [1].

$$\phi_i(n, k) = \sum_{\pi \in \Pi} \frac{1}{n!} \{v(S_{\pi, i}) - v(S_{\pi, i} - \{i\})\}, \quad (4)$$

where Π denotes the set of all permutation on N and $S_{\pi, i}$ denotes the union of $\{i\}$ and the set of all players that precedes player i with respect to π , that is,

$$S_{\pi, i} = \{\pi(j) \mid 1 \leq j \leq \pi^{-1}(i)\}. \quad (5)$$

We note that $S_{\pi, i}$ plays an important role in this paper.

The following proposition is evident from symmetry of the characteristic function.

Proposition 1. *For any $1 \leq i \leq n$, it holds that $\phi_i = \phi_{n-i+1}$.*

Definition 2. *Any element of $W_S := \{i \mid v(S) - v(S - \{i\}) = 1\}$ is called a pivot of S .*

Then the Shapley value (4) is simply written as

$$\phi_i = \frac{1}{n!} \#\{\pi \mid i \in W_{S_{\pi, i}}\}. \quad (6)$$

So it suffices to test whether $i \in S$ is a pivot of S or not in order to compute ϕ_i . For this aim, we introduce two subsets of S , say $I(i; S)$ and $Ker(i; S)$.

Definition 3. *For any $i \in N$ and $S \subset N$ including i , we denote by $I(i; S) \subset S$ the maximum interval including i . We denote by $Ker(i; S)$ the remainder of $I(i; S)$ after removing intervals of length k from both sides of i as much as possible with keeping i .*




FIGURE 3. When $k = 3$ and S consists of circles, $I(i; S)$ consists of gray circles and $Ker(i; S)$ consists of black circles.

Theorem 1. *The following conditions are equivalent to each others.*

- (i) $i \in W_S$,
- (ii) $i \in W_{I(i; S)}$,
- (iii) $i \in W_{Ker(i; S)}$,
- (iv) $\#Ker(i; S) \geq k$.

Proof. (i) \Leftrightarrow (ii): Obvious. (ii) \Leftrightarrow (iii): It is enough to consider the case that $[j_1, j_2] := I(i; S) \neq Ker(i; S)$. Because of symmetry, we may assume that $j_1 + k < i$. Then, since $v([j_1, j_2]) = v([j_1 + k, j_2]) + 1$ and $v([j_1, j_2] - \{i\}) = v([j_1 + k, j_2] - \{i\}) + 1$, we have

$$\begin{aligned} i \in W_{I(i; S)} &\Leftrightarrow v([j_1, j_2]) - v([j_1, j_2] - \{i\}) = 1 \\ &\Leftrightarrow v([j_1 + k, j_2]) - v([j_1 + k, j_2] - \{i\}) = 1 \\ &\Leftrightarrow i \in W_{I(i; S) - [j_1, j_1 + k - 1]}. \end{aligned}$$

Repeating this procedure, we finally get $Ker(i; S)$ as the remainder and see the equivalence of (ii) and (iii). (iii) \Leftrightarrow (iv): Since we can not remove any interval of length k from $Ker(i; S)$ without deleting i , this assertion is clear. \square

3. THE SHAPLEY VALUE OF PLAYER 1

In this section, we compute the Shapley value ϕ_1 . It follows from Theorem 1 that

$$1 \in W_S \Leftrightarrow \#Ker(1; S) \geq k \Leftrightarrow Ker(1; S) = [1, k]. \quad (7)$$

Since $I(1; S)$ is obtained by adding disjoint intervals of length k to $Ker(1; S)$, we get from (7) that

$$\{I(1; S) \mid 1 \in W_S\} = \{[1, mk] \mid 1 \leq m \leq p\}. \quad (8)$$

Lemma 1. *Let $1 \leq m \leq p$. Then π satisfies $I(1; S_{\pi, 1}) = [1, mk]$ if and only if*

$$\pi^{-1}(j) < \pi^{-1}(1) \quad \forall j \in [2, mk] \quad (9)$$

and either (a) $\pi^{-1}(mk + 1) > \pi^{-1}(1)$ or (b) $mk = n$ holds.

Proof. Necessity: Since 1 joins $I(1; S_{\pi, 1}) = [1, mk]$ last, (9) is clear. If $mk < n$ and $mk + 1$ joins $S_{\pi, 1}$ before 1, then the interval $I(1; S_{\pi, 1})$ contains $[1, mk + 1]$. Sufficiency is evident. \square

Theorem 2.

$$\phi_1 = \begin{cases} \sum_{m=1}^{p-1} \frac{1}{mk(mk+1)} + \frac{1}{pk} & \text{if } r = 0, \\ \sum_{m=1}^p \frac{1}{mk(mk+1)} & \text{if } r \neq 0, \end{cases} \quad (10)$$

where $n = pk + r$ ($0 \leq r \leq k - 1$).

Proof. By (6), it suffices to compute $\#\{\pi \mid 1 \in W_{S_{\pi, 1}}\}$. Combining (7) and (8), it equals $\#\{\pi \mid 1 \leq \exists m \leq p, I(1; S_{\pi, 1}) = [1, mk]\}$. Further, it equals

$$\#\{\pi \mid \pi \text{ satisfies (9) and (a)}\} + \#\{\pi \mid \pi \text{ satisfies (9) and (b)}\}. \quad (11)$$

For each m , the first term of (11) is given by

$$\binom{n}{mk+1} (n - mk - 1)! (mk - 1)! = \frac{n!}{mk(mk+1)}. \quad (12)$$

Indeed, such a permutation π satisfies

$$\pi^{-1}(j) < \pi^{-1}(1) < \pi^{-1}(mk + 1) \quad \forall j \in [2, mk]. \quad (13)$$

There are ${}_nC_{mk+1}$ ways to choose $P := \pi^{-1}([1, mk+1]) \subset N$. Since $\pi^{-1}([2, mk])$ can freely share the first $mk-1$ places of P , and since the complement of P can be freely shared by other $n-mk-1$ numbers, we get (12). On the other hand, since case (b) occurs only when $m=p$ and $n=pk$ (so that $r=0$), we similarly see that the second term of (11) is given by

$$(n-1)!. \quad (14)$$

In the case of $r=0$, ϕ_1 is equal to the total sum of (12)/ $n!$ ($m=1, \dots, p-1$) and (14)/ $n!$. Otherwise, ϕ_1 is equal to the total sum of (12)/ $n!$ ($m=1, \dots, p$). \square

4. A RECURRENCE RELATION OF $\{\phi_i\}$: CASE 1

Starting with ϕ_1 , we compute ϕ_2, ϕ_3 , and so on. For this aim, we compute the difference between ϕ_i and ϕ_{i+1} . Because of symmetry of the game, it suffices to consider $1 \leq i \leq \lfloor \frac{n-1}{2} \rfloor$, where $\lfloor \cdot \rfloor$ denotes Gauss's symbol. We will deal with this problem in three cases.

Case 1: $n-k+1 \leq i \leq k-1$ (This is the case that $k \leq \frac{n}{2} + 1$),

Case 2: $1 \leq i \leq \min\{n-k, k-1\}$,

Case 3: $k \leq i \leq \lfloor \frac{n-1}{2} \rfloor$.

Before dealing with Case 1, we present a lemma that is applicable to any case.

Lemma 2.

$$\phi_{i+1} = \phi_i + \delta_i^+ - \delta_i^-, \quad (15)$$

where

$$\delta_i^+ := \#\{\pi \mid i \notin W_{S_{\pi, i+1}}, i+1 \in W_{S_{\pi, i+1}}\}/n! \quad (16)$$

and

$$\delta_i^- := \#\{\pi \mid i \in W_{S_{\pi, i}}, i+1 \notin W_{S_{\pi, i}}\}/n!. \quad (17)$$

Proof. Since

$$\begin{aligned} & n!(\phi_{i+1} - \phi_i) \\ &= \#\{\pi \mid i+1 \in W_{S_{\pi, i+1}}\} - \#\{\pi \mid i \in W_{S_{\pi, i}}\} \\ &= \#\{\pi \mid i, i+1 \in W_{S_{\pi, i+1}}\} + \#\{\pi \mid i \notin W_{S_{\pi, i+1}}, i+1 \in W_{S_{\pi, i+1}}\} \\ &\quad - \#\{\pi \mid i, i+1 \in W_{S_{\pi, i}}\} - \#\{\pi \mid i \in W_{S_{\pi, i}}, i+1 \notin W_{S_{\pi, i}}\}. \end{aligned}$$

it suffices to prove

$$\#\{\pi \mid i, i+1 \in W_{S_{\pi, i+1}}\} = \#\{\pi \mid i, i+1 \in W_{S_{\pi, i}}\}. \quad (18)$$

We define a bijection $f : \Pi \rightarrow \Pi$ by $f(\pi) := (i, i+1) \circ \pi$, where $(i, i+1)$ is a transposition. Then, for any $\pi \in \Pi$ such that $i, i+1 \in W_{S_{\pi, i}}$, due to definition of W_S ,

$$v(S_{\pi, i}) - v(S_{\pi, i} - \{i\}) = v(S_{\pi, i}) - v(S_{\pi, i} - \{i+1\}) = 1. \quad (19)$$

Since $S_{\pi, i} = S_{f(\pi), i+1}$, (19) implies that $i, i+1 \in W_{S_{f(\pi), i+1}}$. That is, $f(\{\pi \mid i, i+1 \in W_{S_{\pi, i}}\}) \subset \{\pi \mid i, i+1 \in W_{S_{\pi, i+1}}\}$. Since f is an injection, we have

$$\#\{\pi \mid i, i+1 \in W_{S_{\pi, i}}\} \leq \#\{\pi \mid i, i+1 \in W_{S_{\pi, i+1}}\}. \quad (20)$$

The converse inequality is similarly obtained. \square

Let us now consider Case 1.

Theorem 3. For any i such that $n - k + 1 \leq i \leq k - 1$, it holds that

$$\delta_i^+ = \delta_i^- = 0. \quad (21)$$

Therefore, $\phi_{n-k+1} = \phi_{n-k+2} = \cdots = \phi_{k-1}$.

Proof. By Theorem 1, $i \in W_{S_{\pi,i}}$ if and only if $\#Ker(i; S_{\pi,i}) \geq k$. For such a π , since $n - k + 1 \leq i \leq k - 1$ and since we can not remove any intervals of length k from $Ker(i; S_{\pi,i})$ without removing i , we have

$$\{Ker(i; S_{\pi,i}) \mid i \in W_{S_{\pi,i}}\} = \left\{ \begin{array}{cccc} [1, k], & [1, k+1], & \dots & [1, n], \\ & [2, k+1], & \dots & [2, n], \\ & & \ddots & \vdots \\ & & & [n-k+1, n] \end{array} \right\}. \quad (22)$$

We similarly see that $\{Ker(i+1; S_{\pi,i+1}) \mid i+1 \in W_{S_{\pi,i+1}}\}$ coincides with set (22). So $\{\pi \mid i \in W_{S_{\pi,i}}, i+1 \notin W_{S_{\pi,i}}\}$ is empty. (Remark that not $i+1 \notin W_{S_{\pi,i+1}}$ but $i+1 \notin W_{S_{\pi,i}}$.) Indeed, if π is an element of this set, then $Ker(i; S_{\pi,i})$ is one of the intervals in (22) and i is its element. Since $i+1 \leq k$, $i+1$ belongs to the interval, which implies that $i+1$ is also an element of $S_{\pi,i}$. Since the length of the interval is greater than or equal to k , we see from Theorem 1 and (22) that $i+1 \in W_{S_{\pi,i}}$. Therefore $\delta_i^- = 0$. Similarly, we have $\delta_i^+ = 0$. \square

5. A RECURRENCE RELATION OF $\{\phi_i\}$: CASE 2

In this section, we consider the case that $1 \leq i \leq \min\{n - k, k - 1\}$.

Since $i \leq k$ and $k + i - 1 \leq n$, we get

$$\{Ker(i; S_{\pi,i}) \mid i \in W_{S_{\pi,i}}\} = \left\{ \begin{array}{cccc} [1, k], & [1, k+1], & \dots & [1, k+i-1], \\ & [2, k+1], & \dots & [2, k+i-1], \\ & & \ddots & \vdots \\ & & & [i, k+i-1] \end{array} \right\} \quad (23)$$

as well as (22), where the difference between (22) and (23) is caused from $k + i - 1 \leq n$. Since $i + 1 \leq k$ and $k + i \leq n$, we similarly see that $\{Ker(i+1; S_{\pi,i+1}) \mid i+1 \in W_{S_{\pi,i+1}}\}$ equals

$$\left\{ \begin{array}{cccc} [1, k], & [1, k+1], & \dots & [1, k+i-1], & [1, k+i], \\ & [2, k+1], & \dots & [2, k+i-1], & \vdots \\ & & \ddots & \vdots & \vdots \\ & & & [i, k+i-1], & [i, k+i], \\ & & & & [i+1, k+i] \end{array} \right\}. \quad (24)$$

Comparing (23) and (24), we get $\delta_i^- = 0$ as well as Theorem 3. On the other hand,

$$\begin{aligned} & \{Ker(i+1; S_{\pi,i+1}) \mid i \notin W_{S_{\pi,i+1}}, i+1 \in W_{S_{\pi,i+1}}\} \\ &= \{[j_1, k+i] \mid 1 \leq j_1 \leq i+1\}. \end{aligned} \quad (25)$$

Indeed, since i belongs to any interval in (24) except $[i + 1, k + i]$, Theorem 1 asserts that $i \notin W_{S_{\pi, i+1}}$ implies $\#Ker(i; S_{\pi, i+1}) < k$. The intervals in (24) that satisfy this condition are those in the last column of (24). So we get (25).

Since $I(i + 1; S_{\pi, i+1})$ is an interval obtained by adding disjoint intervals of length k to $Ker(i + 1; S_{\pi, i+1})$, we get from (25) that

$$\begin{aligned} & \{I(i + 1; S_{\pi, i+1}) \mid i \notin W_{S_{\pi, i+1}}, i + 1 \in W_{S_{\pi, i+1}}\} \\ &= \{[j_1, mk + i] \mid 1 \leq j_1 \leq i + 1, m \geq 1, mk + i \leq n\}. \end{aligned} \quad (26)$$

Lemma 3. *Let $m \geq 1$ satisfy $mk + i \leq n$. Then there exists $1 \leq j_1 \leq i + 1$ such that $I(i + 1; S_{\pi, i+1}) = [j_1, mk + i]$ if and only if*

$$\pi^{-1}(j) < \pi^{-1}(i + 1) \quad \forall j \in [i + 2, mk + i] \quad (27)$$

and either (c) $\pi^{-1}(mk + i + 1) > \pi^{-1}(i + 1)$ or (d) $mk + i = n$ holds.

Proof. Necessity: Since $i + 1$ joins $S_{\pi, i+1}$ last, $I(i + 1; S_{\pi, i+1}) = [j_1, mk + i]$ implies that (27) and $mk + i + 1$ dose not joint $S_{\pi, i+1}$ before $i + 1$ if $mk + i < n$. Conversely, it follows from (c) or (d) that any number greater than $mk + i$ does not join $S_{\pi, i+1}$ before $i + 1$. Hence $mk + i$ is the maximum number of $I(i + 1; S_{\pi, i+1})$. Since $I(i + 1; S_{\pi, i+1})$ is an interval, it has a form of $[j_1, mk + i]$ for some $1 \leq j_1 \leq i + 1$. \square

Theorem 4. *In the case of $1 \leq i \leq \min\{n - k, k - 1\}$, it holds that*

$$\delta_i^+ = \begin{cases} \sum_{m=1}^p \frac{1}{mk(mk + 1)} & 1 \leq i \leq r - 1, \\ \sum_{m=1}^{p-1} \frac{1}{mk(mk + 1)} + \frac{1}{pk} & i = r, \\ \sum_{m=1}^{p-1} \frac{1}{mk(mk + 1)} & r + 1 \leq i \leq k - 1, \end{cases} \quad (28)$$

$$\delta_i^- = 0, \quad (29)$$

where $n = pk + r$ ($0 \leq r \leq k - 1$).

Proof. Assume that π satisfies that $i \notin W_{S_{\pi, i+1}}$ and $i + 1 \in W_{S_{\pi, i+1}}$. Then it is easily seen from (26) and Lemma 3 that (27) and either (c) or (d) hold. The number of π 's satisfying (27) and (c) is given by

$$\binom{n}{mk + 1} (mk - 1)! (n - mk - 1)! = \frac{n!}{mk(mk + 1)}. \quad (30)$$

Indeed, such a permutation π satisfies

$$\pi^{-1}(j) < \pi^{-1}(i + 1) < \pi^{-1}(mk + i + 1) \quad \forall j \in [i + 2, mk + i]. \quad (31)$$

There are ${}_n C_{mk+1}$ ways to choose $P := \pi^{-1}([i + 1, mk + i + 1])$. Since $\pi^{-1}([i + 2, mk + i])$ can freely share the first $mk - 1$ places of P , and since the complement of P can be freely shared by other $n - mk - 1$ numbers, we get (30).

Since case (d) occurs only when $m = p$, we similarly see that the number of π 's satisfying (27) and (d) is given by

$$\binom{n}{pk} (pk-1)! (n-pk)! = \frac{n!}{pk}. \quad (32)$$

In the cases of $0 \leq i < r$, since $mk + i < n$ for any $1 \leq m \leq p$, δ_i^+ equals the total sum of (30)/ $n!$ ($m = 1, \dots, p$). In the cases of $i = r$, since $mk + i$ equals n only when $m = p$, δ_i^+ equals the total sum of (30)/ $n!$ ($m = 1, \dots, p-1$) and (32)/ $n!$. In the cases of $i > r$, since $mk + i < n$ for any $1 \leq m \leq p-1$, δ_i^+ equals the total sum of (30)/ $n!$ ($m = 1, \dots, p-1$). \square

6. A RECURRENCE RELATION OF $\{\phi_i\}$: CASE 3

In this section, we consider the case of $k \leq i \leq \lfloor \frac{n-1}{2} \rfloor$. Then i is expressed as

$$i = qk + s \quad (33)$$

for some $q \geq 1$ and $0 \leq s \leq k-1$. Since $i+k \leq n$, we get from Theorem 1 that $\{Ker(i; S_{\pi,i}) \mid i \in W_{S_{\pi,i}}\}$ is given by

$$\left\{ \begin{array}{cccc} [i-k+1, i] & [i-k+1, i+1] & \dots & [i-k+1, i+k-1] \\ & [i-k+2, i+1] & \dots & [i-k+2, i+k-1] \\ & & \ddots & \vdots \\ & & & [i, i+k-1] \end{array} \right\} \quad (34)$$

and $\{Ker(i+1; S_{\pi,i+1}) \mid i+1 \in W_{S_{\pi,i+1}}\}$ is given by

$$\left\{ \begin{array}{cccc} [i-k+2, i+1] & \dots & [i-k+2, i+k-1] & [i-k+2, i+k] \\ & \ddots & \vdots & \vdots \\ & & [i, i+k-1] & [i, i+k] \\ & & & [i+1, i+k] \end{array} \right\}. \quad (35)$$

So, as well as (25), we have

$$\begin{aligned} \{Ker(i+1; S_{\pi,i+1}) \mid i \notin W_{S_{\pi,i+1}}, i+1 \in W_{S_{\pi,i+1}}\} \\ = \{[j_1, i+k] \mid i-k+2 \leq j_1 \leq i+1\} \end{aligned} \quad (36)$$

and

$$\begin{aligned} \{Ker(i; S_{\pi,i}) \mid i \in W_{S_{\pi,i}}, i+1 \notin W_{S_{\pi,i}}\} \\ = \{[i-k+1, j_2] \mid i \leq j_2 \leq i+k-1\}. \end{aligned} \quad (37)$$

As well as (26), we get from (36) and (37) that

$$\begin{aligned} \{I(i+1; S_{\pi,i+1}) \mid i \notin W_{S_{\pi,i+1}}, i+1 \in W_{S_{\pi,i+1}}\} \\ = \{[j_1, mk+i] \mid 1 \leq j_1 \leq i+1, m \geq 1, mk+i \leq n\} \end{aligned} \quad (38)$$

and

$$\begin{aligned} \{I(i; S_{\pi,i}) \mid i \in W_{S_{\pi,i}}, i+1 \notin W_{S_{\pi,i}}\} \\ = \{[i - mk + 1, j_2] \mid i \leq j_2 \leq n, m \geq 1, 1 \leq i - mk + 1\}. \end{aligned} \quad (39)$$

Lemma 4. *Let $m \geq 1$ satisfy $mk + i \leq n$. Then there exists $i \leq j_2 \leq n$ such that $I(i; S_{\pi,i}) = [i - mk + 1, j_2]$ if and only if*

$$\pi^{-1}(j) < \pi^{-1}(i) \quad \forall j \in [i - mk + 1, i - 1] \quad (40)$$

and either (e) $\pi^{-1}(i - mk) > \pi^{-1}(i)$ or (f) $i - mk + 1 = 1$ holds.

Proof. Almost same with Lemma 3. The only difference is that we make $I(i; S_{\pi,i})$ by attaching intervals of length k to $\text{Ker}(i; S_{\pi,i})$ from not the right side of i but the left side of i . \square

Theorem 5. *In the case of $k \leq i \leq \lfloor \frac{n-1}{2} \rfloor$, it holds that*

$$\delta_i^+ = \begin{cases} \sum_{m=1}^{p-q} \frac{1}{mk(mk+1)} & 0 \leq s \leq r-1, \\ \sum_{m=1}^{p-q-1} \frac{1}{mk(mk+1)} + \frac{1}{(p-q)k} & s = r, \\ \sum_{m=1}^{p-q-1} \frac{1}{mk(mk+1)} & r+1 \leq s \leq k-1, \end{cases} \quad (41)$$

and

$$\delta_i^- = \begin{cases} \sum_{m=1}^{q-1} \frac{1}{mk(mk+1)} + \frac{1}{qk} & s = 0, \\ \sum_{m=1}^q \frac{1}{mk(mk+1)} & s \neq 0, \end{cases} \quad (42)$$

where q and j are defined by (33).

Proof. One can easily prove (41) as well as (28). The only difference is that p is replaced by $p - q$. The difference comes from that i is expressed as $i = qk + s$. So, the condition $m \geq 1$ and $mk + i \leq n$ in (38) is equivalent to $m \geq 1$ and $(m + q)k + s \leq n$. When $s > r$, the latter implies that $1 \leq m \leq p - q - 1$. When $s \leq r$, it implies that $1 \leq m \leq p - q$. In particular, when $s = r$, $m = p - q$ corresponds to (d) in Lemma 3.

We use (39) and Lemma 4 to prove (42). By Lemma 4, π satisfies $i \in W_{S_{\pi,i}}$ and $i + 1 \notin W_{S_{\pi,i}}$ if and only if π satisfies (40) and either (e) or (f). The condition $m \geq 1$ and $1 \leq i - mk + 1$ in (39) is equivalent to $1 \leq m \leq q$. In particular, when $s = 0$, $m = q$ corresponds to (f). So we get (42). \square

Following is the graphs of the Shapley values of 28 players.




FIGURE 4. Shapley values of 28 players. Left: $k = 3$, Right: $k = 12$




FIGURE 5. Shapley values of 28 players when $k = 20$.

7. THE MAXIMAL VALUES OF THE SHAPLEY VALUE

In Figure 4, the maximum value of the Shapley value is attained by $i = k$ and $i = n + 1 - k$. The aim of this section is to show that this is true for any $2 \leq k \leq n/2$. Otherwise, the graph of the Shapley value has a shape in Figure 5. We first consider the case of $k \geq n/2 + 1$.

Theorem 6. *When $k \geq n/2 + 1$, it holds that*

$$\phi_1 < \phi_2 < \cdots < \phi_{n-k+1} = \cdots = \phi_k > \phi_{k+1} > \cdots > \phi_n.$$

Proof. The assertion is a direct consequence of Theorems 3 and 4. \square

Next, we consider the case of $2 \leq k \leq n/2$. We list up the maximal values of the Shapley value. By virtue of symmetry of the Shapley value, it suffices to consider $i \leq [(n-1)/2]$, so that $i+1 \leq [(n+1)/2]$. Since n and i are expressed as $n = pk + r$ and $i = qk + s$, $i \leq [(n-1)/2]$ implies that

$$(p-2q)k + r - 2s - 1 \geq 0. \quad (43)$$

Since $r \leq k$, we see from (43) that $p \geq 2q$.

Theorem 7. *The maximal points of $\{\phi_i\}_i$ on the interval $[1, [(n+1)/2]]$ are $\{k, 2k, \dots\}$.*

Proof. Figure 6 below shows the outline of the proof. Step 1. It follows from Theorem 4 that $\phi_1 < \phi_2 < \cdots < \phi_k$. Step 2. We show

$$\phi_{qk+1} - \phi_{qk} < 0 \quad (q = 1, 2, \dots). \quad (44)$$




FIGURE 6. $\phi_{qk} > \phi_{qk+1} \leq \phi_{qk+2} \leq \dots \leq \phi_{q(k+1)}$.

We get from Theorem 5 that, for any $q = 1, 2, \dots$,

$$\phi_{qk+1} - \phi_{qk} = \begin{cases} \sum_{m=q}^{p-q} \frac{1}{mk(mk+1)} - \frac{1}{qk} & r > 0 \\ \sum_{m=q}^{p-q-1} \frac{1}{mk(mk+1)} + \frac{1}{(p-q)k} - \frac{1}{qk} & r = 0. \end{cases} \quad (45)$$

Indeed, since $s = 0$ for $i = qk$, the first two cases in (41) and the first case in (42) are applicable, and we easily get (45). Further, we get (44) from (45). Indeed, for any $r > 0$ and $p = 2q$, we have

$$\phi_{qk+1} - \phi_{qk} = \frac{1}{qk} \left(\frac{1}{qk+1} - 1 \right) < 0. \quad (46)$$

For any $r > 0$ and $p > 2q$, we have

$$\begin{aligned} \phi_{qk+1} - \phi_{qk} &= \sum_{m=q}^{p-q-1} \frac{1}{mk(mk+1)} + \frac{1}{(p-q)k\{(p-q)k+1\}} - \frac{1}{qk} \\ &< \sum_{m=q}^{p-q-1} \frac{1}{mkmk} + \frac{1}{(p-q)k} - \frac{1}{qk} \\ &= \frac{1}{k} \left(\frac{1}{k} \sum_{m=q}^{p-q-1} \frac{1}{m^2} + \frac{1}{p-q} - \frac{1}{q} \right). \end{aligned}$$

For $r = 0$, we have from (45) that

$$\phi_{qk+1} - \phi_{qk} < \frac{1}{k} \left(\frac{1}{k} \sum_{m=q}^{p-q-1} \frac{1}{m^2} + \frac{1}{p-q} - \frac{1}{q} \right).$$

So, letting $f(p) := \frac{1}{k} \sum_{m=q}^{p-q-1} \frac{1}{m^2} + \frac{1}{p-q} - \frac{1}{q}$, we see that

$$0 \geq f(2q+1) > f(2q+2) > \dots > f(p). \quad (47)$$

In fact,

$$f(2q+1) = \frac{1}{kq^2} + \frac{1}{q+1} - \frac{1}{q} = \frac{1-q(k-1)}{kq^2(q+1)} \leq 0$$

and

$$f(2q+j+1) - f(2q+j) = \frac{1}{k(q+j)^2} + \frac{1}{q+j+1} - \frac{1}{q+j} = \frac{1-(q+j)(k-1)}{k(q+j)^2(q+j+1)} < 0.$$

Hence f is nonincreasing, so that (44) has been proved. Step 3. We show $\phi_{qk+s} \leq \phi_{qk+s+1}$ for any $s \geq 1$ and $q \neq 0$. (i) When $1 \leq s \leq r-1$, we see from the first case of (41) and the second case of (42) that

$$\phi_{qk+s+1} - \phi_{qk+s} = \sum_{m=q+1}^{p-q} \frac{1}{mk(mk+1)} \geq 0, \quad (48)$$

where the summation equals 0 when $p-q < q+1$. (ii) When $s = r$, it follows from the second case of (41) and the second case of (42) that

$$\phi_{qk+s+1} - \phi_{qk+s} = \sum_{m=q+1}^{p-q-1} \frac{1}{mk(mk+1)} + \frac{1}{(p-q)k} \geq 0. \quad (49)$$

(iii) When $s > r$, it follows from the third case of (41) and the second case of (42) that

$$\phi_{qk+s+1} - \phi_{qk+s} = \sum_{m=q+1}^{p-q-1} \frac{1}{mk(mk+1)} \geq 0. \quad (50)$$

Therefore $\phi_{qk+s+1} \geq \phi_{qk+s}$. \square

Theorem 8. When $2 \leq k \leq n/2$, the maximum points of $\{\phi_i\}$ are $i = k$ and $i = n - k + 1$.

Proof. By Theorem 7, the maximum value is attained by either $i = qk$. In the case of $r = 0$, it follows from the second case of (45) and (50) that

$$\phi_{(q+1)k} - \phi_{qk} = \sum_{s=0}^{k-1} (\phi_{qk+s+1} - \phi_{qk+s}) = k \sum_{m=q+1}^{p-q-1} \frac{1}{mk(mk+1)} + \frac{1}{(p-q)k} - \frac{1}{qk+1}. \quad (51)$$

Here, remark that the summations in (45) and (48) are taken from $m = q+1$ to not $p-q-1$ but $p-q$. So, in the case of $r > 0$, it follows from the first case of (45), (48), (49), and (50) that

$$\begin{aligned} \phi_{(q+1)k} - \phi_{qk} &= \sum_{s=0}^{k-1} (\phi_{qk+s+1} - \phi_{qk+s}) \\ &= k \sum_{m=q+1}^{p-q-1} \frac{1}{mk(mk+1)} + \frac{r}{(p-q)k\{(p-q)k+1\}} + \frac{1}{(p-q)k} - \frac{1}{qk+1}. \end{aligned} \quad (52)$$

Since (52) reduces to (51) when $r = 0$, (52) is correct for $r = 0$.

For $1 \leq q$, $2q+2 \leq p$, $0 \leq r \leq k-1$, and $2 \leq k$, let $f_1(p, q, r, k) := \phi_{(q+1)k} - \phi_{qk}$. Then, it is obvious that $f_1(p, q, r, k) \leq f_1(p, q, k-1, k) =: f_2(p, q, k)$ for any $0 \leq r \leq k-1$. Further,

$$f_2(p+1, q, k) - f_2(p, q, k) = \frac{-(k-1)}{(p-q+1)\{(p-q+1)k+1\}\{(p-q)k+1\}} < 0.$$

Hence $f_2(p, q, k)$ is the strict decreasing w.r.t. p . So, let $f_3(q, k) := f_2(2q+2, q, k)$. Then

$$\begin{aligned} f_3(q, k) &= \frac{1}{(q+1)\{(q+1)k+1\}} + \frac{k-1}{(q+2)k\{(q+2)k+1\}} + \frac{1}{(q+2)k} - \frac{1}{qk+1} \\ &= \frac{-2q^2k^2 + 2qk^2 - 5qk^2 + 3qk - 4k^2 + 2q + k + 3}{(q+1)(q+2)(qk+1)(qk+k+1)(qk+2k+1)}. \end{aligned}$$

Since the numerator of the right-hand side is expressed as

$$-(k-1) \left\{ 2k \left(q + \frac{5k+2}{4k} \right)^2 - \frac{(5k+2)^2}{8k} + 4k + 3 \right\},$$

the maximum value of $f_3(q, k)$ on $q \geq 1$ is attained by $q = 1$. Then the numerator of $f_3(1, k)$ is $-(k-1)(11k+5) < 0$. So, $f_1(p, q, k, r) \leq f_2(p, q, k) \leq f_3(q, k) < 0$ as desired. \square

REFERENCES

- [1] Aumann, R. J., and Hart, S., Editors, *Handbook of game theory with Economic Applications Vol. 1*, Elsevier, Amsterdam, (1992).
- [2] Gelfand, I. M., and Fomin, S. V., *Calculus of variations*. Prentice Hall, (1963).
- [3] Kawasaki, H., *Conjugate points for a nonlinear programming problem with constraints*, Journal of Nonlinear and Convex Analysis 1, (2000) 87–293.
- [4] Kawasaki, H., *A conjugate points theory for a nonlinear programming problem*, SIAM Journal Control and Optimization 40, (2001) 54–63.
- [5] Kawasaki, H., *Analysis of conjugate points for constant tridiagonal Hesse matrices of a class of extremal problems*, Optimization Methods and Software 18, (2003) 197–205.
- [6] Kawasaki H., *A game-theoretic aspect of conjugate sets for a nonlinear programming problem*, Proceedings of the third International Conference on Nonlinear Analysis and Convex Analysis, Yokohama Publishers, (2004) 159–168.
- [7] Kawasaki H., *Conjugate-set game for a nonlinear programming problem*, in *Game theory and applications 10*, eds. L.A. Petrosjan and V.V. Mazalov, Nova Science Publishers, New York, USA, (2005) 87–95.

TAKEAKI FUCHIKAMI

Graduate school of Mathematics, Kyushu University 33, Fukuoka 812-8581, Japan

E-mail address: ma204047@math.kyushu-u.ac.jp

HIDEFUMI KAWASAKI

Faculty of Mathematics, Kyushu University 33, Fukuoka 812-8581, Japan

E-mail address: kawasaki@math.kyushu-u.ac.jp

List of MHF Preprint Series, Kyushu University

21st Century COE Program

Development of Dynamic Mathematics with High Functionality

- MHF2003-1 Mitsuhiro T. NAKAO, Kouji HASHIMOTO & Yoshitaka WATANABE
A numerical method to verify the invertibility of linear elliptic operators with applications to nonlinear problems
- MHF2003-2 Masahisa TABATA & Daisuke TAGAMI
Error estimates of finite element methods for nonstationary thermal convection problems with temperature-dependent coefficients
- MHF2003-3 Tomohiro ANDO, Sadanori KONISHI & Seiya IMOTO
Adaptive learning machines for nonlinear classification and Bayesian information criteria
- MHF2003-4 Kazuhiro YOKOYAMA
On systems of algebraic equations with parametric exponents
- MHF2003-5 Masao ISHIKAWA & Masato WAKAYAMA
Applications of Minor Summation Formulas III, Plücker relations, Lattice paths and Pfaffian identities
- MHF2003-6 Atsushi SUZUKI & Masahisa TABATA
Finite element matrices in congruent subdomains and their effective use for large-scale computations
- MHF2003-7 Setsuo TANIGUCHI
Stochastic oscillatory integrals - asymptotic and exact expressions for quadratic phase functions -
- MHF2003-8 Shoki MIYAMOTO & Atsushi YOSHIKAWA
Computable sequences in the Sobolev spaces
- MHF2003-9 Toru FUJII & Takashi YANAGAWA
Wavelet based estimate for non-linear and non-stationary auto-regressive model
- MHF2003-10 Atsushi YOSHIKAWA
Maple and wave-front tracking — an experiment
- MHF2003-11 Masanobu KANEKO
On the local factor of the zeta function of quadratic orders
- MHF2003-12 Hidefumi KAWASAKI
Conjugate-set game for a nonlinear programming problem

- MHF2004-1 Koji YONEMOTO & Takashi YANAGAWA
Estimating the Lyapunov exponent from chaotic time series with dynamic noise
- MHF2004-2 Rui YAMAGUCHI, Eiko TSUCHIYA & Tomoyuki HIGUCHI
State space modeling approach to decompose daily sales of a restaurant into time-dependent multi-factors
- MHF2004-3 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA
Cubic pencils and Painlevé Hamiltonians
- MHF2004-4 Atsushi KAWAGUCHI, Koji YONEMOTO & Takashi YANAGAWA
Estimating the correlation dimension from a chaotic system with dynamic noise
- MHF2004-5 Atsushi KAWAGUCHI, Kentarou KITAMURA, Koji YONEMOTO, Takashi YANAGAWA & Kiyofumi YUMOTO
Detection of auroral breakups using the correlation dimension
- MHF2004-6 Ryo IKOTA, Masayasu MIMURA & Tatsuyuki NAKAKI
A methodology for numerical simulations to a singular limit
- MHF2004-7 Ryo IKOTA & Eiji YANAGIDA
Stability of stationary interfaces of binary-tree type
- MHF2004-8 Yuko ARAKI, Sadanori KONISHI & Seiya IMOTO
Functional discriminant analysis for gene expression data via radial basis expansion
- MHF2004-9 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA
Hypergeometric solutions to the q -Painlevé equations
- MHF2004-10 Raimundas VIDŪNAS
Expressions for values of the gamma function
- MHF2004-11 Raimundas VIDŪNAS
Transformations of Gauss hypergeometric functions
- MHF2004-12 Koji NAKAGAWA & Masakazu SUZUKI
Mathematical knowledge browser
- MHF2004-13 Ken-ichi MARUNO, Wen-Xiu MA & Masayuki OIKAWA
Generalized Casorati determinant and Positon-Negaton-Type solutions of the Toda lattice equation
- MHF2004-14 Nalini JOSHI, Kenji KAJIWARA & Marta MAZZOCCO
Generating function associated with the determinant formula for the solutions of the Painlevé II equation

- MHF2004-15 Kouji HASHIMOTO, Ryohei ABE, Mitsuhiro T. NAKAO & Yoshitaka WATANABE
Numerical verification methods of solutions for nonlinear singularly perturbed problem
- MHF2004-16 Ken-ichi MARUNO & Gino BIONDINI
Resonance and web structure in discrete soliton systems: the two-dimensional Toda lattice and its fully discrete and ultra-discrete versions
- MHF2004-17 Ryuei NISHII & Shinto EGUCHI
Supervised image classification in Markov random field models with Jeffreys divergence
- MHF2004-18 Kouji HASHIMOTO, Kenta KOBAYASHI & Mitsuhiro T. NAKAO
Numerical verification methods of solutions for the free boundary problem
- MHF2004-19 Hiroki MASUDA
Ergodicity and exponential β -mixing bounds for a strong solution of Lévy-driven stochastic differential equations
- MHF2004-20 Setsuo TANIGUCHI
The Brownian sheet and the reflectionless potentials
- MHF2004-21 Ryuei NISHII & Shinto EGUCHI
Supervised image classification based on AdaBoost with contextual weak classifiers
- MHF2004-22 Hideki KOSAKI
On intersections of domains of unbounded positive operators
- MHF2004-23 Masahisa TABATA & Shoichi FUJIMA
Robustness of a characteristic finite element scheme of second order in time increment
- MHF2004-24 Ken-ichi MARUNO, Adrian ANKIEWICZ & Nail AKHMEDIEV
Dissipative solitons of the discrete complex cubic-quintic Ginzburg-Landau equation
- MHF2004-25 Raimundas VIDŪNAS
Degenerate Gauss hypergeometric functions
- MHF2004-26 Ryo IKOTA
The boundedness of propagation speeds of disturbances for reaction-diffusion systems
- MHF2004-27 Ryusuke KON
Convex dominates concave: an exclusion principle in discrete-time Kolmogorov systems

- MHF2004-28 Ryusuke KON
Multiple attractors in host-parasitoid interactions: coexistence and extinction
- MHF2004-29 Kentaro IHARA, Masanobu KANEKO & Don ZAGIER
Derivation and double shuffle relations for multiple zeta values
- MHF2004-30 Shuichi INOKUCHI & Yoshihiro MIZOGUCHI
Generalized partitioned quantum cellular automata and quantization of classical CA
- MHF2005-1 Hideki KOSAKI
Matrix trace inequalities related to uncertainty principle
- MHF2005-2 Masahisa TABATA
Discrepancy between theory and real computation on the stability of some finite element schemes
- MHF2005-3 Yuko ARAKI & Sadanori KONISHI
Functional regression modeling via regularized basis expansions and model selection
- MHF2005-4 Yuko ARAKI & Sadanori KONISHI
Functional discriminant analysis via regularized basis expansions
- MHF2005-5 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA
Point configurations, Cremona transformations and the elliptic difference Painlevé equations
- MHF2005-6 Kenji KAJIWARA, Tetsu MASUDA, Masatoshi NOUMI, Yasuhiro OHTA & Yasuhiko YAMADA
Construction of hypergeometric solutions to the q Painlevé equations
- MHF2005-7 Hiroki MASUDA
Simple estimators for non-linear Markovian trend from sampled data:
I. ergodic cases
- MHF2005-8 Hiroki MASUDA & Nakahiro YOSHIDA
Edgeworth expansion for a class of Ornstein-Uhlenbeck-based models
- MHF2005-9 Masayuki UCHIDA
Approximate martingale estimating functions under small perturbations of dynamical systems
- MHF2005-10 Ryo MATSUZAKI & Masayuki UCHIDA
One-step estimators for diffusion processes with small dispersion parameters from discrete observations
- MHF2005-11 Junichi MATSUKUBO, Ryo MATSUZAKI & Masayuki UCHIDA
Estimation for a discretely observed small diffusion process with a linear drift

- MHF2005-12 Masayuki UCHIDA & Nakahiro YOSHIDA
AIC for ergodic diffusion processes from discrete observations
- MHF2005-13 Hiromichi GOTO & Kenji KAJIWARA
Generating function related to the Okamoto polynomials for the Painlevé IV equation
- MHF2005-14 Masato KIMURA & Shin-ichi NAGATA
Precise asymptotic behaviour of the first eigenvalue of Sturm-Liouville problems with large drift
- MHF2005-15 Daisuke TAGAMI & Masahisa TABATA
Numerical computations of a melting glass convection in the furnace
- MHF2005-16 Raimundas VIDŪNAS
Normalized Leonard pairs and Askey-Wilson relations
- MHF2005-17 Raimundas VIDŪNAS
Askey-Wilson relations and Leonard pairs
- MHF2005-18 Kenji KAJIWARA & Atsushi MUKAIHIRA
Soliton solutions for the non-autonomous discrete-time Toda lattice equation
- MHF2005-19 Yuu HARIYA
Construction of Gibbs measures for 1-dimensional continuum fields
- MHF2005-20 Yuu HARIYA
Integration by parts formulae for the Wiener measure restricted to subsets in \mathbb{R}^d
- MHF2005-21 Yuu HARIYA
A time-change approach to Kotani's extension of Yor's formula
- MHF2005-22 Tadahisa FUNAKI, Yuu HARIYA & Mark YOR
Wiener integrals for centered powers of Bessel processes, I
- MHF2005-23 Masahisa TABATA & Satoshi KAIZU
Finite element schemes for two-fluids flow problems
- MHF2005-24 Ken-ichi MARUNO & Yasuhiro OHTA
Determinant form of dark soliton solutions of the discrete nonlinear Schrödinger equation
- MHF2005-25 Alexander V. KITAEV & Raimundas VIDŪNAS
Quadratic transformations of the sixth Painlevé equation
- MHF2005-26 Toru FUJII & Sadanori KONISHI
Nonlinear regression modeling via regularized wavelets and smoothing parameter selection

- MHF2005-27 Shuichi INOKUCHI, Kazumasa HONDA, Hyen Yeal LEE, Tatsuro SATO,
Yoshihiro MIZOGUCHI & Yasuo KAWAHARA
On reversible cellular automata with finite cell array
- MHF2005-28 Toru KOMATSU
Cyclic cubic field with explicit Artin symbols
- MHF2005-29 Mitsuhiro T. NAKAO, Kouji HASHIMOTO & Kaori NAGATOU
A computational approach to constructive a priori and a posteriori error
estimates for finite element approximations of bi-harmonic problems
- MHF2005-30 Kaori NAGATOU, Kouji HASHIMOTO & Mitsuhiro T. NAKAO
Numerical verification of stationary solutions for Navier-Stokes problems
- MHF2005-31 Hidefumi KAWASAKI
A duality theorem for a three-phase partition problem
- MHF2005-32 Hidefumi KAWASAKI
A duality theorem based on triangles separating three convex sets
- MHF2005-33 Takeaki FUCHIKAMI & Hidefumi KAWASAKI
An explicit formula of the Shapley value for a cooperative game induced from
the conjugate point