

Binding through coupling to a radiation field*

Fumio Hiroshima

平成 18 年 2 月 13 日

1 Introduction

1.1 Definition

This is a joint work with H. Spohn¹. We consider a system of one electron interacting with a quantized radiation field. In particular we investigate the so called *Pauli-Fierz* [13] model². Although the Pauli-Fierz model is a nonrelativistic model, it correctly describes the interaction between low energy electrons and photons in a sense. Actually the Lamb shift and gyromagnetic ratio shift were described by using the Pauli-Fierz model. See [2, 14, 12].

In this paper we take the dipole approximation for simplicity. Moreover we suppose that the electron is spinless, moves in the d -dimensional space, and has the $d - 1$ transverse degrees of freedom. Throughout this paper we assume

$$d \geq 3.$$

The Hamiltonian of the system is of the form

$$H(\alpha) = \frac{1}{2m} (p \otimes I - \alpha I \otimes A)^2 + V \otimes I + I \otimes H_f \quad (1.1)$$

*賺 1208

¹ Zentrum Mathematik, Technische Universität München, Archisstraße 21, D-80290, München, Germany. e-mail: spohn@mathematik.tu-muenchen.de

² See [10] for recent advances of the Pauli-Fierz model.

acting on the Hilbert space

$$\mathcal{H} := L^2(\mathbb{R}^d) \otimes \mathcal{F}_{\text{EM}}.$$

Here \mathcal{F}_{EM} denotes the Boson Fock space over $W := \bigoplus^{d-1} L^2(\mathbb{R}^d)$

$$\mathcal{F}_{\text{EM}} := \bigoplus_{n=0}^{\infty} [\otimes_s^n W],$$

where $\otimes_s^n W$ denotes the n -fold symmetric tensor product of W with $\otimes_s^0 W := \mathbb{C}$. m is the bare mass of the electron and α a coupling constant. We adopt the unit $\hbar = 1 = c$. Then $\alpha \approx \sqrt{137}$. $p = -i\vec{\nabla}$ is the momentum operator canonically conjugate to the position operator x in $L^2(\mathbb{R}^d)$, and $V = V(x)$ an external potential for which precise conditions will be specified below. The smeared radiation field is defined by

$$A_\mu := \sum_{r=1}^{d-1} \frac{1}{\sqrt{2}} \int e_\mu^r(k) \left\{ \frac{\hat{\varphi}(-k)}{\sqrt{(2\pi)^d \omega(k)}} a^{\dagger r}(k) + \frac{\hat{\varphi}(k)}{\sqrt{(2\pi)^d \omega(k)}} a^r(k) \right\} d^d k,$$

and the free Hamiltonian by

$$H_f := \sum_{r=1}^{d-1} \int \omega(k) a^{\dagger r}(k) a^r(k) dk,$$

where the dispersion relation is given by

$$\omega(k) := |k|.$$

$a^{\dagger r}(k)$ and $a^r(k)$ denote the annihilation and creation operators, respectively. They satisfy the canonical commutation relations,

$$[a^r(k), a^{\dagger s}(k')] = \delta_{rs} \delta(k - k'), \quad [a^r(k), a^s(k')] = [a^{\dagger r}(k), a^{\dagger s}(k')] = 0.$$

The vectors, $e^r(k) = (e_1^r(k), \dots, e_d^r(k))$, $r = 1, \dots, d-1$, denote polarization vectors satisfying

$$e^r(k) \cdot e^s(k) = \delta_{rs}, \quad k \cdot e^r(k) = 0.$$

Finally $\hat{\varphi}$ denotes a form factor serving as an ultraviolet cutoff. We assume that

$$\hat{\varphi}/\sqrt{\omega} \in L^2(\mathbb{R}^d), \tag{1.2}$$

and

$$\overline{\hat{\varphi}(k)} = \hat{\varphi}(-k). \quad (1.3)$$

(1.2) and (1.3) ensure that $H(\alpha)$ is a well defined symmetric operator in \mathcal{H} . It is known that

$$\text{Spec}(H_f) = [0, \infty)$$

and

$$\text{Spec}_p(H_f) = \{0\}.$$

The multiplicity of $\{0\}$ is one, and

$$H_f \Omega = 0,$$

where $\Omega := 1 \oplus 0 \oplus 0 \oplus \dots$ is the Fock vacuum in \mathcal{F}_{EM} .

1.2 Problems

Suppose that V is relatively bounded with respect to $-\Delta$ with a sufficiently small relative bound. Then it is proven [8] that $H(\alpha)$ is self-adjoint on $D(\Delta \otimes I) \cap D(I \otimes H_f)$ and bounded from below for arbitrary couplings. Moreover by investigating the integral kernel of $e^{-tH(\alpha)}$, $t \geq 0$, the uniqueness of the ground state, if it exists, is established in [6]³.

In the case when $-\frac{1}{2m}\Delta + V$ has the positive spectral gap,

$$\inf \text{Spec}_{\text{ess}}\left(-\frac{1}{2m}\Delta + V\right) - \inf \text{Spec}\left(-\frac{1}{2m}\Delta + V\right) > 0,$$

the existence of the ground state of the full Pauli-Fierz Hamiltonian is established in [3, 5, 9, 4]. In particular, Bach, Fröhlich and Sigal [3] proved it under *no* assumption of infrared cutoff condition⁴ but sufficiently weak couplings. For arbitrary couplings, it is established in [4] due to Griesemer, Lieb and Loss.

³ For the *full* Pauli-Fierz Hamiltonian, self-adjointness and the uniqueness of the ground state are established in [8] and [6], respectively.

⁴ The condition $\int_{\mathbb{R}^d} |\hat{\varphi}(k)|^2 / \omega(k)^3 dk < \infty$ is called the *infrared cutoff* condition. In the case of $d = 3$ this condition implies $0 = \hat{\varphi}(0) = (2\pi)^{-3/2} \int \varphi(x) dx$, i.e., physically the electron charge turns out to be zero!


The main purpose of this paper is to prove the existence of the ground state of $H(\alpha)$ under *no* assumption of the positive spectral gap. In the zero spectral gap case, $-\frac{1}{2m}\Delta + V$ may have no ground state. That is, we show that strong couplings produce the ground state. The physical reasoning behind such a result is as follows. As the electron binds photons it acquires the effective mass

$$m \rightarrow m + \delta m(\alpha^2)$$

which is increasing in $|\alpha|$. Roughly speaking $H(\alpha)$ may be replaced by

$$H(\alpha) \sim -\frac{1}{2(m + \delta m(\alpha^2))}\Delta + V, \quad (1.4)$$

and, for the sufficiently large $|\alpha|$, the right hand side of (1.4) may have ground states. Needless to say (1.4) has no sharp mathematical meaning, we show, however, the associated phenomena in this paper.




⊠ 1: $H(0)$

This paper is organized as follows. In Section 2 we prove the binding. In Section 3 we give some examples of the external potentials. Finally in Section 4 we give some remarks.

2 Binding

We suppose the following assumptions on V .



☒ 2: $H(\alpha)$

- (1) $\|Vf\| \leq a\|\Delta f\| + b\|f\|$ for $f \in D(\Delta)$ with sufficiently small $a \geq 0$, and positive $b \geq 0$.
- (2) $V \in C^1(\mathbb{R}^d)$ and $\partial_\mu V \in L^\infty(\mathbb{R}^d)$, $\mu = 1, \dots, d$.
- (3) There exist $\mu_0 \geq 1$ and $r_0 > 0$ such that for all $\eta > \mu_0$

$$\inf \text{Spec}\left(-\frac{1}{2m}\Delta + \eta V\right) \leq -r_0,$$

and

$$\text{Spec}_{\text{ess}}\left(-\frac{1}{2m}\Delta + \eta V\right) = [0, \infty).$$

It is of interest to investigate sufficiently shallow external potentials. Since $d \geq 3$, for such a shallow V , $-\frac{1}{2m}\Delta + V$ may have no ground state. If $-\frac{1}{2m}\Delta + V$ has no ground state, then the decoupled Hamiltonian

$$H(\alpha = 0) = \left(-\frac{1}{2m}\Delta + V\right) \otimes I + I \otimes H_f$$

also has no ground state.

For later use we define the dilatation unitary of $L^2(\mathbb{R}^d)$ by

$$D(\kappa)f(k) := \kappa^{d/2}f(k/\kappa),$$

where $\kappa > 0$ denotes the scaling parameter. The scaled Hamiltonian is defined by

$$H(\alpha, \kappa)$$

$$\begin{aligned}
& := \kappa^2 D(\kappa)^{-1} \left\{ \frac{1}{2m} (p \otimes I - \alpha I \otimes A)^2 + I \otimes H_f + \frac{1}{\kappa^2} V(x/\kappa) \otimes I \right\} D(\kappa) \\
& = \frac{1}{2m} (p \otimes I - \kappa \alpha I \otimes A)^2 + V \otimes I + \kappa^2 I \otimes H_f.
\end{aligned}$$

We suppose the following technical assumptions on $\hat{\varphi}$.

- (1) $\hat{\varphi}(k) = \hat{\varphi}(|k|)$.
- (2) $\omega^{n/2} \hat{\varphi} \in L^2(\mathbb{R}^d)$ for $n = -5, -4, -3, -2, -1, 0, 1, 2$.
- (3) $|\hat{\varphi}(\sqrt{s})| s^{(d-1)/2} \in L^\epsilon([0, \infty), ds)$, $0 < \epsilon < 1$, and is Lipschitz continuous of order strictly less than one.
- (4) $\|\hat{\varphi} \omega^{(d-2)/2}\|_\infty < \infty$ and $\|\hat{\varphi} \omega^{(d-1)/2}\|_\infty < \infty$.
- (5) $\hat{\varphi}(k) \neq 0$ for all $k \neq 0$.

Thus (1)–(5) ensure the following lemmas⁵.

Lemma 2.1 *There exist the unitary operator $U(\kappa)$ such that*

$$U(\kappa)^{-1} H(\alpha, \kappa) U(\kappa) = H_{\text{eff}} + \kappa^2 H_f + \kappa^2 \alpha^2 g + \delta V,$$

where

$$H_{\text{eff}} := -\frac{1}{2m_{\text{eff}}} \Delta + V,$$

$$m_{\text{eff}} = m_{\text{eff}}(\alpha^2) := m + \alpha^2 \left(\frac{d-1}{d} \right) \|\hat{\varphi}/\omega\|^2,$$

and

$$g := \frac{d-1}{2\pi} \int_{-\infty}^{\infty} \frac{t^2 \|\hat{\varphi}/(t^2 + \omega^2)\|^2}{m + \alpha^2 \left(\frac{d-1}{d} \right) \|\hat{\varphi}/\sqrt{t^2 + \omega^2}\|^2} dt.$$

Moreover

$$\delta V = \delta V(\alpha, \kappa) := U(\kappa)^{-1} (V \otimes I) U(\kappa) - V \otimes I.$$

⁵ See [1, 11] for details.

Lemma 2.2 *We have*

$$-\frac{D(\alpha)}{\kappa}(H_f + I) \leq \delta V \leq \frac{D(\alpha)}{\kappa}(H_f + I)$$

in the sense of form, where $D(\alpha)$ is a real number satisfying

$$\lim_{|\alpha| \rightarrow \infty} D(\alpha) = 0.$$

Let

$$\alpha_{\text{critical}} := \sqrt{m(\mu_0 - 1)} \sqrt{\frac{d}{d-1}} \|\hat{\varphi}/\omega\|^{-1}.$$

We see

$$H_{\text{eff}} = \frac{m}{m_{\text{eff}}} \left(-\frac{1}{2m} \Delta + \frac{m_{\text{eff}}}{m} V \right).$$

Then, in the case of $|\alpha| > \alpha_{\text{critical}}$, it follows that

$$\inf_x V(x) \leq \inf \text{Spec}(H_{\text{eff}}) < -r_0 \frac{m}{m_{\text{eff}}}.$$

In particular the ground states of H_{eff} exist.

Theorem 2.3 *Let $\kappa = 1$. There exists $\alpha_* > \alpha_{\text{critical}}$ such that for all $|\alpha| \geq \alpha_*$ the ground state of $H(\alpha)$ exists and it is unique.*

Proof: Let N be the number operator in \mathcal{F}_{EM} and $0 < \nu$. By a momentum lattice approximation we see that $H(\alpha) + \nu N$ has the normalized ground state Φ_ν . Let E_I denote the spectral projection of H_{eff} to $I \subset \mathbb{R}$ and P_Ω the projection to Ω . Let $P = E_{(-\infty, -r_0 m/m_{\text{eff}})} \otimes P_\Omega$ and $\Sigma := \inf \text{Spec}(H_{\text{eff}})$. Then we can see that

$$(\Phi_\nu, P\Phi_\nu) \geq 1 - \left(\frac{|\alpha|\epsilon}{m_{\text{eff}}} \right)^2 - \frac{D(\alpha)/2}{|\Sigma| - D(\alpha)/2} \quad (2.1)$$

with some constant ϵ . Note that

$$\lim_{|\alpha| \rightarrow 0} \frac{|\alpha|}{m_{\text{eff}}(\alpha^2)} = 0$$

and

$$\lim_{|\alpha| \rightarrow 0} \Sigma = \inf_x V(x).$$

Thus for sufficiently large $|\alpha|$ the right hand side of (2.1) is strictly positive. Take a subsequence ν' such that $\Phi_{\nu'} \rightarrow \Phi$ as $\nu \rightarrow 0$ weakly. Since P is a finite rank operator, $P\Phi_{\nu'}$ strongly converges to $P\Phi$ and

$$(\Phi, P\Phi) \geq 1 - \left(\frac{|\alpha|\epsilon}{m_{\text{eff}}} \right)^2 - \frac{D(\alpha)/2}{|\Sigma| - D(\alpha)/2}$$

holds. In particular $\Phi \neq 0$. Hence Φ is the ground state. \square

By the assumptions H_{eff} has ground states for $|\alpha| > \alpha_{\text{critical}}$. We have to make sure that $H(\alpha)$ has the same properties.

Theorem 2.4 *We suppose that κ is sufficiently large. Set $V_{\kappa}(x) := \kappa^{-2}V(x/\kappa)$. Then the ground state of*

$$H_{\kappa}(\alpha) = \frac{1}{2m}(p \otimes I - \alpha I \otimes A)^2 + V_{\kappa} \otimes I + I \otimes H_{\text{f}}$$

exists for all $|\alpha| > \alpha_{\text{critical}}$ and it is unique.

Proof: We have

$$H_{\kappa}(\alpha) = \frac{1}{\kappa^2} D(\kappa) H(\alpha, \kappa) D(\kappa)^{-1}.$$

Thus it is enough to prove the existence of the ground states of $H(\alpha, \kappa)$. From the momentum lattice approximation we see that $H(\alpha, \kappa) + \nu N$ has the ground state Φ_{ν} . Moreover we have the inequality

$$(\Phi_{\nu}, P\Phi_{\nu}) \geq 1 - \frac{1}{\kappa^6} \left(\frac{|\alpha|\epsilon}{m_{\text{eff}}} \right)^2 - \frac{1}{\kappa^2} \left(\frac{D(\alpha)/2}{|\Sigma| - D(\alpha)/2} \right).$$

Then the theorem follows in the same way as in Theorem 2.3. \square

3 Example

Suppose that

$$V(x) \leq 0.$$

Let

$$N(V) := a_d \int_{\mathbb{R}^d} |mV(x)|^{d/2} dx,$$

where a_d is a universal constant. The following is known as the Lieb-Thirring equality

$$N(V) = \#\{\text{the nonnegative eigenvalues of } -\frac{1}{2m}\Delta + V\}.$$

Suppose that

$$N(V) < 1.$$

Then $H(0)$ has no ground state, $H(\alpha)$ for sufficiently large $|\alpha|$, however, has the ground state and it is unique by Theorem 2.3.

Remark 3.1 *If $-\frac{1}{2m}\Delta + V$ has the ground state with a positive spectral gap, then $H(\alpha)$ has the ground state for arbitrary $\alpha \in \mathbb{R}$.*

4 Concluding remarks

(1) The full Pauli-Fierz Hamiltonian is defined by

$$H(\alpha) = \frac{1}{2m}(p \otimes I - \alpha A)^2 + V \otimes I + I \otimes H_f.$$

Here under the identification $\mathcal{H} \cong \int_{\mathbb{R}^d}^{\oplus} \mathcal{F}_{\text{EM}} dx$


$$A_\mu := \int_{\mathbb{R}^d}^{\oplus} A_\mu(x) dx,$$

and


$$A_\mu(x) := \sum_{r=1}^{d-1} \frac{1}{\sqrt{2}} \times \\ \times \int e_\mu^r(k) \left\{ \frac{\hat{\varphi}(-k)}{\sqrt{(2\pi)^d \omega(k)}} e^{-ikx} a^{\dagger r}(k) + \frac{\hat{\varphi}(k)}{\sqrt{(2\pi)^d \omega(k)}} e^{ikx} a^r(k) \right\} d^d k.$$

For the full Pauli-Fierz Hamiltonian, it seems to be unknown the binding.

- (2) For α such that $0 < |\alpha| < \alpha_{\text{critical}}$, *no* existence of the ground state is not known.



☒ 3: $H_{\text{Nelson}}(0)$



☒ 4: $H_{\text{Nelson}}(\alpha)$

- (3) The Nelson Hamiltonian with two charged particles is defined by

$$H_{\text{Nelson}}(\alpha) := \left(-\frac{1}{2m} \Delta + V \right) \otimes I + I \otimes H_{\text{N}} + \alpha \phi$$

acting on

$$\mathcal{H} := L^2(\mathbb{R}^d \times \mathbb{R}^d) \otimes \mathcal{F},$$

where \mathcal{F} denotes the Boson Fock space over $L^2(\mathbb{R}^d)$. The free Hamiltonian is defined by

$$H_{\text{N}} := \int \omega(k) a^\dagger(k) a(k) dk$$

and the scalar field by

$$\phi := \int_{\mathbb{R}^d \times \mathbb{R}^d}^{\oplus} \phi(x) dx,$$

$$\phi(x) := \sum_{j=1}^2 \frac{1}{\sqrt{2}} \int \hat{\lambda}(-k) e^{-ikx^j} a^\dagger(k) + \hat{\lambda}(k) e^{ikx^j} a(k) dk.$$

Roughly speaking $H_{\text{Nelson}}(\alpha)$ may be replaced by

$$H_{\text{Nelson}}(\alpha) \sim -\frac{1}{2m} \Delta + V + V_{\text{eff}},$$

$$V_{\text{eff}}(x^1, x^2) = -\frac{\alpha^2}{2} \int_{\mathbb{R}^d} \frac{\hat{\lambda}(k)^2}{\omega(k)} e^{-ik(x^1 - x^2)} dk.$$

Then we can also prove the binding of the Nelson Hamiltonian under certain conditions. We omit details.

参考文献

- [1] A. Arai, Rigorous theory of spectra and radiation for a model in quantum electrodynamics, *J. Math. Phys.* **24** (1983), 1896–1910.
- [2] H. A. Bethe, The electromagnetic shift of energy levels, *Phys. Rev.* **72** (1947), 339–342.
- [3] V. Bach, J. Fröhlich and I. M. Sigal, Spectral analysis for systems of atoms and molecules coupled to the quantized radiation fields, *Commun. Math. Phys.* **207** (1999), 249–290.
- [4] M. Grieseman, E. Lieb and M. Loss, Ground states in non-relativistic quantum electrodynamics, preprint, mp-arc 00–313, 2000.
- [5] F. Hiroshima, Ground states of a model in nonrelativistic quantum electrodynamics I, *J. Math. Phys.* **40** (1999), 6209–6222.
- [6] F. Hiroshima, Ground states of a model in nonrelativistic quantum electrodynamics II, *J. Math. Phys.* **41** (2000), 661–674.
- [7] F. Hiroshima, Essential self-adjointness of translation invariant quantum field models for arbitrary coupling constants, *Commun. Math. Phys.* **211** (2000), 585–613
- [8] F. Hiroshima, The self-adjointness and a relative bound of the Pauli-Fierz Hamiltonian in quantum electrodynamics for arbitrary coupling constants, preprint, mp-ar 01-97, 2001.
- [9] F. Hiroshima, Ground states and spectrum of quantum electrodynamics of non-relativistic particles, (2001), *Trans. Amer. Math. Soc.*
- [10] F. Hiroshima, The analysis of the ground states of atoms interacting with a quantized radiation field, to be published in *International J. Mod. Phys. B*, 2001.
- [11] F. Hiroshima and H. Spohn, Enhanced binding through coupling to a quantum field, preprint, mp-arc 01-39, 2001.

- [12] Z. Koba, Semi-classical treatment of the reactive corrections. I., *Prog. Theoret. Phys.* **4** (1949), 319–330.
- [13] W. Pauli and M. Fierz, Zur Theorie der Emission langwelliger Lichtquanten, *Nuovo Cimento* **15** (1938), 167–188.
- [14] T. Welton, Some observable effects of the quantum-mechanical fluctuations of the electromagnetic field, *Phys. Rev.* **74** (1948), 1157–1167.