

# CAP automorphic representations of low rank groups <sup>\*</sup>

Takuya KONNO <sup>†</sup>

## Abstract

In this talk, I report my recent joint work with K. Konno on non-tempered automorphic representations on low rank groups [KK]. We obtain a fairly complete classification of such automorphic representations for the quasisplit unitary groups in four variables.

## 1 CAP forms

The term CAP in the title is a short hand for the phrase “Cuspidal but Associated to Parabolic subgroups”. This is the name given by Piatetski-Shapiro [PS83] to those cuspidal automorphic representations which apparently contradict the generalized Ramanujan conjecture. More precisely, let  $G$  be a connected reductive group defined over a number field  $F$ , and  $G^*$  be its quasisplit inner form. We write  $\mathbb{A} = \mathbb{A}_F$  for the adèle ring of  $F$ . An irreducible cuspidal representation  $\pi = \bigotimes_v \pi_v$  is a *CAP form* if there exists a residual discrete automorphic representation  $\pi^* = \bigotimes_v \pi_v^*$  such that, at all but finite number of  $v$ ,  $\pi_v$  and  $\pi_v^*$  share the same absolute values of Hecke eigenvalues.

It is a consequence of the result of Jacquet-Shalika [JS81a], [JS81b] and Mœglin-Waldspurger [MW89] that there are no CAP forms on the general linear groups. On the other hand, for a central division algebra  $D$  of dimension  $n^2$  over  $F^\times$ , the trivial representation of  $D^\times(\mathbb{A})$  is clearly a CAP form which shares the same local component, at any place  $v$  where  $D$  is unramified, with the residual representation  $\mathbb{1}_{GL(n, \mathbb{A})}$ . On the other hand, a quasisplit unitary group  $U_{E/F}(3)$  of 3-variables already have non-trivial CAP forms, which can be obtained as  $\theta$ -lifts of some automorphic characters of  $U_{E/F}(1)$  [GR90], [GR91]. But the first and the most well-known example of CAP forms are the analogues of the  $\theta_{10}$  representation by Howe-Piatetski-Shapiro [Sou88] and the Saito-Kurokawa representations of  $Sp_4$  [PS83]. Also Gan-Gurevich-Jiang obtained very interesting example of CAP forms on the split group of type  $G_2$  [GGJ02].

In any case, the local components of CAP forms at almost all places are non-trivial Langlands quotients by definition, and hence non-tempered in an apparent way. To put such forms into the framework of Langlands’ conjecture, J. Arthur proposed a series of conjectures [Art89].

---

<sup>\*</sup>Talk at the seminar “Groupes réductifs et formes automorphes”, Université Denis Diderot (Paris VII), 6 March, 2003

<sup>†</sup>Graduate School of Mathematics, Kyushu University, 812-8581 Hakozaki, Higashi-ku, Fukuoka, Japan

*E-mail:* takuya@math.kyushu-u.ac.jp

*URL:* <http://knmac.math.kyushu-u.ac.jp/~tkonno/>

The author is partially supported by the Grants-in-Aid for Scientific Research No. 12740019, the Ministry of Education, Science, Sports and Culture, Japan

The conjectural description is through the so-called *A-parameters*, homomorphisms  $\psi$  from the direct product of the hypothetical Langlands group  $\mathcal{L}_F$  of  $F$  with  $SL(2, \mathbb{C})$  to the  $L$ -group  ${}^L G$  of  $G$  [Bor79]:

$$\psi : \mathcal{L}_F \times SL(2, \mathbb{C}) \longrightarrow {}^L G,$$

considered modulo  $\widehat{G}$ -conjugation. We write  $\Psi(G)$  for the set of  $\widehat{G}$ -conjugacy classes of  $A$ -parameters for  $G$ . By restriction, we obtain the local component

$$\psi_v : \mathcal{L}_{F_v} \times SL(2, \mathbb{C}) \rightarrow {}^L G_v$$

of  $\psi$  at each place  $v$ . Here the local Langlands group  $\mathcal{L}_{F_v}$  is defined in [Kot84, §12], and  ${}^L G_v$  is the  $L$ -group of the scalar extension  $G_v = G \otimes_F F_v$ . The local conjecture, among other things, associates to each  $\psi_v$  a finite set  $\Pi_{\psi_v}(G_v)$  of isomorphism classes of irreducible unitarizable representations of  $G(F_v)$ , called an *A-packet*. At all but finite number of  $v$ ,  $\Pi_{\psi_v}(G_v)$  is expected to contain a unique unramified element  $\pi_v^1$ . Using such elements, we can form the global  $A$ -packet associated to  $\psi$

$$\Pi_\psi(G) := \left\{ \bigotimes_v \pi_v \mid \begin{array}{l} \text{(i)} \quad \pi_v \in \Pi_{\psi_v}(G_v), \forall v; \\ \text{(ii)} \quad \pi_v = \pi_v^1, \forall v \end{array} \right\}.$$

Arthur's conjecture predicts the multiplicity of each element in  $\Pi_\psi(G)$  in the discrete spectrum of the right regular representation of  $G(\mathbb{A})$  on  $L^2(G(F)\backslash G(\mathbb{A}))$ . Here  $\mathfrak{A}_G$  is the maximal  $\mathbb{R}$ -vector subgroup in the center of the infinite component  $G(\mathbb{A}_\infty)$  of  $G(\mathbb{A})$ .

We say an  $A$ -parameter  $\psi$  is of *CAP type* if

- (i)  $\psi$  is *elliptic*. This is the condition for  $\Pi_\psi(G)$  to contain an element which occurs in the discrete spectrum.
- (ii)  $\psi|_{SL(2, \mathbb{C})}$  is non-trivial.

According to the conjecture, the CAP automorphic representations of  $G(\mathbb{A})$  is contained in some of the global  $A$ -packets associated to such  $A$ -parameters. In this talk, we shall classify the CAP forms by such parameters along the line of Arthur's conjecture, in the case of the quasisplit unitary group  $U_{E/F}(4)$  of four variables. Although our description of such forms tells nothing about the character relations conjectured in [Art89], it is quite explicit and fairly complete. We hope to apply this to certain analysis of the cohomology of the Shimura variety attached to  $GU_{E/F}(4)$ .

## 2 Parameter consideration

**Global case** Take a quadratic extension  $E/F$  of number fields and write  $\sigma$  for the generator of the Galois group of this extension. Let  $G = G_n := U_{E/F}(n)$  be the quasisplit unitary groups in  $n$  variables associated to  $E/F$ . Later we shall mainly be concerned with the case  $n = 4$ . The  $L$ -group  ${}^L G$  is the semi-direct product of  $\widehat{G} = GL(n, \mathbb{C})$  by the absolute Weil group  $W_F$  of  $F$ , where  $W_F$  acts through  $W_F/W_E \simeq \text{Gal}(E/F)$  by

$$\rho_G(\sigma)g = \text{Ad}(I_n)^t g^{-1}, \quad I_n := \begin{pmatrix} & & & 1 \\ & & -1 & \\ & \ddots & & \\ (-1)^{n-1} & & & \end{pmatrix}.$$

Thus an  $A$ -parameter  $\psi$  for  $G$  is determined by its restriction to  $\mathcal{L}_E \times SL(2, \mathbb{C})$ , which is just a completely reducible representation:

$$\psi|_{\mathcal{L}_E \times SL(2, \mathbb{C})} = \bigoplus_{i=1}^r \varphi_{\Pi_i} \otimes \rho_{d_i}.$$

Here  $\Pi_i$  is an irreducible cuspidal representation of  $GL(m_i, \mathbb{A}_E)$  enjoying the following properties:

- $\sigma(\Pi_i) := \Pi_i \circ \sigma$  is isomorphic to the contragredient  $\Pi_i^\vee$ .
- Its central character  $\omega_{\Pi_i}$  restricted to  $\mathbb{A}^\times$  equals  $\omega_{E/F}^{n-d_i-m_i+1}$ , where  $\omega_{E/F}$  is the quadratic character associated to  $E/F$  by the classfield theory.
- Some condition on the order of its twisted Asai  $L$ -functions at  $s = 1$ .

$\rho_d$  is the  $d$ -dimensional irreducible representation of  $SL(2, \mathbb{C})$ . We note that  $\psi$  is elliptic if and only if its irreducible components  $\varphi_{\Pi_i} \otimes \rho_{d_i}$  are distinct to each other. The  $S$ -group

$$\mathcal{S}_\psi(G) := \pi_0(\text{Cent}(\psi, \widehat{G})/Z(\widehat{G}))$$

is isomorphic to  $(\mathbb{Z}/2\mathbb{Z})^{r-1}$ , where  $\pi_0(\bullet)$  stands for the group of connected components. This plays a central role in the conjectural multiplicity formula.

**Local case** Similar description for the  $A$ -packets of the unitary group  $G = G_n$  associated to a quadratic extension  $E/F$  of local fields is also valid. For each  $A$ -parameter  $\psi$ , we have the associated non-tempered Langlands parameter

$$\phi_\psi : \mathcal{L}_F \ni w \longmapsto \psi\left(w, \begin{pmatrix} |w|_F^{1/2} & 0 \\ 0 & |w|_F^{-1/2} \end{pmatrix}\right) \in {}^L G.$$

Here the “absolute value”  $|\cdot|_F$  on  $\mathcal{L}_F$  is the composite  $|\cdot|_F : \mathcal{L}_F \twoheadrightarrow W_F^{\text{ab}} \xrightarrow{\text{rec}} F^\times \xrightarrow{|\cdot|_F} \mathbb{R}_+^\times$ . (rec denotes the reciprocity map in the local classfield theory.) In Arthur’s conjecture, it was imposed that the  $L$ -packet  $\Pi_{\phi_\psi}(G)$  associated to  $\phi_\psi$  should be contained in  $\Pi_\psi(G)$ . We also have the  $S$ -group  $\mathcal{S}_\psi(G)$  as in the global case. We postulate the following:

**Assumption 2.1.** *There exists a bijection  $\Pi_\psi(G) \ni \pi \longmapsto (\bar{s} \mapsto \langle \bar{s}, \pi \rangle_\psi) \in \Pi(\mathcal{S}_\psi(G))$ . Here  $\Pi(\mathcal{S}_\psi(G))$  is the set of isomorphism classes of irreducible representations of  $\mathcal{S}_\psi(G)$ .*

Now for  $n = 4$ , the possibilities of  $\{(d_i, m_i)\}_i$  for elliptic  $A$ -parameters with non-trivial  $SL(2, \mathbb{C})$ -component are given as follows.

- (1) Stable cases.  $\{(4, 1)\}, \{(2, 2)\}$ .
- (2) Endoscopic cases.
  - (a)  $\{(3, 1), (1, 1)\}$ ;
  - (b)  $\{(2, 1), (1, 2)\}$ ;

- (c)  $\{(2, 1), (2, 1)\}$ ;
- (d)  $\{(2, 1), (1, 1), (1, 1)\}$ .

In the cases (1), (2.a), it follows from Assumption 2.1 that  $\Pi_{\phi_\psi}(G) = \Pi_\psi(G)$ , and we know from [Kon98] that all the contribution of the corresponding global  $A$ -packets belong to the residual spectrum. On the other hand,  $\Pi_\psi(G) \setminus \Pi_{\phi_\psi}(G)$  is expected to be non-empty in the rest cases. We shall use the local  $\theta$ -correspondence to construct the missing members.

### 3 Local $\theta$ -correspondence

**Local Howe duality** First let us recall the local  $\theta$ -correspondence. We consider an  $m$ -dimensional (non-degenerate) hermitian space  $(V, (\cdot, \cdot))$  and  $n$ -dimensional skew-hermitian space  $(W, \langle \cdot, \cdot \rangle)$  over  $E$ . We write  $G(V)$  and  $G(W)$  for the unitary groups of  $V$  and  $W$ , respectively. If we define the symplectic space  $(\mathbb{W}, \langle\langle \cdot, \cdot \rangle\rangle)$  by

$$\mathbb{W} := V \otimes_E W, \quad \langle\langle v \otimes w, v' \otimes w' \rangle\rangle := \frac{1}{2} \text{Tr}_{E/F}[(v, v')\sigma(\langle w, w' \rangle)],$$

Then  $(G(V), G(W))$  form a so-called *dual reductive pair* in the symplectic group  $Sp(\mathbb{W})$  of this symplectic space:

$$\iota_{V,W} : G(V) \times G(W) \ni (g, g') \longmapsto g \otimes g' \in Sp(\mathbb{W}).$$

Fixing a non-trivial character  $\psi_F$  of  $F$ , we have the metaplectic group of  $\mathbb{W}$  which is a central extension

$$1 \longrightarrow \mathbb{C}^1 \longrightarrow Mp_{\psi_F}(\mathbb{W}) \longrightarrow Sp(\mathbb{W}) \longrightarrow 1.$$

This admits a unique Weil representation  $\omega_{\psi_F}$  on which  $\mathbb{C}^1$  acts by the multiplication [RR93]. For each pair  $\underline{\xi} = (\xi, \xi')$  of characters of  $E^\times$  satisfying  $\xi|_{F^\times} = \omega_{E/F}^n$ ,  $\xi'|_{F^\times} = \omega_{E/F}^m$ , we have the corresponding lifting  $\tilde{\iota}_{V,W,\underline{\xi}} : G(V) \times G(W) \rightarrow Mp_{\psi_F}(\mathbb{W})$  of  $\iota_{V,W}$ :

$$\begin{array}{ccc} G(V) \times G(W) & \xrightarrow{\tilde{\iota}_{V,W,\underline{\xi}}} & Mp_{\psi_F}(\mathbb{W}) \\ \parallel & & \downarrow \\ G(V) \times G(W) & \xrightarrow{\iota_{V,W}} & Sp(\mathbb{W}) \end{array}$$

The composite  $\omega_{V,W,\underline{\xi}} := \omega_\psi \circ \tilde{\iota}_{V,W,\underline{\xi}}$  is the *Weil representation* of the dual reductive pair  $(G(V), G(W))$  associated to  $\underline{\xi}$ . It is the product of the Weil representations  $\omega_{W,\xi}$  of  $G(V)$  and  $\omega_{V,\xi'}$  of  $G(W)$ .

We write  $\mathcal{R}(G(V), \omega_{W,\xi})$  for the set of isomorphism classes of irreducible admissible representations of  $G(V)$  which appear as quotients of  $\omega_{W,\xi}$ . For  $\pi_V \in \mathcal{R}(G(V), \omega_{W,\xi})$ , the maximal  $\pi_V$ -isotypic quotient of  $\omega_{V,W,\underline{\xi}}$  is of the form  $\pi_V \otimes \Theta_\xi(\pi_V, W)$  for some smooth representation  $\Theta_\xi(\pi_V, W)$  of  $G(W)$ . Similarly we have  $\mathcal{R}(G(W), \omega_{V,\xi'})$  and  $\Theta_\xi(\pi_W, V)$  for each  $\pi_W \in \mathcal{R}(G(W), \omega_{V,\xi'})$ . The local Howe duality conjecture, which was proved by R. Howe himself if  $F$  is archimedean [How89] and by Waldspurger if  $F$  is a non-archimedean local field of odd residual characteristic [Wal90], asserts the following:

- (i)  $\Theta_{\underline{\xi}}(\pi_V, W)$  (resp.  $\Theta_{\underline{\xi}}(\pi_W, V)$ ) is an admissible representation of finite length of  $G(W)$  (resp.  $G(V)$ ), so that it admits an irreducible quotient.
- (ii) Moreover its irreducible quotient  $\theta_{\underline{\xi}}(\pi_V, W)$  (resp.  $\theta_{\underline{\xi}}(\pi_W, V)$ ) is unique.
- (iii)  $\pi_V \mapsto \theta_{\underline{\xi}}(\pi_V, W), \pi_W \mapsto \theta_{\underline{\xi}}(\pi_W, V)$  are bijections between  $\mathcal{R}(G(V), \omega_{W, \underline{\xi}})$  and  $\mathcal{R}(G(W), \omega_{V, \underline{\xi}'})$  converse to each other.

**Adams' conjecture** A link between the local  $\theta$ -correspondence and  $A$ -packets is given by the following conjecture of J. Adams [Ada89]. Suppose  $n \geq m$ . Then we have an  $L$ -embedding  $i_{V, W, \underline{\xi}} : {}^L G(V) \rightarrow {}^L G(W)$  given by

$$i_{V, W, \underline{\xi}}(g \rtimes w) := \begin{cases} \xi' \xi^{-1}(w) \begin{pmatrix} g & \\ & \mathbf{1}_{n-m} \end{pmatrix} \rtimes w & \text{if } w \in W_E, \\ \begin{pmatrix} & g \\ J_{n-m}^{n-m-1} & \end{pmatrix} \rtimes w_\sigma & \text{if } w = w_\sigma, \end{cases}$$

where  $w_\sigma$  is a fixed element in  $W_F \setminus W_E$  and

$$J_n := \begin{pmatrix} 1 & & & \\ & -1 & & \\ & & \ddots & \\ & & & (-1)^{n-1} \end{pmatrix}$$

Let  $T : SL(2, \mathbb{C}) \rightarrow \text{Cent}(i_{V, W, \underline{\xi}}, \widehat{G}(W))$  be the homomorphism which corresponds to a regular unipotent element in  $\text{Cent}(i_{V, W, \underline{\xi}}, \widehat{G}(W)) \simeq GL(n - m, \mathbb{C})$  (the *tail representation* of  $SL(2, \mathbb{C})$ ). Using this, we define the  $\theta$ -lifting of  $A$ -parameters by

$$\theta_{V, W, \underline{\xi}} : \Psi(G(V)) \ni \psi \mapsto (i_{V, W, \underline{\xi}} \circ \psi^\vee) \cdot T \in \Psi(G(W)).$$

**Conjecture 3.1** ([Ada89] Conj.A). *The local  $\theta$ -correspondence should be subordinated to the map of  $A$ -packets:  $\Pi_\psi(G(V)) \mapsto \Pi_{\theta_{V, W, \underline{\xi}}(\psi)}(G(W))$ .*

Here we have said subordinated because  $\mathcal{R}(G(V), \omega_{W, \underline{\xi}})$  is not compatible with  $A$ -packets, that is,  $\Pi_\psi(G(V)) \cap \mathcal{R}(G(V), \omega_{W, \underline{\xi}})$  is often strictly smaller than  $\Pi_\psi(G(V))$ . But when these two are assured to coincide, we can expect more:

**Conjecture 3.2** ([Ada89] Conj.B). *For  $V, W$  in the stable range, that is, the Witt index of  $W$  is larger than  $m$ , we have*

$$\Pi_{\theta_{V, W, \underline{\xi}}(\psi)}(G(W)) = \bigcup_{V; \dim_E V = m} \theta_{\underline{\xi}}(\Pi_\psi(G(V)), W).$$

Now we note that our situation is precisely that of Conj. 3.2 with  $m = 2$  and  $W = V \oplus -V$ . Moreover, we find that the  $A$ -parameters in the cases (2.b), (2.c), (2.d) in § 2 are exactly those of the form

$$\theta_{V, W, \underline{\xi}}(\psi), \quad \psi \in \Psi(G(V)).$$

**$\varepsilon$ -dichotomy** We explain the construction of the  $A$ -packets when  $F$  is non-archimedean. We need one more ingredient.

**Proposition 3.3** ( $\varepsilon$ -dichotomy). *Suppose  $\dim_E V = 2$  and write  $W_1$  for the hyperbolic skew-hermitian space  $(E^2, \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix})$ . Take an  $L$ -packet  $\Pi$  of  $G_2(F) = G(W)$  and  $\tau \in \Pi$  [Rog90, Ch.11].*

(i)  $\tau \in \mathcal{R}(G(W), \omega_{V, \xi'})$  if and only if

$$\varepsilon(1/2, \Pi \times \xi \xi'^{-1}, \psi_F) \omega_\Pi(-1) \lambda(E/F, \psi_F)^{-2} = \omega_{E/F}(-\det V).$$

Here the  $\varepsilon$ -factor on the right hand side is the standard  $\varepsilon$ -factor for  $G_2$  twisted by  $\xi \xi'^{-1}$  defined by the Langlands-Shahidi theory [Sha90].  $\omega_\Pi$  is the central character of the elements of  $\Pi$  and  $\lambda(E/F, \psi_F)$  is Langlands'  $\lambda$ -factor [Lan70].

(ii) If this is the case, we have  $\theta_\xi(\tau, V) = (\xi^{-1} \xi')_{G(V)} \tau_V^\vee$ . Here  $(\xi^{-1} \xi')_{G(V)}$  denotes the character of  $G(V)$  given by the composite


$$G(V) \xrightarrow{\det} U_{E/F}(1, F) \ni z/\sigma(z) \mapsto \xi^{-1} \xi'(z) \in \mathbb{C}^\times.$$

$\tau_V$  stands for the Jacquet-Langlands correspondent<sup>1</sup> of  $\tau$ .

This is a special case of the  $\varepsilon$ -dichotomy of the local  $\theta$ -correspondence for unitary groups over  $p$ -adic fields, which was proved for general unitary groups (at least for supercuspidal representations) in [HKS96]. But since we need to combine this with our description of the residual spectrum [Kon98], we have to use the Langlands-Shahidi  $\varepsilon$ -factors instead of Piatetski-Shapiro-Rallis's doubling  $\varepsilon$ -factors adopted by them. By this reason, we deduced this proposition from the analogous result for the unitary similitude groups [Har93] combined with the following description of the base change for  $G_2$ .

**Lemma 3.4.** *Let  $\tilde{\pi} = \omega \otimes \pi'$  be an irreducible admissible representation of the unitary similitude group  $GU_{E/F}(2) \simeq (E^\times \times GL(2, F))/\Delta F^\times$ , and write  $\Pi(\tilde{\pi})$  for the associated  $L$ -packet of  $G_2(F)$  consisting of the irreducible components of  $\tilde{\pi}|_{G_2(F)}$ . Then the standard base change of  $\Pi(\tilde{\pi})$  to  $GL(2, E)$  [Rog90, 11.4] is given by  $\omega(\det) \pi'_E$ , where  $\pi'_E$  is the base change lift of  $\pi'$  to  $GL(2, E)$  [Lan80].*

Now we construct the  $A$ -packets. Our construction is summarized in the following picture.



<sup>1</sup>In fact, the Jacquet-Langlands correspondence for unitary groups in two variables is defined only for  $L$ -packets and not for each member of the packets [LL79]. We know that  $\tau \mapsto \tau_V$  certainly defines a bijection between  $\Pi$  and its Jacquet-Langlands correspondent. But we do not specify the bijection explicitly here. See Rem. 3.6 also.

Each  $A$ -parameter of our concern is of the form

$$\psi|_{\mathcal{L}_E \times SL(2, \mathbb{C})} = \psi_1|_{\mathcal{L}_E \times SL(2, \mathbb{C})} \oplus (\xi' \xi^{-1} \otimes \rho_2),$$

where  $\psi_1$  is some  $A$ -parameter for  $G_2$ . Take  $\tau \in \Pi_{\psi_1}(G_2)$  and let  $(V, (, ))$  be the 2-dimensional hermitian space such that the condition of Prop. 3.3 (i) holds. If we write  $\pi_V := \theta_{\xi}(\tau, V) \simeq (\xi \xi'^{-1})_{G(V)} \tau_V^{\vee}$ , then the result of [Kud86] tells us  $\pi_+ := \theta_{\xi}(\pi_V, W_2)$ , ( $\tau \in \Pi_{\psi_1}(G_2)$ ) form the local residual  $L$ -packet  $\Pi_{\phi_{\psi}}(G_4)$ . We now suppose that there exists a Jacquet-Langlands correspondent  $\pi_{V'} \simeq (\xi \xi'^{-1})_{G(V')} \tau_{V'}^{\vee}$  of  $\pi_V$  on the unitary group  $G(V')$  of the other (isometry class of) 2-dimensional hermitian space. Then Prop. 3.3 (i) tells us that  $\pi_{V'} \notin \mathcal{R}(G(V'), \omega_{W_1, \xi})$ . Yet its local  $\theta$ -lifting  $\pi_- := \theta_{\xi}(\pi_{V'}, W_2)$  to the larger group  $G_4(F)$  still exists. This is the so-called *early lift* or the *first occurrence*. Following Conj. 3.2, we define

$$\Pi_{\psi}(G_4) := \{\pi_{\pm} \mid \tau \in \Pi_{\psi}(G_2)\}.$$

This gives sufficiently many members of the packet as predicted by Assumption 2.1.

**Example 3.5.** (i) Suppose  $\Pi_{\psi_1}(G_2)$  is an  $L$ -packet consisting of supercuspidal elements. For  $\tau \in \Pi_{\psi_1}(G_2)$ ,  $\pi_+$  is the Langlands quotient  $J_{P_1}^{G_4}(\xi' \xi^{-1} | \cdot |_E^{1/2} \otimes \tau)$ , where  $P_1$  is a parabolic subgroup with the Levi factor  $R_{E/F} \mathbb{G}_m \times G_2$ . On the other hand the early lift  $\pi_-$  of the supercuspidal  $\tau$  is again supercuspidal. Thus  $\Pi_{\psi}(G_4)$  consists of non-tempered members and supercuspidal elements.

(ii) On the contrary, we take  $\xi = \xi'$  and consider  $\Pi_{\psi_1}(G_2)$  consists of either the Steinberg representation  $\delta_{G_2}$  or the trivial representation  $\mathbb{1}_{G_2}$ .

- $\delta_{G_2}$  lifts to  $\pi_V = \mathbb{1}_{G(V)}$ , where  $V$  is anisotropic.  $\pi_{V'} = \delta_{G_2}$ .  $\pi_+ = J_{P_1}^{G_4}(| \cdot |_E^{1/2} \otimes \delta_{G_2})$  and  $\pi_-$  is an irreducible tempered but not square integrable representation.
- $\mathbb{1}_{G_2}$  lifts to  $\pi_V = \mathbb{1}_{G(V)}$  but  $V$  is hyperbolic this time.  $\pi_{V'}$  is again  $\mathbb{1}_{G(V')}$  but this should be viewed as the Jacquet-Langlands correspondent of the  $A$ -packet  $\{\mathbb{1}_{G(V)}\}$ . We have  $\pi_+ = J_{P_2}^{G_4}(I_{\mathbf{B}}^{GL(2)E}(\mathbb{1} \otimes \mathbb{1}) | \det |_E^{1/2})$ , where  $P_2$  is the so-called Siegel parabolic subgroup with the Levi factor  $GL(2, E)$ . Obviously  $\pi_- = J_{P_1}^{G_4}(| \cdot |_E^{1/2} \otimes \delta_{G_2})$ . This last representation is shared by the two packets considered here.

**Real case** We end this section by some comments on the case  $E/F = \mathbb{C}/\mathbb{R}$ . Similar results are obtained by applying the argument of Adams-Barbasch [AB95]. In fact, the local  $\theta$ -correspondence between unitary groups of the same size is described quite explicitly and in full generality in [Pau98]. Their argument also works in the present case. Let me explain some example.

We write  $G_{p,q} = U(p, q)$ . For a regular integral infinitesimal character  $\lambda = (\lambda_1, \lambda_2)$  for  $G_{1,1}$ , consider the extended  $L$ -packet:

$$\Pi_{\lambda} = \{\delta_{1,1}^+, \delta_{1,1}^-, \delta_{2,0}, \delta_{0,2}\}$$

consisting of the discrete series representation of various  $G_{p,q}$  with the infinitesimal character  $\lambda$ . The subscript  $p, q$  indicates that  $\delta_{p,q}^{\bullet}$  lives on  $G_{p,q}$ . We can write  $\xi' \xi^{-1}(z) = (z/\bar{z})^n$ ,  $\forall z \in \mathbb{C}$  for

some  $n \in \mathbb{Z}$ . An analogue of Prop. 3.3 in the real case asserts that the local  $\theta$ -correspondence under the Weil representation  $\omega_{V,W,\underline{\xi}}$  gives a bijection

$$\theta_{\underline{\xi}} : \Pi_{\lambda} \xrightarrow{\sim} \Pi_{n-\lambda},$$

where  $n - \lambda = (n - \lambda_2, n - \lambda_1)$ .

If  $\lambda$  is sufficiently regular, by which we mean  $|\lambda_i - n| > 1$ , then it is proved by J.-S. Li [Li90] that  $\theta_{\underline{\xi}}(\theta_{\underline{\xi}}(\delta_{1,1}^{\pm}), W_2)$  is a non-tempered cohomological representation  $A_{\mathfrak{q}}(\lambda')$ , where the Levi factor of the  $\theta$ -stable parabolic subalgebra  $\mathfrak{q}$  is  $\mathfrak{u}(1, 1) \oplus \mathfrak{u}(1)^2$ . As for the other elements  $\delta_{p,q} \in \Pi_{n-\lambda}$ ,  $\theta_{\underline{\xi}}(\delta_{p,q}, W_2)$  is a discrete series representation  $A_{\mathfrak{q}}(\lambda')$ . This time  $\mathfrak{q}$  has the Levi factor  $\mathfrak{u}(2) \oplus \mathfrak{u}(1)^2$ . The resulting  $A$ -packet  $\theta_{\underline{\xi}}(\Pi_{n-\lambda})$  is exactly the cohomological  $A$ -packet defined by Adams-Johnson [AJ87].

For the complete list of the packets both in the archimedean and non-archimedean case, see our paper [KK].

One can easily check that the  $S$ -groups in the cases (2.b), (2.c), (2.d) satisfy  $\mathcal{S}_{\psi}(G_4) \simeq \mathcal{S}_{\psi_1}(G_2) \times \mathbb{Z}/2\mathbb{Z}$ . Now we define the bijection in Assumption 2.1 by

- $\langle \bar{s}, \pi_{\pm} \rangle_{\psi} := \langle \bar{s}, \tau \rangle_{\psi_1}$  on  $\bar{s} \in \mathcal{S}_{\psi_1}(G_2)$ ;
- $\langle \cdot, \pi_{\pm} \rangle_{\psi}$  on  $\mathbb{Z}/2\mathbb{Z}$  equals the sign character if  $\pi_{-}$  and trivial character otherwise.

For the other cases, only the first one in this definition is enough to give a complete bijection. This finishes our local task.

**Remark 3.6.** *In the above, we do not mention the definition of the pairing  $\langle \cdot, \cdot \rangle_{\psi_1}$ . There are several choices for this, and we can choose one by fixing a non-trivial character  $\psi_F$  of  $F$  [LL79]. Also we did not specify the correspondence  $\pi_V \mapsto \pi_{V'}$ , which is again a subtle problem. In fact, we need to make a choice of (absolute) transfer factor as in [LL79] which again involves a choice of  $\psi_F$  (appearing in  $\lambda(E/F, \psi_F)$  in the transfer factor). Using this specific transfer, we label the members of endoscopic  $L$ -packets of anisotropic unitary group. The correspondence  $\pi_V \mapsto \pi_{V'}$  can be described in terms of these data, but we do not go into details here.*

## 4 Multiplicity formula

We now go back to the global situation where  $E/F$  is a quadratic extension of number fields. We note that there always exists a homomorphism  $\mathcal{S}_{\psi}(G_4) \ni \bar{s} \mapsto \bar{s}(v) \in \mathcal{S}_{\psi_v}(G_{4,v})$ . We can now state the main result of this talk. Although we treat only the number field case, we believe the result holds also over function fields of one variable over a finite field of odd characteristic.

**Theorem 4.1.** *Let  $\psi$  be an  $A$ -parameter of CAP type for  $G_4 = U_{E/F}(4)$ . As was explained in § 1, we form the global  $A$ -packet  $\Pi_{\psi}(G_4) := \bigotimes_v \Pi_{\psi_v}(G_{4,v})$ . Then the multiplicity  $m(\pi)$  of  $\pi = \bigotimes_v \pi_v \in \Pi_{\psi}(G_4)$  in  $L^2(G(F) \backslash G(\mathbb{A}))$  is given by*

$$m(\pi) = \frac{1}{|\mathcal{S}_{\psi}(G_4)|} \sum_{\bar{s} \in \mathcal{S}_{\psi}(G_4)} \epsilon_{\psi}(\bar{s}) \prod_v \langle \bar{s}(v), \pi_v \rangle_{\psi_v},$$



where the sign character  $\epsilon_\psi$  is defined by

$$\epsilon_\psi = \begin{cases} \text{sgn}_{\mathcal{S}_\psi(G_4)} & \text{if } \psi_1 \text{ is a stable } L\text{-parameter} \\ & \text{and } \varepsilon(1/2, \psi_1 \otimes \xi\xi'^{-1}) = -1, \\ \mathbb{1} & \text{otherwise.} \end{cases}$$

Here  $\varepsilon(s, \psi_1 \otimes \xi\xi'^{-1})$  is the Artin root number attached to  $\psi_1$ , which equals the standard  $\varepsilon$ -function for  $\Pi_{\psi_1}(G_2) \times \xi\xi'^{-1}$ .

The proof divides into two parts. Our local construction together with the global  $\theta$ -correspondence shows that the multiplicity is no less than the right hand side. Note that we also relies on the multiplicity formula of Labesse-Langlands for unitary groups in two variables [LL79], [Rog90]. Then we prove a characterization of the image of such  $\theta$ -lifts by poles of certain  $L$ -functions, which gives the converse inequality. This also shows that all the CAP forms for  $U_{E/F}(4)$  are obtained in the above as the contribution of the  $A$ -packets we constructed. In particular the  $A$ -packets contains the sufficiently many members at least for global purposes, so that our Assumption 2.1 is justified.

## References

- [AB95] Jeffrey Adams and Dan Barbasch. Reductive dual pair correspondence for complex groups. *J. Funct. Anal.*, 132(1):1–42, 1995.
- [Ada89] Jeffrey Adams.  $L$ -functoriality for dual pairs. *Astérisque*, (171-172):85–129, 1989. Orbites unipotentes et représentations, II.
- [AJ87] Jeffrey Adams and Joseph F. Johnson. Endoscopic groups and packets for non-tempered representations. *Composit. Math.*, 64:271–309, 1987.
- [Art89] James Arthur. Unipotent automorphic representations: conjectures. *Astérisque*, (171-172):13–71, 1989. Orbites unipotentes et représentations, II.
- [Bor79] A. Borel. Automorphic  $L$ -functions. In *Automorphic forms, representations and  $L$ -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 2*, pages 27–61. Amer. Math. Soc., Providence, R.I., 1979.
- [GGJ02] Wee Teck Gan, Nadya Gurevich, and Dihua Jiang. Cubic unipotent Arthur parameters and multiplicities of square integrable automorphic forms. *Invent. Math.*, 149:225–265, 2002.
- [GR90] Stephen S. Gelbart and Jonathan D. Rogawski. Exceptional representations and Shimura’s integral for the local unitary group  $U(3)$ . In *Festschrift in honor of I. I. Piatetski-Shapiro on the occasion of his sixtieth birthday, Part I (Ramat Aviv, 1989)*, pages 19–75. Weizmann, Jerusalem, 1990.
- [GR91] Stephen S. Gelbart and Jonathan D. Rogawski.  $L$ -functions and Fourier-Jacobi coefficients for the unitary group  $U(3)$ . *Invent. Math.*, 105(3):445–472, 1991.

- [Har93] Michael Harris.  $L$ -functions of  $2 \times 2$  unitary groups and factorization of periods of Hilbert modular forms. *J. Amer. Math. Soc.*, 6(3):637–719, 1993.
- [HKS96] Michael Harris, Stephen S. Kudla, and William J. Sweet. Theta dichotomy for unitary groups. *J. Amer. Math. Soc.*, 9(4):941–1004, 1996.
- [How89] Roger Howe. Transcending classical invariant theory. *J. Amer. Math. Soc.*, 2:535–552, 1989.
- [JS81a] H. Jacquet and J.A. Shalika. On Euler products and the classification of automorphic forms, I. *Amer. J. Math.*, 103(3):499–558, 1981.
- [JS81b] H. Jacquet and J.A. Shalika. On Euler products and the classification of automorphic forms, II. *Amer. J. Math.*, 103(4):777–815, 1981.
- [KK] Kazuko Konno and Takuya Konno. CAP automorphic representations of  $U_{E/F}(4)$  I. Local  $A$ -packets. Preprint, (Kyushu Univ. 2003 No. 4), downloadable from <http://knmac.math.kyushu-u.ac.jp/~tkonno/>.
- [Kon98] T. Konno. The residual spectrum of  $U(2, 2)$ . *Trans. Amer. Math. Soc.*, 350(4):1285–1358, 1998.
- [Kot84] Robert E. Kottwitz. Stable trace formula: cuspidal tempered terms. *Duke Math. J.*, 51(3):611–650, 1984.
- [Kud86] Stephen S. Kudla. On the local theta-correspondence. *Invent. Math.*, 83(2):229–255, 1986.
- [Lan70] R. P. Langlands. On Artin’s  $L$ -function. *Rice Univ. Studies*, 56:23–28, 1970.
- [Lan80] Robert P. Langlands. *Base change for  $GL(2)$* . Princeton University Press, Princeton, N.J., 1980.
- [Li90] Jian-Shu Li. Theta lifting for unitary representations with nonzero cohomology. *Duke Math. J.*, 61(3):913–937, 1990.
- [LL79] J.-P. Labesse and R. P. Langlands.  $L$ -indistinguishability for  $SL(2)$ . *Canad. J. Math.*, 31(4):726–785, 1979.
- [MW89] C. Mœglin and J.-L. Waldspurger. Le spectre résiduel de  $GL(n)$ . *Ann. Sci. École Norm. Sup. (4)*, 22(4):605–674, 1989.
- [Pau98] Annegret Paul. Howe correspondence for real unitary groups. *J. Funct. Anal.*, 159(2):384–431, 1998.
- [PS83] I. I. Piatetski-Shapiro. On the Saito-Kurokawa lifting. *Invent. Math.*, 71(2):309–338, 1983.
- [Rog90] Jonathan D. Rogawski. *Automorphic representations of unitary groups in three variables*. Princeton University Press, Princeton, NJ, 1990.

- [RR93] R. Ranga Rao. On some explicit formulas in the theory of Weil representation. *Pacific J. Math.*, 157(2):335–371, 1993.
- [Sha90] Freydoon Shahidi. A proof of Langlands’ conjecture on Plancherel measures; complementary series for  $p$ -adic groups. *Ann. of Math. (2)*, 132(2):273–330, 1990.
- [Sou88] David Soudry. The CAP representations of  $GS(4, \mathbb{A})$ . *J. reine angew. Math.*, 383:87–108, 1988.
- [Wal90] J.-L. Waldspurger. Démonstration d’une conjecture de dualité de Howe dans le cas  $p$ -adique,  $p \neq 2$ . In *Festschrift in honor of I. I. Piatetski-Shapiro on the occasion of his sixtieth birthday, Part I (Ramat Aviv, 1989)*, pages 267–324. Weizmann, Jerusalem, 1990.