

DECAY OF CORRELATIONS IN SUSPENSION SEMI-FLOWS OF ANGLE-MULTIPLYING MAPS

MASATO TSUJII

ABSTRACT. We consider suspension semi-flows of angle-multiplying maps on the circle for C^r ceiling functions with $r \geq 3$. Under a C^r generic condition on the ceiling function, we show that there exists a Hilbert space (anisotropic Sobolev space) contained in the L^2 space such that the Perron-Frobenius operator for the time- t -map acts naturally on it and that the essential spectral radius of that action is bounded by the square root of the inverse of the minimum expansion rate. This leads to a precise description on decay of correlations. Furthermore, the Perron-Frobenius operator for the time- t -map is quasi-compact for a C^r open and dense set of ceiling functions.

1. INTRODUCTION

Decay of correlations and related topics for hyperbolic dynamical systems have been studied for more than three decades since the works of Bowen[5], Ruelle[18] and Sinai[19]. For the cases of discrete dynamical systems such as iterations of expanding maps and Anosov diffeomorphisms, exponential decay of correlations for Hölder observables was already known in the early stage of the study [5, 18, 19] and we nowadays have fairly good understanding on the speed of decay and also on the spectra of the Perron-Frobenius operators. (See [2, 3, 10, 11] and the references therein.) On the contrary, for the cases of continuous dynamical systems such as Anosov flows, the corresponding argument is much subtler and our knowledge is less satisfactory at present. A simple reason for the subtleness in the cases of flows is that the time- t -maps of hyperbolic flows are *not* hyperbolic (but partially hyperbolic) as there is no expansion or contraction in the flow direction. In [7], Dolgopyat showed exponential decay of correlations for Anosov flows under the assumption that the stable and unstable foliations are both C^1 and are jointly non-integrable. More recently, Liverani[15] extended the result to Anosov flows preserving contact structures and, in particular, to all geodesic flows in strictly negative curvature. However, even in such cases, we do not know whether the semi-groups of Perron-Frobenius operators are quasi-compact on some appropriate Banach spaces. Neither do we know whether we observe exponential decay of correlations for mixing (or generic) Anosov flows.

The aim of this paper is to study decay of correlations not for Anosov flows but for a class of expanding semi-flows, suspension semi-flows of angle-multiplying maps on the circle, which we would like to view as a simplified model of the Anosov flow. We consider Perron-Frobenius operators for the time- t -maps of such semi-flows and let them act on the anisotropic Sobolev spaces introduced in [3]. Our main result is that, under a C^r generic condition on the ceiling function, the essential spectral

radius of the action is bounded by the square root of the inverse of the minimum expansion rate. This leads to a precise description on decay of correlations, which resembles the results known for hyperbolic discrete dynamical systems[3, 10], and extends the earlier result of M. Pollicott[17] on exponential decay. As a byproduct of our methods, we show that the Perron-Frobenius operators are quasi-compact for a C^r open and dense set of ceiling functions.

Actually a prototype of the argument in this paper has appeared in [1], where a class of volume-expanding hyperbolic endomorphisms, called *fat solenoidal attractors*, were studied. In this paper, we will apply essentially the same idea to analyze the time- t -maps of the class of expanding semi-flows mentioned above. We emphasize that our intension is to display the idea in a simple setting and present this paper as a study for the cases of hyperbolic flows¹⁾. For this reason, we will confine our argument to a rather restrictive setting, though it is not very difficult to extend it to more general classes of expanding semi-flows and transfer operators. (See Remark 1.5.)

We fix integers $\ell \geq 2$ and $r \geq 3$. Let $\tau : S^1 \rightarrow S^1$ be the angle-multiplying map on the circle $S^1 = \mathbb{R}/\mathbb{Z}$ defined by $\tau(x) = \ell x$. Let $C_+^r(S^1)$ be the space of positive-valued C^r functions on S^1 . For each $f \in C_+^r(S^1)$, we consider the subset

$$X_f = \{(x, s) \in S^1 \times \mathbb{R} \mid 0 \leq s < f(x)\}$$

of the cylinder $S^1 \times \mathbb{R}$. The suspension semi-flow $\mathbf{T}_f = \{T_f^t : X_f \rightarrow X_f\}_{t \geq 0}$ of τ is the semi-flow on X_f in which each point on X_f moves right upward (or the s -direction) with the unit speed and, at the instant it reaches to the upper boundary of X_f , it jumps down to the lower boundary with the x -coordinate transferred by τ . (See Figure 1.) The precise expression for its time- t -map is

$$T_f^t(x, s) = (\tau^{n(x, s+t; f)}(x), s + t - f^{(n(x, s+t; f))}(x))$$

where $f^{(n)}(x) = \sum_{i=0}^{n-1} f(\tau^i(x))$ and $n(x, t; f) = \max\{n \geq 0 \mid f^{(n)}(x) \leq t\}$.




FIGURE 1. The semi-flow \mathbf{T}_f

Let $m = m_f$ be the normalization of the restriction of the standard Lebesgue measure on $S^1 \times \mathbb{R}$ to X_f . This is an ergodic invariant probability measure for \mathbf{T}_f . For a point $z = (x, s) \in X_f$ and $t \geq 0$, $E(z, t; f) := \ell^{n(x, s+t; f)}$ is the expansion rate

¹⁾It is possible to apply the idea presented in this paper to Anosov flows with sufficiently smooth stable foliations. But we suspect that we need to overcome essential difficulties to treat more general Anosov flows.

along the orbit of z up to time t . The minimum expansion rate of \mathbf{T}_f is naturally defined by

$$\lambda_{\min}(\mathbf{T}_f) = \lim_{t \rightarrow \infty} \left(\min_{z \in X_f} E(z, t; f) \right)^{1/t}.$$

For functions ψ and φ in $L^2(X_f) = L^2(X_f, m_f)$, we consider the correlation

$$\mathbf{Cor}_t(\psi, \varphi) = \int \psi \cdot \varphi \circ T_f^t dm_f - \left(\int \varphi dm_f \right) \left(\int \psi dm_f \right) \quad \text{for } t \geq 0.$$

Suppose that $\int \psi dm_f = 0$ for simplicity. If the semi-flow \mathbf{T}_f is mixing, we have that $\lim_{t \rightarrow \infty} \mathbf{Cor}_t(\psi, \varphi) = 0$. The question is the rate of convergence in this limit. In [17], M. Pollicott showed, under a mild condition on f , that the rate is exponential: $|\mathbf{Cor}_t(\psi, \varphi)| < \text{const. exp}(-\epsilon t)$ for some $\epsilon > 0$. (See also [4] for a generalization.) Under a C^r generic condition on f , our results give a more precise description on asymptotic behavior of the correlation as $t \rightarrow \infty$: For any real number $\mu > (\lambda_{\min}(\mathbf{T}_f))^{-1/2}$, there exists finitely many complex numbers²⁾ $\lambda_i \in \mathbb{C}$ with $\mu \leq |\lambda_i| < 1$ and integers $m_i \geq 0$ for $1 \leq i \leq k$, such that

$$(1) \quad \left| \mathbf{Cor}_t(\psi, \varphi) - \sum_{i=1}^k H_i(\psi, \varphi) \cdot t^{m_i} \lambda_i^t \right| \leq H_0(\psi, \varphi) \mu^t \quad \text{for } t \geq 0$$

for any $\varphi \in L^2(X_f)$ and any C^1 function ψ with $\int \psi dm_f = 0$ supported on the interior of X_f , where $H_i(\psi, \varphi)$ are coefficients that depend on ψ and φ (and μ). As we will see later, the complex numbers λ_i above are peripheral eigenvalues of the Perron-Frobenius operator for the time-1-map and each integer m_i is bounded by the geometric multiplicity of the eigenvalue λ_i .

In order to state the main results, we introduce some more notation. The differential $(DT_f^t)_z$ of T_f^t at $z \in X_f$ is defined in the usual way if both z and $T^t(z)$ belong to the interior of X_f and, otherwise, is defined by

$$(DT_f^t)_z = \lim_{\epsilon \rightarrow +0} (DT_f^t)_{z+(0,\epsilon)}.$$

Fix³⁾ a real number $\ell^{-1} < \gamma_0 < 1$. Put

$$\theta_f = \max_{x \in S^1} |f'(x)| / (\gamma_0 \ell - 1)$$

and

$$\mathbf{C}_f = \mathbf{C}(\theta_f) := \{(x, y) \in \mathbb{R}^2 \mid |y| \leq \theta_f |x|\}.$$

By the definition of θ_f , we see that the cone \mathbf{C}_f is strictly invariant for \mathbf{T}_f in the sense that

$$(DT_f^t)_z(\mathbf{C}_f) \subset \mathbf{C}(\gamma_0 \theta_f) \subset \mathbf{C}_f \quad \text{for all } z = (x, s) \in X_f \text{ and } t \geq f(x) - s.$$

Note that, for large t , the inverse image $(T_f^t)^{-1}(z)$ of a point $z \in X_f$ consists of many points and thus there are many narrow cones $(DT_f^t)_\zeta(\mathbf{C}_f)$ for $\zeta \in (T_f^t)^{-1}(z)$ in the tangent space at z . As a measure for transversality between such cones, we introduce the quantity

$$\mathbf{m}(f, t) = \max_{z \in X_f} \max_{w \in (T_f^t)^{-1}(z)} \sum_{\zeta: \zeta \ni w} \frac{1}{E(\zeta, t; f)}$$

²⁾The complex numbers λ_i need not be distinct from each other.

³⁾It is certainly better to choose γ_0 close to 1, though it is not necessary.

where $\sum_{\zeta: \zeta \not\sim w}$ is the sum over the points $\zeta \in (T_f^t)^{-1}(z)$ such that

$$(DT_f^t)_\zeta(\mathbf{C}_f) \cap (DT_f^t)_w(\mathbf{C}_f) \neq \{0\}.$$

Note that we always have $\mathbf{m}(f, t) \leq 1$ because

$$(2) \quad \sum_{\zeta \in (T_f^t)^{-1}(z)} \frac{1}{E(\zeta, t; f)} = 1 \quad \text{for any } t \geq 0 \text{ and } z \in X_f.$$

Finally we define the exponent

$$\mathbf{m}(f) = \limsup_{t \rightarrow \infty} \mathbf{m}(f, t)^{1/t} \leq 1.$$

The Perron-Frobenius operator $\mathcal{P}_f^t : L^1(X_f) \rightarrow L^1(X_f)$ for $t \geq 0$ is defined by

$$\mathcal{P}_f^t(u)(z) = \sum_{w \in (T_f^t)^{-1}(z)} \frac{u(w)}{\det(DT_f^t)_w},$$

so that we have $\mathbf{Cor}_t(\psi, \varphi) = \int \mathcal{P}_f^t \psi \cdot \varphi \, dm_f$ provided that $\int \psi dm_f = 0$.

Let $C^1(X_f)$ be the set of functions φ on X_f such that $\mathcal{P}_f^t(\varphi)$ is C^1 on the interior of X_f for any $t \geq 0$ ⁴⁾. This contains all the functions that are supported and C^1 on the interior of X_f .

Now the main results are stated as follows:

Theorem 1.1. *There exists a Hilbert space $W_*(X_f)$ such that*

$$C^1(X_f) \subset W_*(X_f) \subset L^2(X_f)$$

and that the Perron-Frobenius operator \mathcal{P}_f^t for large $t \geq 0$ restricts to the bounded operator $\mathcal{P}_f^t : W_*(X_f) \rightarrow W_*(X_f)$ whose essential spectral radius is bounded by $\mathbf{m}(f)^{t/2}$.

Theorem 1.2. *For each $\rho > 1$, there exists an open and dense subset \mathcal{R} in $C_+^r(S^1)$ such that, for $f \in \mathcal{R}$, the corresponding semi-flow $\mathbf{T}_f = \{T_f^t\}$ is weakly mixing and satisfies $\mathbf{m}(f) \leq \rho \cdot \lambda_{\min}^{-1}(\mathbf{T}_f)$.*

From these theorems, we obtain the following corollary.

Corollary 1.3. *For a C^r generic $f \in C_+^r(S^1)$, the semi-flow \mathbf{T}_f is weakly mixing and the essential spectral radius of the Perron-Frobenius operator \mathcal{P}_f^t acting on $W_*(X_f)$ is bounded by $\lambda_{\min}(\mathbf{T}_f)^{-t/2}$ for any sufficiently large t .*

The estimate (1) for C^r generic f is an immediate consequence of this corollary.

Proof of (1). Take large $t > 0$. By Corollary 1.3, we have the decomposition $W_*(X_f) = E \oplus V$ where E is the sum of the generalized eigenspaces for \mathcal{P}_f^t corresponding to the eigenvalues not smaller than μ^t in absolute value and

$$V = \{\varphi \in W_*(X_f) \mid \mu^{-s} \|\mathcal{P}_f^s \varphi\|_* \rightarrow 0 \text{ as } s \rightarrow \infty\}$$

where $\|\cdot\|_*$ denotes the norm on $W_*(X_f)$. The finite dimensional subspace E is invariant under \mathcal{P}_f^s for $s \geq 0$ by commutativity of \mathcal{P}_f^s and \mathcal{P}_f^t . Thus we have $\mathcal{P}_f^s|_E = \exp(sB)$ for a linear map $B : E \rightarrow E$. Now it is easy to get the formula (1), expressing ψ as the sum of an element of V and generalized eigenvectors of B . \square

⁴⁾This condition imposes a restriction on the behavior of the function in the neighborhood of the boundary of X_f , in addition to that it should be C^1 on the interior of X_f .

Artur Avila told the author the following interesting observation related to the results above. Here we quote it under his permission. (But the author is responsible for the statement and proof.) We say that $f \in C_+^r(S^1)$ is cohomologous to a constant function if $f(x) = \varphi(\tau(x)) - \varphi(x) + c$ for some $\varphi \in C^r(S^1)$ and $c \in \mathbb{R}$.

Theorem 1.4. *The following conditions for $f \in C_+^r(S^1)$ are all equivalent:*

- (i) $\mathbf{m}(f) < 1$,
- (ii) \mathbf{T}_f is weakly mixing, and
- (iii) f is not cohomologous to a constant function.

In particular, either (or all) of these conditions holds for a C^r open and dense set of ceiling functions $f \in C_+^r(S^1)$.

This theorem and Theorem 1.1 (or Dolgopyat's argument[17]) imply that, once \mathbf{T}_f is weakly mixing, it is exponentially mixing and there are no intermediate rates in correlation decay (for functions in $C^1(X_f)$ at least). We will give the proof in Appendix A.

Remark 1.5. We can generalize the argument in this paper to a more general class of expanding semi-flows without much difficulty. For instance, the main results remains true with obvious changes in the related definitions when we consider arbitrary C^3 expanding maps on the circle in the place of τ . The proof of Theorem 1.1 can be translated almost literally to such cases using the standard estimates on the distortion of expanding maps, while we need a slight modification to translate the proof of Theorem 1.2.

Acknowledgement: The author would like to thank Artur Avila, Sébastien Gouëzel, Michihiro Hirayama and the anonymous referee for valuable comments that were crucial in improving this paper.

2. PROOF OF THEOREM 1.1

In this section we consider the semi-flow $\mathbf{T}_f = \{T_f^t\}_{t \geq 0}$ for some fixed $f \in C_+^r(S^1)$. For simplicity, we write T^t and \mathcal{P}^t for T_f^t and \mathcal{P}_f^t respectively.

2.1. Local charts on X_f . We set up a system of local charts ⁵⁾ on X_f . To begin with, we consider two small real numbers $\eta > 0$ and $\delta > 0$, and set

$$R = (-\eta, \eta) \times (\delta, 2\delta) \subset Q = (-2\eta, 2\eta) \times (0, 3\delta).$$

For each $a = (x_0, s_0) \in X_f$ such that $[x_0 - 2\eta, x_0 + 2\eta] \times \{s_0\} \subset X_f$, we consider two mappings

$$\kappa_a : Q \rightarrow X_f, \quad \kappa_a(x, s) = T^s(x_0 + x, s_0)$$

and

$$\tilde{\kappa}_a : Q \rightarrow S^1 \times \mathbb{R}, \quad \tilde{\kappa}_a(x, s) = (x_0 + x, s_0 + s).$$

Note that κ_a and $\tilde{\kappa}_a$ coincide when the image of $\tilde{\kappa}_a$ does not meet the upper boundary of X_f . Let η and δ be so small that κ_a is injective on Q whenever it is defined.

Next we take a finite subset A of X_f so that the mapping κ_a for $a \in A$ are defined and that the images $\tilde{\kappa}_a(R)$, $a \in A$, cover the subset

$$\tilde{X}_f := \{(x, s) \in S^1 \times \mathbb{R} \mid \delta/3 \leq s \leq f(x) + 2\delta/3\}.$$

⁵⁾The system of local charts does *not* give the structure of (branched) manifold on X_f .

We may and do suppose that the intersection multiplicity of $\{\tilde{\kappa}_a(R)\}_{a \in A}$ is bounded by an absolute constant (say 100). (This is in fact possible if we let the ratio η/δ be small.) Clearly the images of $\kappa_a(R)$ for $a \in A$ cover X_f .

Let $C^r(R)$ be the set of C^r functions supported on R . We take the family $\{h_a\}_{a \in A}$ of functions in $C^\infty(R)$ as follows. First take a C^∞ function $\beta_0 : \mathbb{R} \rightarrow [0, 1]$ such that $\beta_0(s) = 1$ if $s \leq \delta/3$ and $\beta_0(s) = 0$ if $s \geq 2\delta/3$. We define $\beta : S^1 \times \mathbb{R} \rightarrow [0, 1]$ by

$$\beta(x, s) = \begin{cases} \beta_0(s - f(x)), & \text{if } s \geq f(x); \\ 1, & \text{if } \delta < s < f(x); \\ 1 - \beta_0(s), & \text{if } s \leq \delta. \end{cases}$$

This is a C^∞ function supported on \tilde{X}_f . From the choice of the finite subset A , we can take a family $\{\tilde{h}_a : S^1 \times \mathbb{R} \rightarrow [0, 1]\}_{a \in A}$ of C^∞ functions so that the support of each \tilde{h}_a is contained in $\tilde{\kappa}_a(R)$ and that we have $\sum_a \tilde{h}_a \equiv \beta$ on $S^1 \times \mathbb{R}$. We then define the C^∞ function $h_a : \mathbb{R}^2 \rightarrow [0, 1]$ for $a \in A$ by

$$h_a = \begin{cases} \tilde{h}_a \circ \tilde{\kappa}_a & \text{on } R; \\ 0 & \text{on } \mathbb{R}^2 \setminus R. \end{cases}$$

2.2. Anisotropic Sobolev spaces. We recall the anisotropic Sobolev space and the and related definitions introduced in [3]. For a cone $\mathbf{C} \subset \mathbb{R}^2$, we define its dual by

$$\mathbf{C}^* = \{v \in \mathbb{R}^2 \mid (v, u) = 0 \text{ for some } u \in \mathbf{C} \setminus \{0\}\}.$$

For two cones $\mathbf{C}, \mathbf{C}' \subset \mathbb{R}^2$, we write $\mathbf{C} \Subset \mathbf{C}'$ if the closure of \mathbf{C} is contained in the interior of \mathbf{C}' *except for the origin*.

A *polarization* Θ is a combination $\Theta = (\mathbf{C}_+, \mathbf{C}_-, \varphi_+, \varphi_-)$ of closed cones \mathbf{C}_\pm in \mathbb{R}^2 and C^∞ functions $\varphi_\pm : S^1 \rightarrow [0, 1]$ on the unit circle $S^1 \subset \mathbb{R}^2$ that satisfy $\mathbf{C}_+ \cap \mathbf{C}_- = \{0\}$ and

$$(3) \quad \varphi_+(\xi) = \begin{cases} 1, & \text{if } \xi \in S^1 \cap \mathbf{C}_+; \\ 0, & \text{if } \xi \in S^1 \cap \mathbf{C}_-, \end{cases} \quad \varphi_-(\xi) = 1 - \varphi_+(\xi).$$

For two polarizations $\Theta = (\mathbf{C}_+, \mathbf{C}_-, \varphi_+, \varphi_-)$ and $\Theta' = (\mathbf{C}'_+, \mathbf{C}'_-, \varphi'_+, \varphi'_-)$, we write $\Theta < \Theta'$ if $\mathbb{R}^2 \setminus \mathbf{C}'_+ \Subset \mathbf{C}_-$.

Fix a C^∞ function $\chi : \mathbb{R} \rightarrow [0, 1]$ satisfying

$$\chi(s) = \begin{cases} 1 & \text{for } s \leq 1; \\ 0 & \text{for } s \geq 2. \end{cases}$$

For a polarization $\Theta = (\mathbf{C}_+, \mathbf{C}_-, \varphi_+, \varphi_-)$, an integer $n \geq 0$ and $\sigma \in \{+, -\}$, we define the C^∞ function $\psi_{\Theta, n, \sigma} : \mathbb{R}^2 \rightarrow [0, 1]$ by

$$\psi_{\Theta, n, \sigma}(\xi) = \begin{cases} \varphi_\sigma(\xi/|\xi|) \cdot (\chi(2^{-n}|\xi|) - \chi(2^{-n+1}|\xi|)), & \text{if } n \geq 1; \\ \chi(|\xi|)/2, & \text{if } n = 0. \end{cases}$$

This family of functions, $\psi_{\Theta, n, \sigma}$ for $n \geq 0$ and $\sigma \in \{+, -\}$, is a C^∞ partition of unity on \mathbb{R}^2 .

For a function $u \in C^r(R)$, we define

$$u_{\Theta, n, \sigma}(x) = \psi_{\Theta, n, \sigma}(D)u(x) := (2\pi)^{-2} \int e^{i(x-y)\xi} \psi_{\Theta, n, \sigma}(\xi) u(y) dy d\xi$$

where $\psi_{\Theta,n,\sigma}(D)$ is the pseudo-differential operator with symbol $a(x, \xi) = \psi_{\Theta,n,\sigma}(\xi)$. Note that the pseudo-differential operator $\psi_{\Theta,n,\sigma}(D)$ may be viewed as the composition $\mathcal{F}^{-1} \circ \Psi_{\Theta,n,\sigma} \circ \mathcal{F}$ where \mathcal{F} is the Fourier transform and $\Psi_{\Theta,n,\sigma}$ is the multiplication operator by $\psi_{\Theta,n,\sigma}$.

For a polarization $\Theta = (\mathbf{C}_+, \mathbf{C}_-, \varphi_+, \varphi_-)$ and a real number p , we define the semi-norms $\|\cdot\|_{\Theta,p}^+$ and $\|\cdot\|_{\Theta,p}^-$ on $C^r(R)$ by

$$\|u\|_{\Theta,p}^\sigma = \left(\sum_{n \geq 0} 2^{2pn} \|u_{\Theta,n,\sigma}\|_{L^2}^2 \right)^{1/2} = \left(\sum_{n \geq 0} 2^{2pn} \|\psi_{\Theta,n,\sigma} \cdot \mathcal{F}u\|_{L^2}^2 \right)^{1/2}.$$

Then we define the anisotropic Sobolev norm $\|\cdot\|_{\Theta,p,q}$ for real numbers p and q by

$$\|u\|_{\Theta,p,q} = \left((\|u\|_{\Theta,p}^+)^2 + (\|u\|_{\Theta,q}^-)^2 \right)^{1/2}.$$

Clearly this norm is associated to a scalar product.

Actually we will not use the anisotropic Sobolev norms for general p , q and Θ but those for the following specific cases. Fix small $0 < \epsilon < 1/2$ and put

$$\|\cdot\|_{\Theta}^+ := \|\cdot\|_{\Theta,1}^+, \quad \|\cdot\|_{\Theta}^- := \|\cdot\|_{\Theta,0}^-, \quad \|\cdot\|_{\Theta} := \|\cdot\|_{\Theta,1,0}$$

and

$$|\cdot|_{\Theta}^+ := \|\cdot\|_{\Theta,1-\epsilon}^+, \quad |\cdot|_{\Theta}^- := \|\cdot\|_{\Theta,-\epsilon}^-, \quad |\cdot|_{\Theta} := \|\cdot\|_{\Theta,1-\epsilon,-\epsilon}.$$

In view of Parseval's identity, we have

$$(4) \quad \|\cdot\|_{\Theta} \geq \|\cdot\|_{L^2} / \sqrt{6}$$

where 6 is (a bound for) the intersection multiplicity of the supports of $\psi_{\Theta,n,\sigma}$. The anisotropic Sobolev spaces $W_*(R; \Theta)$ and $W_{\dagger}(R; \Theta)$ are the completion of $C^\infty(R)$ with respect to the norms $\|\cdot\|_{\Theta}$ and $|\cdot|_{\Theta}$ respectively. By (4), we see that the space $W_*(R; \Theta)$ is naturally embedded in $L^2(R)$. Further we recall the next lemma from [3]: Let $W^s(R)$ be the (usual) Sobolev space of order s , that is,

$$W^s(R) = \{u \in \mathcal{D}'(\mathbb{R}^2) \mid \text{supp}(u) \subset \text{closure}(R), (1 + |\xi|^2)^{s/2} \mathcal{F}u(\xi) \in L^2(\mathbb{R}^2)\}.$$

Lemma 2.1. *For any polarizations $\Theta' < \Theta$, we have*

- (a) $C^1(R) \subset W^1(R) \subset W_*(R; \Theta) \subset L^2(R)$ and $W^{1-\epsilon}(R) \subset W_{\dagger}(R; \Theta) \subset W^{-\epsilon}(R)$.
- (b) $W_*(R; \Theta) \subset W_*(R; \Theta')$ and $W_{\dagger}(R; \Theta) \subset W_{\dagger}(R; \Theta')$.
- (c) *The inclusion $W_*(R; \Theta) \subset W_{\dagger}(R; \Theta)$ is compact.*

Proof. We can check (a) and (b) easily by using Parseval's identity. For the claim (c), we refer [3, Proposition 5.1]. \square

As for the polarization Θ , we will consider three polarizations

$$\check{\Theta}_0 = (\check{\mathbf{C}}_{0,\pm}, \check{\varphi}_{0,\pm}) < \Theta_0 = (\mathbf{C}_{0,\pm}, \varphi_{0,\pm}) < \hat{\Theta}_0 = (\hat{\mathbf{C}}_{0,\pm}, \hat{\varphi}_{0,\pm})$$

such that

$$(\mathbf{C}(\gamma_0 \theta_f))^* \Subset \hat{\mathbf{C}}_{0,-} \Subset (\mathbb{R}^2 \setminus \check{\mathbf{C}}_{0,+}) \Subset (\mathbf{C}(\theta_f))^*.$$

The Hilbert space $W_*(X_f)$ in Theorem 1.1 is defined as follows. Consider the projection operator (regarding functions as densities)

$$\Pi : (L^2(R))^A \rightarrow L^2(X_f), \quad \Pi((u_a)_{a \in A}) = \sum_{a \in A} \pi_a(u_a)$$

where $\pi_a : L^2(R) \rightarrow L^2(X_f)$ for $a \in A$ is defined by

$$\pi_a(u)(z) = \begin{cases} u(w)/\det D\kappa_a(w), & \text{if } z = \kappa_a(w) \text{ for some } w \in R; \\ 0, & \text{otherwise.} \end{cases}$$

We equip the product spaces $(W_*(R; \Theta_0))^A \subset (L^2(R))^A$ and $(W_+(R; \Theta_0))^A$ with the norms

$$\|\mathbf{u}\| := \left(\sum_{a \in A} \|u_a\|_{\Theta_0}^2 \right)^{1/2} \quad \text{and} \quad |\mathbf{u}| := \left(\sum_{a \in A} |u_a|_{\Theta_0}^2 \right)^{1/2} \quad \text{where } \mathbf{u} = (u_a)_{a \in A}$$

respectively, so that they are Hilbert spaces. Then we put

$$W_*(X_f) = \Pi((W_*(R; \Theta_0))^A)$$

and equip it with the norm $\|u\| = \inf\{\|\mathbf{u}\| \mid \Pi(\mathbf{u}) = u\}$. This space $W_*(X_f)$ is isomorphic to the orthogonal complement of the kernel of Π in $(W_*(R; \Theta_0))^A$ and, hence, is a Hilbert space.

2.3. Transfer operators on local charts. We consider the time- t -map T^t and the corresponding Perron-Frobenius operator \mathcal{P}^t for t large enough⁶). One technical difficulty in treating the time- t -map T^t directly is that it is *not* continuous at the boundary of X_f . To avoid this problem, we consider a system of transfer operators on the local charts $\{\kappa_a\}_{a \in A}$ as a lift of \mathcal{P}^t , in which we do not find any trace of the discontinuity of T^t .

For $a, b \in A$, let $Q(a, b, t)$ be the set of points $z \in Q$ such that, for some $(x, s) \in Q$,

- (i) $T^t \circ \kappa_a(z) = \kappa_b((x, s))$ and
- (ii) $T^{t-\eta} \circ \kappa_a(z) \in \kappa_b(Q)$ for $0 \leq \eta < s$.

Notice that the second condition (ii) does not follow from (i) when $\tilde{\kappa}_b(Q)$ meets the upper boundary of X_f . In fact, the second condition (ii) is important to avoid discontinuity in the following argument.

For $a, b \in A$ with $Q(a, b, t) \neq \emptyset$, we consider the C^r mapping

$$T_{ab}^t : Q(a, b, t) \rightarrow Q, \quad T_{ab}^t(z) = \kappa_b^{-1} \circ T^t \circ \kappa_a(z).$$

This is nothing but the mapping T^t viewed in the local charts κ_a and κ_b . Notice however that we restrict the domain of definition to $Q(a, b, t)$.

For $a, b \in A$ with $Q(a, b, t) \neq \emptyset$, we consider the transfer operator

$$\mathcal{P}_{ab} : L^2(R) \rightarrow L^2(R), \quad \mathcal{P}_{ab}^t u(z) = \sum_{w \in (T_{ab}^t)^{-1}(z)} \frac{h_b(z)u(w)}{\det(DT_{ab}^t)_w}$$

where the sum is taken over $w \in Q(a, b, t)$ such that $T_{ab}^t(w) = z$. And we define the system of transfer operators on local charts,

$$\mathbf{P}^t : L^2(R)^A \rightarrow L^2(R)^A,$$

by

$$\mathbf{P}^t(\mathbf{u}) = \left(\sum_{a \in A} \mathcal{P}_{ab}^t(u_a) \right)_{b \in A} \quad \text{for } \mathbf{u} = (u_a)_{a \in A} \in L^2(R)^A.$$

⁶)It is enough to consider t with $t \geq \max_{x \in S^1} f(x) + 3\delta$

It is then not difficult to check that the following diagram commutes:

$$(5) \quad \begin{array}{ccc} L^1(R)^A & \xrightarrow{\mathbf{P}^t} & L^1(R)^A \\ \downarrow \Pi & & \downarrow \Pi \\ L^1(X_f) & \xrightarrow{\mathcal{P}^t} & L^1(X_f) \end{array}$$

In the following subsections, we will prove

Proposition 2.2. *The operator \mathbf{P}^t restricts to the bounded operator*

$$\mathbf{P}^t : W_*(R; \Theta_0)^A \rightarrow W_*(R; \Theta_0)^A.$$

Also \mathbf{P}^t extends to the bounded operator

$$\mathbf{P}^t : W_{\dagger}(R; \Theta_0)^A \rightarrow W_{\dagger}(R; \Theta_0)^A.$$

Further we have the Lasota-Yorke type inequality

$$(6) \quad \|\mathbf{P}^t(\mathbf{u})\| \leq C_{\sharp} \cdot \mathbf{m}(f, t)^{1/2} \|\mathbf{u}\| + C|\mathbf{u}| \quad \text{for } \mathbf{u} \in W_*(R)^A$$

where the constant C_{\sharp} does not depend on t while the constant C may.

We can deduce Theorem 1.1 from this proposition and Lemma 2.1(c).

Proof of Theorem 1.1. Since $W_*(R; \Theta_0)^A$ is compactly embedded in $W_{\dagger}(R; \Theta_0)^A$ from Lemma 2.1(c), the Lasota-Yorke type inequality in Proposition 2.2 implies that the essential spectral radius of the operator $\mathbf{P}^t : W_*(R; \Theta_0)^A \rightarrow W_*(R; \Theta_0)^A$ is bounded by $C_{\sharp} \cdot \mathbf{m}(f, t)^{1/2}$. (See [13, 14].) By the definition of the space $W_*(X_f)$, the commutative diagram (5) restricts to

$$(7) \quad \begin{array}{ccc} W_*(R; \Theta_0)^A & \xrightarrow{\mathbf{P}^t} & W_*(R; \Theta_0)^A \\ \downarrow \Pi & & \downarrow \Pi \\ W_*(X_f) & \xrightarrow{\mathcal{P}^t} & W_*(X_f). \end{array}$$

Recall also that the space $W_*(X_f)$ is identified with the orthogonal complement of the kernel of Π in $W_*(R)^A$. Through this identification, \mathcal{P}^t corresponds to the composition of \mathbf{P}^t with the orthogonal projection along the kernel of Π . Thus the essential spectral radius of $\mathcal{P}^t : W_*(X_f) \rightarrow W_*(X_f)$ is bounded by that of $\mathbf{P}^t : W_*(R; \Theta_0)^A \rightarrow W_*(R; \Theta_0)^A$ or $C_{\sharp} \mathbf{m}(f, t)^{1/2}$. Since this holds for any t large enough, the essential spectral radius of $\mathcal{P}^t : W_*(X_f) \rightarrow W_*(X_f)$ is bounded by $\mathbf{m}(f)^{1/2}$. The inclusions $C^1(X_f) \subset W_*(X_f) \subset L^2(X_f)$ follows from Lemma 2.1(a). \square

2.4. Two lemmas on the anisotropic Sobolev norms. In this section, we give two basic lemmas on the anisotropic Sobolev norms. These lemmas and their proofs are slight modification of those given in [3]. For convenience of the reader, we give the proofs in Appendix B and C. (See also Remark 2.5.)

Lemma 2.3. *Let $g_i : \mathbb{R}^2 \rightarrow [0, 1]$, $1 \leq i \leq I$, be a family of C^r functions such that $\sum_{i=1}^I g_i(x) \leq 1$ for $x \in Q$ and that $\text{supp}(g_i) \subset Q$. Let Θ and Θ' be polarizations such that $\Theta' < \Theta$. Then we have*

$$(8) \quad \left[\sum_{1 \leq i \leq I} \|g_i u\|_{\Theta'}^2 \right]^{1/2} \leq C_0 \|u\|_{\Theta} + C|u|_{\Theta} \quad \text{for } u \in W_*(R; \Theta),$$

where C_0 is a constant that does not depend on $\{g_i\}$ while the constant C may. Further, if $\sum_{i=1}^I g_i(x) \equiv 1$ for all $x \in R$ in addition, we also have

$$(9) \quad \|u\|_{\Theta'} \leq \nu \left[\sum_{1 \leq i \leq I} \|g_i u\|_{\Theta}^2 \right]^{1/2} + C \sum_{i=1}^I |g_i u|_{\Theta} \quad \text{for } u \in W_*(R; \Theta),$$

where ν is the intersection multiplicity of the supports of the functions g_i , $1 \leq i \leq I$.

In the next lemma, we consider the following situation. For a C^{r-1} -function $h : \mathbb{R}^2 \rightarrow \mathbb{R}$ supported on a closed subset $K \subset R$ and for a C^r -diffeomorphism $S : U \rightarrow S(U) \subset \mathbb{R}^2$ defined on an open neighborhood U of K , we consider a transfer operator $L : C^{r-1}(R) \rightarrow C^{r-1}(R)$ defined by

$$Lu(x) = \begin{cases} h(x) \cdot u \circ S(x), & \text{if } x \in K; \\ 0 & \text{otherwise.} \end{cases}$$

Assume that, for polarizations $\Theta = (\mathbf{C}_+, \mathbf{C}_-, \varphi_+, \varphi_-)$ and $\Theta' = (\mathbf{C}'_+, \mathbf{C}'_-, \varphi'_+, \varphi'_-)$, we have

$$(DS_{\zeta})^{tr}(\mathbb{R}^2 \setminus \mathbf{C}_+) \Subset \mathbf{C}'_- \quad \text{for all } \zeta \in K,$$

where $(DS_{\zeta})^{tr}$ denotes the transpose of DS_{ζ} . Put

$$\gamma(S) = \min_{\zeta \in K} |\det DS_{\zeta}|$$

and

$$\Lambda(S, \Theta', K) = \sup \left\{ \frac{\|(DS_{\zeta})^{tr}(v)\|}{\|v\|} \mid \zeta \in K, (DS_{\zeta})^{tr}(v) \notin \mathbf{C}'_- \right\}.$$

Lemma 2.4. *The operator L extends boundedly to $L : W_*(R; \Theta) \rightarrow W_*(R; \Theta')$ and to $L : W_{\dagger}(R; \Theta) \rightarrow W_{\dagger}(R; \Theta')$. Further we have, for $u \in W_*(R; \Theta)$,*

$$(10) \quad \|Lu\|_{\Theta'} \leq \gamma(S)^{-1/2} \|h\|_{L^\infty} \|u\|_{\Theta}$$

and

$$(11) \quad \|Lu\|_{\Theta'}^{\pm} \leq C_0 \gamma(S)^{-1/2} \Lambda(S, \Theta', K) \|h\|_{L^\infty} \|u\|_{\Theta} + C|u|_{\Theta}$$

where the constant C_0 does not depend on S , h , Θ nor Θ' while the constant C may. In particular, we have, for $u \in W_*(R; \Theta)$,

$$(12) \quad \|Lu\|_{\Theta'} \leq C_0 \gamma(S)^{-1/2} \max\{1, \Lambda(S, \Theta', K)\} \|h\|_{L^\infty} \|u\|_{\Theta} + C|u|_{\Theta}.$$

Remark 2.5. The latter claim (9) of Lemma 2.3 is a special case of [3, Lemma 7.1]. Also, Lemma 2.4 and the former claim (8) in Lemma 2.3 correspond to [3, Proposition 7.2 and 6.1]. However, since we considered only anisotropic Sobolev norms $\|\cdot\|_{\Theta, p, q}$ with $q < 0 < p$ in those propositions in [3], we need to modify the statements and proofs slightly. This is the reason why we give the proofs of Lemma 2.4 and the former claim (8) of Lemma 2.3 in the appendices B and C.

2.5. The Lasota-Yorke type inequality in local charts. To complete the proof of Proposition 2.2, we have only to show the Lasota-Yorke type inequality (6), because the other claims now follow from Lemma 2.4 with S the branches of the inverse of T_{ab}^t . For the proof of the Lasota-Yorke type inequality (6), it is enough to show the following lemma for the components \mathcal{P}_{ab}^t of \mathbf{P}^t .

Lemma 2.6. *For $a, b \in A$ and for sufficiently large t , we have*

$$\|\mathcal{P}_{ab}^t(u)\|_{\Theta_0}^2 \leq C_{\sharp} \cdot \mathbf{m}(f, t) \|u\|_{\Theta_0}^2 + C |u|_{\Theta_0}^2 \quad \text{for } u \in C^r(R)$$

where the constant C_{\sharp} does not depend on t while the constant C may.

Below we prove Lemma 2.6. To begin with, we set up some notation. Fix $a, b \in A$ and consider large $t > 0$. Let $\{D(\omega), \omega \in \Omega\}$, be a finite family of small closed disks whose interiors cover the closure of R . We may and do assume that the intersection multiplicity of this cover is bounded by some absolute constant (say 4). Note that we can take such family of disks with arbitrarily small diameters. Let $D(\omega, i)$, $1 \leq i \leq I(\omega)$, be the connected components of the preimage $(T_{ab}^t)^{-1}(D(\omega))$ that meet the closure of R . Then $\det DT_{ab}^t$ takes constant value on each component $D(\omega, i)$, which is denoted by $e(\omega, i)$. From (2), it holds

$$(13) \quad \sum_{1 \leq i \leq I(\omega)} e(\omega, i)^{-1} \leq 1 \quad \text{for any } \omega \in \Omega.$$

We write $i \pitchfork_{\omega} j$ for $1 \leq i, j \leq I(\omega)$ if

$$(14) \quad (DT_{ab}^t)_z(\mathbf{C}_f) \cap (DT_{ab}^t)_w(\mathbf{C}_f) = \{0\}$$

for any $z \in D(\omega, i)$ and $w \in D(\omega, j)$. Note that (14) holds if and only if

$$(((DT_{ab}^t)_z)^{tr})^{-1}((\mathbf{C}_f)^*) \cap (((DT_{ab}^t)_w)^{tr})^{-1}((\mathbf{C}_f)^*) = \{0\}.$$

Letting the diameters of the disks $D(\omega)$ be small, we may assume

$$\sum_{j \not\pitchfork_{\omega} i} e(\omega, j)^{-1} \leq \mathbf{m}(f, t) \quad \text{for } 1 \leq i \leq I(\omega) \text{ and } \omega \in \Omega$$

where $\sum_{j \not\pitchfork_{\omega} i}$ denotes the sum over $1 \leq j \leq I(\omega)$ such that $j \not\pitchfork_{\omega} i$. Further, we can take polarizations $\Theta(\omega, i) = (\mathbf{C}_{\omega, i, +}, \mathbf{C}_{\omega, i, -}, \varphi_{\omega, i, +}, \varphi_{\omega, i, -})$ for each $\omega \in \Omega$ and $1 \leq i \leq I(\omega)$ such that

$$(((DT_{ab}^t)_z)^{tr})^{-1}(\mathbb{R}^2 \setminus \hat{\mathbf{C}}_{0,+}) \Subset \mathbf{C}_{\omega, i, -} \Subset (\mathbb{R}^2 \setminus \mathbf{C}_{\omega, i, +}) \Subset \hat{\mathbf{C}}_{0,-} \quad \text{for any } z \in D(\omega, i)$$

and that

$$\overline{(\mathbb{R}^2 \setminus \mathbf{C}_{\omega, i, +})} \cap \overline{(\mathbb{R}^2 \setminus \mathbf{C}_{\omega, j, +})} = \{0\} \quad \text{if } i \pitchfork_{\omega} j.$$

Take a family of C^∞ functions $g_\omega : \mathbb{R}^2 \rightarrow [0, 1]$ for $\omega \in \Omega$ such that the support of each g_ω is contained in $D(\omega)$ and that $\sum_{\omega \in \Omega} g_\omega(z) \equiv 1$ for all $z \in R$. We then define the functions $g_{\omega, i} : \mathbb{R}^2 \rightarrow [0, 1]$ for $\omega \in \Omega$ and $1 \leq i \leq I(\omega)$ by

$$g_{\omega, i}(z) = \begin{cases} g_\omega(T_{ab}^t(z)), & \text{if } z \in D(\omega, i) \text{ and} \\ 0, & \text{otherwise.} \end{cases}$$

Now we start the proof of Lemma 2.6. In the following, we will write C_{\sharp} for constants that do not depend on t while write C for constants that may depend on t . (Notice that the value of the constants denoted by C_{\sharp} and C are different from place to place.) We view the operator \mathcal{P}_{ab}^t under consideration as the composition of the four operations

- (i) breaking a function $u \in C^r(R)$ into $u_{\omega, i} := g_{\omega, i}u$, $\omega \in \Omega$, $1 \leq i \leq I(\omega)$,
- (ii) transforming each $u_{\omega, i}$ to $v_{\omega, i} := \mathcal{P}_{ab}^t(u_{\omega, i})$,
- (iii) summing up $v_{\omega, i}$ for $1 \leq i \leq I(\omega)$ to get $v_\omega := \sum_{1 \leq i \leq I(\omega)} v_{\omega, i} = g_\omega \mathcal{P}_{ab}^t u$,
- (iv) summing up v_ω for $\omega \in \Omega$ to get $\mathcal{P}_{ab}^t u = \sum_{\omega} v_\omega$.

For the operation (i), the former claim (8) of Lemma 2.3 gives the estimate

$$\sum_{\omega \in \Omega} \sum_{i=1}^{I(\omega)} \|u_{\omega,i}\|_{\check{\Theta}_0}^2 \leq (C_{\sharp} \|u\|_{\Theta_0} + C|u|_{\Theta_0})^2 \leq C_{\sharp} \|u\|_{\check{\Theta}_0}^2 + C|u|_{\check{\Theta}_0}^2.$$

For the operation (iv), the latter claim (9) of Lemma 2.3 gives the estimate

$$\|\mathcal{P}_{ab}^t u\|_{\check{\Theta}_0}^2 = \left\| \sum_{\omega \in \Omega} g_{\omega} \mathcal{P}_{ab}^t u \right\|_{\check{\Theta}_0}^2 = \left\| \sum_{\omega \in \Omega} v_{\omega} \right\|_{\check{\Theta}_0}^2 \leq C_{\sharp} \sum_{\omega \in \Omega} \|v_{\omega}\|_{\check{\Theta}_0}^2 + C \sum_{\omega \in \Omega} |v_{\omega}|_{\check{\Theta}_0}^2.$$

Below we consider the operations (ii) and (iii). Letting S be a branch of $(T_{ab}^t)^{-1}$ and $K = \text{supp}(g_{\omega})$ in Lemma 2.4, we obtain the estimates

$$(15) \quad |v_{\omega,i}|_{\Theta(\omega,i)} \leq C|u_{\omega,i}|_{\check{\Theta}_0} \quad \text{and}$$

$$(16) \quad \|v_{\omega,i}\|_{\check{\Theta}_0} \leq C_{\sharp} \cdot e(\omega,i)^{-1/2} \|u_{\omega,i}\|_{\check{\Theta}_0} + C|u_{\omega,i}|_{\check{\Theta}_0},$$

$$(17) \quad \|v_{\omega,i}\|_{\check{\Theta}_0}^{\dagger} \leq C_{\sharp} \cdot e(\omega,i)^{-3/2} \|u_{\omega,i}\|_{\check{\Theta}_0} + C|u_{\omega,i}|_{\check{\Theta}_0}.$$

The next lemma is the core of our argument, in which we make use of the transversality condition (14) essentially.

Lemma 2.7. *If $i \pitchfork_{\omega} j$, we have*

$$\sum_{n \geq 0} \left| \left(\psi_{\check{\Theta}_0, n, -}(D)v_{\omega,i}, \psi_{\check{\Theta}_0, n, -}(D)v_{\omega,j} \right)_{L^2} \right| \leq C|v_{\omega,i}|_{\Theta(\omega,i)} |v_{\omega,j}|_{\Theta(\omega,j)}.$$

Proof. Put $w_{i,n} = \psi_{\check{\Theta}_0, n, -}(D)v_{\omega,i}$, $w'_{i,n} = \psi_{\Theta(\omega,i), n, -}(D)v_{\omega,i}$, $w''_{i,n} = w_{i,n} - w'_{i,n}$. For $n > 0$, we have, from the assumption $i \pitchfork_{\omega} j$, that $(w'_{i,n}, w'_{j,n})_{L^2} = 0$, that is,

$$(w_{i,n}, w_{j,n})_{L^2} = (w''_{i,n}, w'_{j,n})_{L^2} + (w'_{i,n}, w''_{j,n})_{L^2} + (w''_{i,n}, w''_{j,n})_{L^2}.$$

Since we have that $\|w'_{i,n}\|_{L^2} \leq 2^{\epsilon n} |v_{\omega,i}|_{\Theta(\omega,i)}$ and that $\|w''_{i,n}\|_{L^2} \leq 2^{-(1-\epsilon)n} |v_{\omega,i}|_{\Theta(\omega,i)}$ by the definition of the norm $|\cdot|_{\Theta(\omega,i)}$ and since $0 < \epsilon < 1/2$, we can get the lemma using Schwarz inequality. \square

It follows from Lemma 2.7 that

$$\left(\|v_{\omega}\|_{\check{\Theta}_0}^- \right)^2 \leq \sum_i \sum_{j \not\pitchfork_{\omega} i} \|v_{\omega,i}\|_{\check{\Theta}_0} \|v_{\omega,j}\|_{\check{\Theta}_0} + C \sum_i |v_{\omega,i}|_{\check{\Theta}_0}^2$$

for some constant $C > 0$ that may depend on $I(\omega)$, where $\sum_{j \not\pitchfork_{\omega} i}$ denotes the sum over $1 \leq j \leq I(\omega)$ such that $j \not\pitchfork_{\omega} i$. Applying the inequality (16) and (15) in the right hand side along, we obtain

$$\begin{aligned} \left(\|v_{\omega}\|_{\check{\Theta}_0}^- \right)^2 &\leq C_{\sharp} \sum_i \sum_{j \not\pitchfork_{\omega} i} \frac{e(\omega,j)^{-1} \|u_{\omega,i}\|_{\check{\Theta}_0}^2 + e(\omega,i)^{-1} \|u_{\omega,j}\|_{\check{\Theta}_0}^2}{2} + C \sum_i |u_{\omega,i}|_{\check{\Theta}_0}^2 \\ &\leq C_{\sharp} \mathbf{m}(f, t) \sum_i \|u_{\omega,i}\|_{\check{\Theta}_0}^2 + C \sum_i |u_{\omega,i}|_{\check{\Theta}_0}^2. \end{aligned}$$

On the other hand we obtain, from (17), that

$$\begin{aligned} \left(\|v_{\omega}\|_{\check{\Theta}_0}^+ \right)^2 &\leq C_{\sharp} \left(\sum_i e(\omega,i)^{-3/2} \|u_{\omega,i}\|_{\check{\Theta}_0} \right)^2 + C \left(\sum_i |u_{\omega,i}|_{\check{\Theta}_0} \right)^2 \\ &\leq C_{\sharp} \mathbf{m}(f, t) \sum_i \|u_{\omega,i}\|_{\check{\Theta}_0}^2 + C \sum_i |u_{\omega,i}|_{\check{\Theta}_0}^2 \end{aligned}$$

where we used Schwarz inequality, (13) and the simple fact $e(\omega, i)^{-1} \leq \mathbf{m}(f, t)$ in the second inequality. Therefore we obtain, for the operations (ii) and (iii),

$$\left(\|v_\omega\|_{\Theta_0}\right)^2 \leq C_{\#} \mathbf{m}(f, t) \sum_i \|u_{\omega, i}\|_{\Theta_0}^2 + C \sum_i |u_{\omega, i}|_{\Theta_0}^2.$$

Letting S be the identity map in Lemma 2.4, we see that $|u_{\omega, i}|_{\Theta_0} \leq C|u|_{\Theta_0}$ and that $|v_{\omega, i}|_{\Theta_0} \leq C|v_{\omega, i}|_{\Theta(\omega, i)}$. These and (15) give

$$|v_\omega|_{\Theta_0} \leq C \sum_{1 \leq i \leq I(\omega)} |v_{\omega, i}|_{\Theta(\omega, i)} \leq C \sum_{1 \leq i \leq I(\omega)} |u_{\omega, i}|_{\Theta_0} \leq C|u|_{\Theta_0}.$$

We can now conclude the Lasota-Yorke type inequality in Lemma 2.6 from the estimates on the operations (i)–(iv) given above.

3. PROOF OF THEOREM 1.2

The proof of Theorem 1.2 below is a modification of the argument in [20].

3.1. Notations. We recall some notation from [20]. Let $\mathcal{A} = \{1, 2, \dots, \ell\}$ and let \mathcal{A}^n be the space of words of length n on \mathcal{A} . For a word $\mathbf{a} = (a_i)_{i=1}^n \in \mathcal{A}^n$ and an integer $0 \leq p \leq n$, let $[\mathbf{a}]_p = (a_i)_{i=1}^p$. For $0 \leq p \leq n$, we define the equivalence relation \sim_p on \mathcal{A}^n so that $\mathbf{a} \sim_p \mathbf{b}$ if and only if $[\mathbf{a}]_p = [\mathbf{b}]_p$.

Let \mathcal{P} be the partition of S^1 into the intervals $\mathcal{P}(k) = [(k-1)/\ell, k/\ell)$ for $k \in \mathcal{A}$. Then $\mathcal{P} = \bigvee_{i=0}^{n-1} \tau^{-i}(\mathcal{P})$ is the partition into the intervals

$$\mathcal{P}(\mathbf{a}) = \bigcap_{i=0}^{n-1} \tau^{-i}(\mathcal{P}(a_{n-i})), \quad \mathbf{a} = (a_i)_{i=1}^n \in \mathcal{A}^n.$$

Let $x_{\mathbf{a}}$ the left end point of the interval $\mathcal{P}(\mathbf{a})$.

Remark 3.1. Notice that \mathbf{a} is the *inverse* of the itinerary of the points in $\mathcal{P}(\mathbf{a})$.

For a point $x \in S^1$ and $\mathbf{a} = (a_i)_{i=1}^n \in \mathcal{A}^n$, we denote by $\mathbf{a}(x)$ the unique point $y \in \mathcal{P}(\mathbf{a})$ such that $\tau^n(y) = x$. For a C^r function $f \in C^r(S^1)$, $x \in S^1$ and $\mathbf{b} \in \mathcal{A}^n$, we put

$$s(x, \mathbf{b}; f) := f^{(n)}(\mathbf{b}(x)) = \sum_{i=1}^n f([\mathbf{b}]_i(x)).$$

Then we have

$$\frac{ds}{dx}(x, \mathbf{b}; f) = \sum_{i=1}^n \ell^{-i} \frac{df}{dx}([\mathbf{b}]_i(x)).$$

We will identify the unit circle S^1 with the lower boundary $S^1 \times \{0\}$ of X_f . If $f^{(n)}(\mathbf{b}(x)) \leq t < f^{(n+1)}(\mathbf{b}(x))$, the image of the horizontal tangent vector $(1, 0)$ at $\mathbf{b}(x) \in S^1 \times \{0\}$ by the mapping T_f^t has slope $\frac{d}{dx}s(x, \mathbf{b}; f)$ and hence

$$(DT_f^t)_{\mathbf{b}(x)}(\mathbf{C}_f) = \left\{ (\xi, \eta) \in \mathbb{R}^2 \mid \left| \eta - \frac{ds}{dx}(x, \mathbf{b}; f) \cdot \xi \right| \leq \ell^{-n} \theta_f |\xi| \right\}.$$

For $K > 1$, let $C_+^r(S^1; K)$ be the set $f \in C_+^r(S^1)$ such that $K^{-1} < f(x) < K$ for $x \in S^1$ and $\|f\|_{C^r} < K$. By virtue of Theorem 1.4, it is sufficient for the proof of Theorem 1.2 to show that, for each $\rho > 1$ and $K > 0$, the condition

$$(18) \quad \mathbf{m}(f) \leq \rho \cdot \lambda_{\min}(\mathbf{T}_f)^{-1}$$

holds for functions f in an open and dense subset of $C_+^r(S^1; K)$. We will prove this claim in the following. We henceforth fix arbitrary $\rho > 1$ and $K > 0$. Note that we have, for $f \in C_+^r(S^1; K)$, that

$$\theta_f \leq \theta_K := K/(\gamma_0 \ell - 1), \quad \ell^{1/K} \leq \lambda_{\min}(\mathbf{T}_f) \leq \ell^K$$

and also that

$$(19) \quad \left| \frac{d^2 s}{d^2 x}(x, \mathbf{b}; f) \right| = \left| \sum_{i=1}^n \ell^{-2i} \frac{d^2 f}{dx^2}([\mathbf{b}]_i(x)) \right| \leq \frac{\ell^{-2} K}{1 - \ell^{-2}} \leq \theta_K$$

for $x \in S^1$ and $\mathbf{b} \in \mathcal{A}^n$.

3.2. Some consequences of the condition $\mathbf{m}(f) > \rho \cdot \lambda_{\min}(\mathbf{T}_f)^{-1}$. In this subsection, we see what kind of singular situation occurs if the condition (18) does not hold. Fix $1 < \gamma < \ell$ such that $\gamma^K < \rho$.

Proposition 3.2. *If $\mathbf{m}(f) > \rho \cdot \lambda_{\min}(\mathbf{T}_f)^{-1}$ for $f \in C_+^r(S^1; K)$, then, for any $n \geq 1$, there exist $\mathbf{c} \in \mathcal{A}^n$ and $B \subset \mathcal{A}^n$ with $\#B \geq \gamma^n$ such that*

$$\left| \frac{ds}{dx}(x_{\mathbf{c}}, \mathbf{a}; f) - \frac{ds}{dx}(x_{\mathbf{c}}, \mathbf{b}; f) \right| \leq 8\theta_K \cdot \ell^{-n} \quad \text{for all } \mathbf{a}, \mathbf{b} \in B.$$

Proof. Take $\gamma < \bar{\gamma} < 1$ so close to γ that $\bar{\gamma}^K < \rho$. Then take $1 < \lambda < \lambda_{\min}(\mathbf{T}_f)$ so close to $\lambda_{\min}(\mathbf{T}_f)$ that $\bar{\gamma}^K \lambda_{\min}(\mathbf{T}_f) < \rho \lambda$. From the assumption, we can take an arbitrarily large $t \geq 0$, $z \in X_f$, $w \in T^{-t}(z)$ and $Z \subset T^{-t}(z)$ such that

$$(20) \quad (DT^t)_\zeta(\mathbf{C}_f) \cap (DT^t)_w(\mathbf{C}_f) \neq \{0\} \quad \text{for } \zeta \in Z$$

and that

$$(21) \quad \sum_{\zeta \in Z} \frac{1}{E(\zeta, t; f)} \geq \rho^t \cdot \lambda_{\min}(\mathbf{T}_f)^{-t}.$$

We may and do assume in addition that $z \in S^1 \times \{0\}$. Put

$$m = \min \{n(x, s + t; f) \mid (x, s) \in Z\}.$$

(Recall the definition of $n(x, t; f)$ in Section 1.) Then we have

$$(22) \quad K^{-1}t < m < Kt \quad \text{and} \quad \ell^m > \lambda^t$$

provided that t is sufficiently large. (To see the second inequality, note that we have $\ell^m \sim \lambda_{\min}(\mathbf{T}_f)^t$ for large t .)

For each $\zeta = (x, s) \in Z$, let $I(\zeta) \in \mathcal{A}^{n(x, s+t; f)}$ be the word such that $\zeta \in \mathcal{P}(I(\zeta))$. Put $A = \{[I(\zeta)]_m \mid \zeta \in Z\} \subset \mathcal{A}^m$. Then, applying the formula (2), we can see that

$$\#A \cdot \ell^{-m} \geq \sum_{\zeta \in Z} \frac{1}{E(\zeta, t; f)}.$$

Combining this inequality with (21), we see

$$(23) \quad \#A \geq \ell^m \rho^t \lambda_{\min}(\mathbf{T}_f)^{-t} > (\lambda \rho \cdot \lambda_{\min}(\mathbf{T}_f)^{-1})^t > \bar{\gamma}^{Kt} > \bar{\gamma}^m.$$

Note that the condition (20) implies

$$(24) \quad \left| \frac{ds}{dx}(z, \mathbf{b}; f) - \frac{ds}{dx}(z, \mathbf{b}'; f) \right| \leq 4\theta_K \cdot \ell^{-m} \quad \text{for all } \mathbf{b}, \mathbf{b}' \in A.$$

Now we start to consider an arbitrary integer $n \geq 1$ in the claim. Taking large t in the beginning, we may and do assume that the integer m above is much larger

than n . For each $0 \leq k \leq m$, we split A into equivalence classes with respect to \sim_k and let $A_k \subset A$ be one of those equivalence classes with maximum cardinality. Then the cardinality $q(k)$ of A_k is a decreasing sequence with respect to k . Further we have $q(0) \geq \bar{\gamma}^m$ by (23) and also $q(m) = 1$ obviously. Therefore we can find an integer $0 \leq m' \leq m - n$ such that $q(m' + n) \leq \gamma^{-n}q(m')$, provided that we took sufficiently large t in the beginning. We fix such integer m' and let $A' \subset \mathcal{A}^{m-m'}$ be the set of words that are obtained by removing the first common m' letters (say \mathbf{c}) from the words in $A_{m'}$, and put $x = \mathbf{c}(z)$. Then it follows from (24) that

$$\left| \frac{ds}{dx}(x, \mathbf{b}; f) - \frac{ds}{dx}(x, \mathbf{b}'; f) \right| \leq 4\theta_K \cdot \ell^{-(m-m')} \leq 4\theta_K \cdot \ell^{-n} \quad \text{for all } \mathbf{b}, \mathbf{b}' \in A'.$$

Put $B = \{[\mathbf{a}]_n \mid \mathbf{a} \in A'\}$. From the condition $q(m' + n) \leq \gamma^{-n}q(m')$ in the choice of m' , we have that $\#B \geq \gamma^n$. Also, since

$$\left| \frac{ds}{dx}(x, [\mathbf{a}]_n; f) - \frac{ds}{dx}(x, \mathbf{a}; f) \right| \leq \theta_K \cdot \ell^{-n} \quad \text{for } \mathbf{a} \in \mathcal{A}^{m-m'},$$

we see that

$$\left| \frac{ds}{dx}(x, \mathbf{b}; f) - \frac{ds}{dx}(x, \mathbf{b}'; f) \right| \leq 6\theta_K \cdot \ell^{-n} \quad \text{for all } \mathbf{b}, \mathbf{b}' \in B.$$

For $\mathbf{b}, \mathbf{c} \in \mathcal{A}^n$ and $f \in C_+^r(S^1; K)$, the variation of $\frac{d}{dx}s(\cdot, \mathbf{b}; f)$ on the interval $\mathcal{P}(\mathbf{c})$ is bounded by $\theta_K \ell^{-n}$, in view of (19). Therefore, translating the point x to the point $x_{\mathbf{c}}$, we obtain the conclusion of the proposition. \square

To state the next proposition, we set up some constants. First take real numbers α and β such that $1 < \beta < \alpha < \gamma$ and then take positive integers p and ν such that

$$\beta^{-p}\ell^2 < 1 \quad \text{and} \quad (\nu + 1)(p + 1)\alpha^{-\nu} < 1.$$

We put

$$\delta = \frac{\log \gamma - \log \alpha}{\log \ell - \log \alpha} \in (0, 1).$$

Then we choose an integer $N > \nu$ such that

$$\ell^\nu \alpha^n < \gamma^n \quad \text{for } n \geq N$$

and that

$$\ell^{-\nu}(\gamma/\beta)^{n'}(1 - (\nu + 1)(p + 1)\alpha^{-\nu}) \geq 1 \quad \text{for } n' \geq \delta N.$$

Proposition 3.3. *If $\mathbf{m}(f) > \rho \cdot \lambda_{\min}(\mathbf{T}_f)^{-1}$ for $f \in C_+^r(S^1; K)$, then, for any $n \geq N$, there exist an integer $\delta n \leq n' \leq n$, a word $\mathbf{d} \in \mathcal{A}^{n'}$ and mutually disjoint subsets $B_i \subset \mathcal{A}^{n'}$ for $1 \leq i \leq (\nu + 1)(p + 1)$ such that*

- (a) $\left| \frac{ds}{dx}(x_{\mathbf{d}}, \mathbf{b}; f) - \frac{ds}{dx}(x_{\mathbf{d}}, \mathbf{b}'; f) \right| \leq 10\theta_K \cdot \ell^{-n'}$ for all $\mathbf{b}, \mathbf{b}' \in \bigcup_{i=1}^{(\nu+1)(p+1)} B_i$,
- (b) $\#B_i \geq \beta^{n'}$ for $1 \leq i \leq (\nu + 1)(p + 1)$, and
- (c) $[\mathbf{a}]_\nu = [\mathbf{b}]_\nu$ for $\mathbf{a} \in B_i$ and $\mathbf{b} \in B_j$ if and only if $i = j$.

Proof. Let $n \geq N$ and let $B \subset \mathcal{A}^n$ be the subset in the conclusion of Proposition 3.2. For $0 \leq k \leq [n/\nu]$, we split B into equivalence classes with respect to $\sim_{k\nu}$ and let $B_k \subset B$ be one of those equivalence classes with maximum cardinality. Then the cardinality $q(k)$ of B_k is decreasing with respect to k and satisfies $q(0) = \#B \geq \gamma^n$

and $q(\lfloor n/\nu \rfloor) \leq \ell^\nu < \gamma^n \alpha^{-\lfloor n/\nu \rfloor \nu}$, where the last inequality follows from the first condition in the choice of N .

Let k_0 be the smallest integer $1 \leq k \leq \lfloor n/\nu \rfloor$ such that $q(k) < \gamma^n \alpha^{-k\nu}$. By this choice of k_0 , we have

$$(25) \quad q(k_0) < \alpha^{-\nu} q(k_0 - 1) \quad \text{and} \quad q(k_0 - 1) \geq \gamma^n \alpha^{-(k_0-1)\nu}.$$

Put $n' = n - (k_0 - 1)\nu$. Since $q(k) \leq \ell^{n-\nu k}$ obviously, we have

$$\ell^{n'} = \ell^{n-(k_0-1)\nu} \geq q(k_0 - 1) \geq \gamma^n \alpha^{n'-n} \quad \text{or} \quad n' \geq \frac{\log \gamma - \log \alpha}{\log \ell - \log \alpha} \cdot n = \delta n.$$

Let $B'_i \subset B_{k_0-1}$, $1 \leq i \leq \ell^\nu$, be the equivalence classes in B_{k_0-1} with respect to the relation \sim_{k_0} , arranged in decreasing order of cardinality⁷⁾. Then we have a simple inequality

$$\begin{aligned} \min_{1 \leq i \leq (\nu+1)(p+1)} \#B'_i &\geq \frac{q(k_0 - 1) - (\nu + 1)(p + 1)q(k_0)}{\ell^\nu} \\ &\geq \ell^{-\nu} \gamma^n \alpha^{-(k_0-1)\nu} (1 - (\nu + 1)(p + 1)\alpha^{-\nu}) \geq \beta^{n'} \end{aligned}$$

where the second inequality follows from (25) and the last from the second condition in the choice of N . Finally let $B_i \subset \mathcal{A}^{n'}$ for $1 \leq i \leq (\nu + 1)(p + 1)$ be the subset of words that are obtained by removing the first common $(k_0 - 1)\nu$ letters (say \mathbf{c}') from the words in B'_i . Then the conditions (b) and (c) hold. From the condition on the subset B in Proposition 3.2, we have

$$\left| \frac{ds}{dx}(x_{\mathbf{c}\mathbf{c}'}, \mathbf{b}; f) - \frac{ds}{dx}(x_{\mathbf{c}\mathbf{c}'}, \mathbf{b}'; f) \right| \leq 8\theta_K \cdot \ell^{-n'} \quad \text{for all } \mathbf{b}, \mathbf{b}' \in \bigcup_{i=0}^{(\nu+1)(p+1)} B_i.$$

Take $\mathbf{d} \in \mathcal{A}^{n'}$ such that $x_{\mathbf{c}\mathbf{c}'} \in \mathcal{P}(\mathbf{d})$. (\mathbf{c} is that in Proposition 3.2.) Then the condition (a) holds because the variations of the functions $\frac{d}{dx}s(\cdot, \mathbf{a}; f)$ for $\mathbf{a} \in \mathcal{A}^{n'}$ on $\mathcal{P}(\mathbf{d})$ are bounded by $\theta_K \ell^{-n'}$, in view of (19). \square

3.3. Generic perturbations. We are going to show that the consequences of the condition $\mathbf{m}(f) > \rho \cdot \lambda_{\min}(\mathbf{T}_f)^{-1}$ given in Proposition 3.3 hold only for very small set of $f \in C_+^r(S^1; K)$. For this purpose, we next consider about perturbations of the function f .

For $f \in C_+^r(S^1; K)$ and $\varphi_i \in C^\infty(S^1)$, $1 \leq i \leq m$, we consider the family

$$(26) \quad f_{\mathbf{t}}(x) = f(x) + \sum_{i=1}^m t_i \cdot \varphi_i(x)$$

with parameter $\mathbf{t} = (t_i)_{i=1}^m \in \mathbb{R}^m$. For a point $x \in S^1$ and a finite subset $\sigma = \{\mathbf{b}_i\}_{0 \leq i \leq p}$ of \mathcal{A}^n , let $G_{x,\sigma} : \mathbb{R}^m \rightarrow \mathbb{R}^p$ be the affine map defined by

$$G_{x,\sigma}(\mathbf{t}) = \left(\frac{ds}{dx}(x, \mathbf{b}_i; f_{\mathbf{t}}) - \frac{ds}{dx}(x, \mathbf{b}_0; f_{\mathbf{t}}) \right)_{i=1}^p.$$

Note that $G_{x,\sigma}(\mathbf{t})$ is independent of f in (26). For an affine map $A : \mathbb{R}^m \rightarrow \mathbb{R}^p$, let $Jac(A)$ be the Jacobian of $DA|_{\ker(DA)^\perp}$, the restriction of the linear part DA to the orthogonal complement of its kernel when A is surjective, and put $Jac(A) = 0$ otherwise. In other words, $Jac(A)$ is the maximum among the Jacobians of the restrictions of DA to p -dimensional subspaces in \mathbb{R}^m . The following is a slight variant of [20, Proposition 16].

⁷⁾Some of B'_i 's may be empty.

Proposition 3.4. *We can choose functions $\varphi_i \in C^\infty(S^1)$, $1 \leq i \leq m$, such that, for any $x \in S^1$ and any subsets $A = \{\mathbf{a}_i\}_{1 \leq i \leq (\nu+1)(p+1)}$ of \mathcal{A}^ν , there exist a subset $A' = \{\mathbf{a}'_i\}_{0 \leq i \leq p}$ of A such that we have $\text{Jac}(G_{x,\sigma}) \geq 1$ whenever a subset $\sigma = \{\mathbf{b}_i\}_{0 \leq i \leq p}$ of \mathcal{A}^n with $n \geq \nu$ satisfies $[\mathbf{b}_i]_\nu = \mathbf{a}'_i$ for $0 \leq i \leq p$.*

The proof of Proposition 3.4 is similar to that of [20, Proposition 16]. For completeness, we give the proof in the last subsection.

3.4. The end of the proof. For $n \geq \nu$, $\mathbf{c} \in \mathcal{A}^n$ and $\sigma = (\mathbf{b}_i)_{i=0}^p \in (\mathcal{A}^n)^{p+1}$, let $Y(n, \mathbf{c}, \sigma)$ be the set of functions $f \in C_+^r(S^1; K)$ such that

$$\left| \frac{ds}{dx}(x_{\mathbf{c}}, \mathbf{b}_i; f) - \frac{ds}{dx}(x_{\mathbf{c}}, \mathbf{b}_0; f) \right| \leq 10\theta_K \cdot \ell^{-n} \quad \text{for all } 1 \leq i \leq p.$$

Note that $Y(n, \mathbf{c}, \sigma)$ is a closed subset in $C_+^r(S^1; K)$.

For $n \geq \nu$, let $Y(n)$ be the set of functions $f \in C_+^r(S^1; K)$ that belongs to $Y(n, \mathbf{c}, \sigma)$ for more than $\lceil \beta^{n(p+1)} \rceil$ combinations of $(\mathbf{c}, \sigma) \subset \mathcal{A}^n \times (\mathcal{A}^n)^{p+1}$ satisfying $\text{Jac}(G_{x_{\mathbf{c}}, \sigma}) \geq 1$. Let $Y_*(n) = \bigcup_{n'=\lfloor \delta n \rfloor}^n Y(n')$. Then $Y(n)$ and $Y_*(n)$ are also closed subsets in $C_+^r(S^1; K)$. Proposition 3.3 tells that, if $\mathbf{m}(f) > \rho \cdot \lambda_{\min}(\mathbf{T}_f)^{-1}$, then f belongs to the closed subset $\bigcap_{n \geq N} Y_*(n)$. To finish the proof of the theorem, we show that the complement of $\bigcap_{n \geq N} Y_*(n)$ is dense in $C_+^r(S^1; K)$.

Take a function $f \in C_+^r(S^1; K)$ arbitrarily and consider the family (26) with $\varphi_i \in C^\infty(S^1)$, $1 \leq i \leq m$, in Proposition 3.4. Take $\epsilon > 0$ so small that $f_{\mathbf{t}} \in C_+^r(S^1; K)$ for all $\mathbf{t} \in [-\epsilon, \epsilon]^m$. Let $X(n, \mathbf{c}, \sigma)$, $X(n)$ and $X_*(n)$ be the set of parameters $\mathbf{t} \in [-\epsilon, \epsilon]^m$ such that $f_{\mathbf{t}} \in Y(n, \mathbf{c}, \sigma)$, that $f_{\mathbf{t}} \in Y(n)$ and that $f_{\mathbf{t}} \in Y_*(n)$, respectively. From the definition of Jacobian in the last subsection, we have $\text{Leb}(X(n, \mathbf{c}, \sigma)) \leq C\ell^{-np}$ for some constant $C > 0$ that depends on θ_K , m and ϵ . Therefore, taking the number of combinations of (\mathbf{c}, σ) into consideration, we get

$$\text{Leb}(X(n)) \leq \frac{C\ell^{-np} \times \ell^n \times \ell^{(p+1)n}}{\beta^{(p+1)n}} < C(\beta^{-p}\ell^2)^n.$$

As we chose p such that $\beta^{-p}\ell^2 < 1$, we have $\text{Leb}(\bigcap_{n \geq N} X_*(n)) = 0$ and hence the complement of $\bigcap_{n \geq N} Y_*(n)$ in $C_+^r(S^1; K)$ is dense.

Remark 3.5. The proof above shows also that the condition $\mathbf{m}(f) \leq \lambda_{\min}(\mathbf{T}_f)^{-1}$ holds for a prevalent subset of $f \in C_+^r(S^1)$ in measure-theoretical sense([12, 21]).

3.5. The proof of Proposition 3.4. To prove Proposition 3.4, it is enough to show the following localized version of the claim.

Proposition 3.6. *For each $y \in S^1$, we can choose functions $\varphi_{y,i} \in C^\infty(S^1)$ for $1 \leq i \leq \ell^\nu$ and a neighborhood U_y of y such that, for any point $x \in U_y$ and any subsets $A = \{\mathbf{a}_i\}_{1 \leq i \leq (\nu+1)(p+1)}$ of \mathcal{A}^ν , there exists a subset $A' = \{\mathbf{a}'_i\}_{0 \leq i \leq p}$ of A such that we have that $\text{Jac}(G_{x,\sigma}) \geq 1$ whenever a subset $\sigma = \{\mathbf{b}_i\}_{0 \leq i \leq p}$ of \mathcal{A}^n with $n \geq \nu$ satisfies $[\mathbf{b}_i]_\nu = \mathbf{a}'_i$ for $0 \leq i \leq p$.*

In fact, once we have Proposition 3.6, we can take a finite subset $\{y(j)\}_{j=1}^J$ in S^1 so that the neighborhoods $U_{y(j)}$ in Proposition 3.6 cover S^1 and, letting $\{\varphi_i\}_{j=1}^m$ be the union of $\{\varphi_{y(j),i}\}_{i=1}^{\ell^\nu}$ for $1 \leq j \leq J$ in the corresponding conclusions of Proposition 3.6, we obtain Proposition 3.4.

Proof of Proposition 3.6. Take a point $y \in S^1$ arbitrarily. For $\mathbf{a}, \mathbf{b} \in \mathcal{A}^\nu$, we write $\mathbf{a} \prec \mathbf{b}$ if $\tau^q(\mathbf{b}(y)) = \mathbf{a}(y)$ for some $q \geq 0$. By simple combinatorial argument, we can show that this is a partial order on \mathcal{A}^ν and that, for each $\mathbf{a} \in \mathcal{A}^\nu$, there exists at most $(\nu+1)$ elements $\mathbf{b} \in \mathcal{A}^\nu$ such that $\mathbf{b} \prec \mathbf{a}$. (See the proof of [20, Proposition 16].)

For $0 < \epsilon < 1/2$ and $\mathbf{a} \in \mathcal{A}^\nu$, let $U(\epsilon)$ be the ϵ -neighborhood of y and $U_{\mathbf{a}}(\epsilon)$ the connected component of $\tau^{-\nu}(U(\epsilon))$ that contains $\mathbf{a}(y)$. We consider an integer $\mu > \nu$ that will be specified later. We then choose $\epsilon_0 > 0$ so small that $\tau^i(U_{\mathbf{b}}(\epsilon_0)) \cap U_{\mathbf{a}}(\epsilon_0) \neq \emptyset$ for some $1 \leq i \leq \mu$ only if $\mathbf{a} \prec \mathbf{b}$. Take functions $\varphi_{\mathbf{a}} \in C^\infty(S^1)$ for $\mathbf{a} \in \mathcal{A}^\nu$ supported on $U_{\mathbf{a}}(\epsilon_0)$ such that

$$\frac{d}{dx}\varphi_{\mathbf{a}}(y) = \ell^\nu \quad \text{on } U_{\mathbf{a}}(\epsilon_0/3) \quad \text{and} \quad \left| \frac{d}{dx}\varphi_{\mathbf{a}}(y) \right| < 2\ell^\nu \quad \text{on } S^1.$$

Finally let $\varphi_{y,i}$, $1 \leq i \leq \ell^\nu$ be a rearrangement of $\varphi_{\mathbf{a}}$, $\mathbf{a} \in \mathcal{A}^\nu$ and let $U_y = U(\epsilon_0/3)$.

We show that the conclusion of the proposition holds for the neighborhood U_y and the functions $\varphi_{y,i}$, $1 \leq i \leq \ell^\nu$, provided that the integer μ is sufficiently large. Consider the family (26) with $\varphi_i = \varphi_{y,i}$ and $m = \ell^\nu$ and suppose that a subset $A = \{\mathbf{a}_i\}_{1 \leq i \leq (\nu+1)(p+1)}$ of \mathcal{A}^ν is given. From the property of the partial order \prec on \mathcal{A}^ν mentioned above, we can choose a subset $A' = \{\mathbf{a}'_i\}_{0 \leq i \leq p}$ of A that consists of maximal elements in A with respect to \prec . Let $\sigma = \{\mathbf{b}_i\}_{0 \leq i \leq p}$ be a subset of \mathcal{A}^n with $n \geq \nu$ such that $[\mathbf{b}_i]_\nu = \mathbf{a}'_i$ for $0 \leq i \leq p$. For $\mathbf{b} \in \mathcal{A}^n$ and $x \in U_y$, we put

$$h_1(x, \mathbf{b}; \mathbf{t}) = \sum_{j=1}^{\min\{n, \mu\}} \ell^{-j} \frac{d}{dx} f_{\mathbf{t}}([\mathbf{b}]_j(x))$$

and

$$h_2(x, \mathbf{b}; \mathbf{t}) = \sum_{j=\min\{n, \mu\}+1}^n \ell^{-j} \frac{d}{dx} f_{\mathbf{t}}([\mathbf{b}]_j(x)),$$

so that

$$\frac{ds}{dx}(x, \mathbf{b}; f_{\mathbf{t}}) = h_1(x, \mathbf{b}; \mathbf{t}) + h_2(x, \mathbf{b}; \mathbf{t}).$$

Accordingly we decompose the affine map $G_{x,\sigma}$ into

$$G_{x,\sigma}^{(1)}(\mathbf{t}) = (h_1(x, \mathbf{b}_i; \mathbf{t}) - h_1(x, \mathbf{b}_0; \mathbf{t}))_{i=1,2,\dots,p} : \mathbb{R}^{\ell^\nu} \rightarrow \mathbb{R}^p$$

and

$$G_{x,\sigma}^{(2)}(\mathbf{t}) = (h_2(x, \mathbf{b}_i; \mathbf{t}) - h_2(x, \mathbf{b}_0; \mathbf{t}))_{i=1,2,\dots,p} : \mathbb{R}^{\ell^\nu} \rightarrow \mathbb{R}^p.$$

Let $\xi : \{1, 2, \dots, p\} \rightarrow \{1, 2, \dots, \ell^\nu\}$ be the correspondence such that $\mathbf{a}'_i(y) \in \text{supp}(\varphi_{y,\xi(i)})$ for $1 \leq i \leq p$, and consider the subspace of \mathbb{R}^{ℓ^ν} ,

$$E = \{\mathbf{t} = (t_j)_{j=1}^{\ell^\nu} \in \mathbb{R}^{\ell^\nu} \mid t_j \neq 0 \text{ only if } j = \xi(i) \text{ for some } 1 \leq i \leq p\},$$

which is naturally identified with \mathbb{R}^p . Take any point $x \in U_y$ and let $L^{(1)}$ and $L^{(2)}$ be the matrices that represent the linear part of the affine mappings $G_{x,\sigma}^{(1)} : E \rightarrow \mathbb{R}^p$ and $G_{x,\sigma}^{(2)} : E \rightarrow \mathbb{R}^p$ respectively. As a consequence of the choice of \mathbf{a}'_i , we can see that $L^{(1)}$ is the identity matrix of size p while all the entries $L^{(2)}$ are bounded by $2\ell^{-\mu+\nu}(1 - \ell^{-1})^{-1}$. Therefore, if we take sufficiently large μ , it holds

$$\text{Jac}(DG_{x,\sigma}) \geq \text{Jac}(DG_{x,\sigma}|_E) \geq 1/2.$$

Multiplying each $\varphi_{y,i}$ by 2, we can replace 1/2 by 1 on the right hand side. \square

APPENDIX A. PROOF OF THEOREM 1.4

We show that the negation of the conditions (i), (ii) and (iii) are all equivalent. First we introduce two quantities $\mathbf{n}(f, t)$ and $\mathbf{n}(f)$, similar to $\mathbf{m}(f, t)$ and $\mathbf{m}(f)$ respectively, as follows. Put $\hat{\mathbf{C}}_f = \{(x, y) \in \mathbb{R}^2 \mid |y| \leq 2\theta_f|x|\} \supset \mathbf{C}_f$. For $t \geq 0$, $z \in X_f$ and a one dimensional subspace $L \subset \mathbb{R}^2$, we define

$$\mathbf{n}(f, t, z, L) = \sum^* \frac{1}{E(\zeta, t; f)} \leq 1$$

where \sum^* is the sum over $\zeta \in (T_f^t)^{-1}(z)$ such that $(DT_f^t)_\zeta(\hat{\mathbf{C}}_f) \supset L$. Then we put

$$\mathbf{n}(f, t) = \max_{z \in X_f} \max_{L \in \mathbb{R}\mathbf{P}^1} \mathbf{n}(f, t, z, L)$$

and

$$\mathbf{n}(f) = \limsup_{t \rightarrow \infty} \mathbf{n}(f, t)^{1/t}.$$

Note that $\mathbf{n}(f, t)$ is *sub-multiplicative* with respect to t : $\mathbf{n}(f, t+s) \leq \mathbf{n}(f, t) \cdot \mathbf{n}(f, s)$. In this point, the quantity $\mathbf{n}(f, t)$ is better than $\mathbf{m}(f, t)$. In particular, the limit in the definition of $\mathbf{n}(f)$ is actually exact.

We show that $\mathbf{m}(f) = 1$ implies $\mathbf{n}(f) = 1$. For this purpose, it is sufficient to prove the claim that

$$\mathbf{m}(f, s) \leq \mathbf{n}(f, t) \quad \text{for any } t \geq 0 \text{ and } s = (b/a)t + b > t$$

where

$$a = \min_{x \in S^1} f(x) \quad \text{and} \quad b = \max_{x \in S^1} f(x).$$

Consider a point $z \in X_f$ and take $w \in T_f^{-s}(z)$. If

$$(27) \quad (DT_f^s)_\zeta(\mathbf{C}_f) \cap (DT_f^s)_w(\mathbf{C}_f) \neq \{0\}$$

for a points $\zeta \in T_f^{-s}(z)$, then we have

$$(28) \quad (DT_f^t)_{\zeta'}(\hat{\mathbf{C}}_f) \supset L := (DT_f^s)_w(\mathbb{R} \times \{0\}) \quad \text{for } \zeta' = T_f^{s-t}(\zeta) \in T_f^{-t}(z).$$

Indeed, this follows from the fact that the differences between the slope of L and those of boundary lines of $(DT_f^s)_w(\mathbf{C}_f)$ are not greater than $\ell^{-[s/b]}\theta_f$, while the differences between the slopes of the boundary lines of $(DT_f^t)_{\zeta'}(\hat{\mathbf{C}}_f)$ and those of the boundary lines of $(DT_f^t)_{\zeta'}(\mathbf{C}_f)$ are greater than $\ell^{-[t/a]-1}\theta_f = \ell^{-[s/b]}\theta_f$. Hence, in view of (2), we have that

$$\sum_{\zeta: \zeta \mapsto w} \frac{1}{E(\zeta, s; f)} \leq \sum^* \frac{1}{E(\zeta, t; f)}$$

where $\sum_{\zeta: \zeta \mapsto w}$ denotes the sum over $\zeta \in T_f^{-s}(z)$ satisfying (27) and \sum^* denotes the same sum as that in the definition of $\mathbf{n}(f, t, z, L)$. Clearly this implies the claim above.

We next show that f is cohomologous to a constant function if $\mathbf{n}(f) = 1$. Suppose $\mathbf{n}(f) = 1$. By submultiplicative property of $\mathbf{n}(f, t)$, we have $\mathbf{n}(f, t) = 1$ for all $t \geq 0$. Therefore we can take sequences of real numbers $t_n \geq 0$, points $z_n \in X_f$ and one-dimensional subspaces $L_n \in \mathbb{R}\mathbf{P}^1$ for $n \geq 1$ such that $\mathbf{n}(f, t_n, z_n, L_n) = 1$ for all $n \geq 1$ and that, as $n \rightarrow \infty$,

- $t_n \rightarrow \infty$,
- z_n converges to some $z_\infty \in X_f$, and

- L_n converges to some L_∞ (in \mathbb{RP}^1).

The condition $\mathbf{n}(f, t_n, z_n, L_n) = 1$ and (2) imply that the cone $(DT_f^{-t_n})_w(\hat{\mathbf{C}}_f)$ contains L_n for all the point $w \in T_f^{-t_n}(z_n)$. Note the unstable subspace (or the tangent space of the unstable manifold) for a backward orbit $(w(t))_{t \leq 0}$ is contained in $(DT_f^{-t})_{w(t)}(\mathbf{C}_f)$ for any $t \leq 0$. Thus, by continuity, we see that the unstable subspaces for *all* backward orbits of z_∞ coincide with each other (and with L_∞). Moreover, such property holds not only for the point z_∞ but for all the points in X_f because the set of points with such property is closed and completely invariant with respect to the flow \mathbf{T}_f .

For $x \in S^1$, let $\psi(x)$ be the slope of the (unique) unstable subspace at $(x, 0) \in X_f$. Invariance of the unstable subspaces implies that we have

$$(29) \quad \psi(\tau(x)) = (f'(x) + \psi(x))/\ell \quad \text{for all } x \in S^1.$$

Inductive use of this equality yields

$$(30) \quad \psi(x) = \sum_{n \geq 1} \sum_{\tau^n(y)=x} \ell^{-2n} f'(y) \quad \text{for all } x \in S^1,$$

where the right hand side converges in C^{r-1} sense. Since we have $\int_{S^1} \psi(x) = 0$ from (30), the function

$$\varphi(x) = \int_0^x \psi(y) dy$$

is well-defined and C^r on S^1 . It follows from (29) that

$$(31) \quad \varphi(\tau(x)) = \varphi(x) + f(x) - c \quad \text{for some constant } c.$$

That is, f is cohomologous to a constant function.

It is easy to show that \mathbf{T}_f is not weakly mixing if f is cohomologous to a constant function. Suppose that $f \in C_+^r(S^1)$ is cohomologous to a constant function, that is, f satisfies (31) for some $\varphi \in C^r(S^1)$ and $c \in \mathbb{R}$. By integrating the both sides over S^1 , we see that $c = \int_{S^1} f(x) dx > 0$. Define

$$\Phi(x, s) = \exp((2\pi i/c)(\varphi(x) + s)) \quad \text{for } (x, s) \in X_f.$$

Then $\Phi \circ T_f^t = e^{(2\pi i/c)t} \Phi$ for $t \geq 0$. Therefore T_f^t is not weakly mixing.

To finish the proof, we show that $\mathbf{m}(f) = 1$ if the semi-flow \mathbf{T}_f^t is not weakly mixing. Suppose that \mathbf{T}_f^t is not weakly mixing. Then we can find a real number $a \neq 0$ and an L^2 function Φ on X_f such that $\Phi \circ T_f^t = e^{iat} \Phi$ for $t \geq 0$. Equivalently there exists an L^2 function φ on S^1 such that $\Phi(x, s) = e^{-ias} \varphi(x)$ and that $\varphi(\tau(x)) = e^{iaf(x)} \varphi(x)$ for $x \in S^1$ and $t \geq 0$. Actually the last equality tells that φ is a C^r function and so is Φ . (For the proof of this fact, we refer that of [16, Proposition 4.2]⁸⁾, for instance.) Let $L(z)$ be the null line of the differential $D_z \Phi$. Then this line field is invariant with respect to the semi-flow \mathbf{T}_f and not tangent to the flow direction. Hence $L(z)$ is contained in the cone $(DT_f^{-t})_{w(t)}(\mathbf{C}_f)$ for any backward orbit $\{w(t)\}_{t \leq 0}$ of z and any $t \leq 0$. This and (2) imply that $\mathbf{m}(f, t) = 1$ for any $t \geq 0$ and hence that $\mathbf{m}(f) = 1$.

⁸⁾To get the proof, replace the symbolic dynamical system $\sigma : X^+ \rightarrow X^+$ and the space of Hölder functions on X^+ in the proof of [16, Proposition 4.2] by $\tau : S^1 \rightarrow S^1$ and $C^r(S^1)$ respectively.

APPENDIX B. PROOF OF LEMMA 2.4

Let $\Gamma = \mathbb{Z}_+ \times \{+, -\}$, $c(+)=1$ and $c(-)=0$. Below we write C_0 for constants that does not depend on S , h , Θ nor Θ' , while we write C for constants that may depend on them. Take an integer $\mu = \mu(S)$ such that

$$2^{-\mu+6} \|\xi\| \leq \|(DS_x)^{tr}(\xi)\| \leq 2^{\mu-6} \|\xi\| \quad \text{for any } x \in K \text{ and any } \xi \in \mathbb{R}^2.$$

Let $\nu \leq \mu - 6$ be an integer such that

$$2^{\nu-1} < \Lambda(S, \Theta', K) \leq 2^\nu.$$

So we have

$$\|DS_x^{tr}(\xi)\| \leq 2^\nu \|\xi\| \quad \text{if } x \in K \text{ and } (DS_x)^{tr}(\xi) \notin \mathbf{C}'_-.$$

We write $(m, \tau) \leftrightarrow (n, \sigma)$ if either

- $(\tau, \sigma) = (+, +)$ and $m - \mu \leq n \leq \max\{0, m + \nu + 6\}$, or
- $(\tau, \sigma) \in \{(-, -), (+, -)\}$ and $m - \mu \leq n \leq m + \mu$.

And we write $(m, \tau) \not\leftrightarrow (n, \sigma)$ otherwise.

Consider a function $u \in C^r(R)$ and put $v := Lu$. For $(n, \sigma), (m, \tau) \in \Gamma$, we define

$$v_{n,\sigma}^{m,\tau} = \psi_{\Theta',n,\sigma}(D)L(u_{\Theta,m,\tau}),$$

so that $v_{\Theta',n,\sigma} = \sum_{(m,\tau) \in \Gamma} v_{n,\sigma}^{m,\tau}$. By using Parseval's identity, we can get

$$(32) \quad \sum_{(n,\sigma) \in \Gamma} \|v_{n,\sigma}^{m,\tau}\|_{L^2}^2 \leq \|L(u_{\Theta,m,\tau})\|_{L^2}^2 \leq C_0 \gamma(S)^{-1} \|h\|_{L^\infty}^2 \|u_{\Theta,m,\tau}\|_{L^2}^2.$$

Also we have the following estimate, whose proof is postponed for a while.

Lemma B.1. *If $(m, \tau) \not\leftrightarrow (n, \sigma)$, we have*

$$(33) \quad \|v_{n,\sigma}^{m,\tau}\|_{L^2} \leq C 2^{-(r-1) \max\{m,n\}} \|u_{\Theta,m,\tau}\|_{L^2}.$$

Remark B.2. If S is an affine map in the lemma above, the Fourier transform of $L(u_{\Theta,m,\tau})$ is supported on $DS^{tr}(\text{supp}(\psi_{\Theta,m,\tau}))$, which does not meet $\text{supp}(\psi_{\Theta',n,\sigma})$ by the assumption $(m, \tau) \not\leftrightarrow (n, \sigma)$, and hence the assertion holds trivially with $v_{n,\sigma}^{m,\tau} = 0$. To prove the lemma above, we will estimate some oscillatory integrals using smoothness of S .

We first show the assertion that $|v|_{\Theta'} \leq C|u|_{\Theta}$ for some constant C , which implies that L extends boundedly to $L : W_{\dagger}(R; \Theta) \rightarrow W_{\dagger}(R; \Theta')$. By definition, we have

$$|v|_{\Theta'}^2 \leq 2 \sum_{(n,\sigma) \in \Gamma} 2^{2(c(\sigma)-\epsilon)n} \left(\left\| \sum_{(m,\tau) \leftrightarrow (n,\sigma)} v_{n,\sigma}^{m,\tau} \right\|_{L^2}^2 + \left\| \sum_{(m,\tau) \not\leftrightarrow (n,\sigma)} v_{n,\sigma}^{m,\tau} \right\|_{L^2}^2 \right)$$

where $\sum_{(m,\tau) \leftrightarrow (n,\sigma)}$ (resp. $\sum_{(m,\tau) \not\leftrightarrow (n,\sigma)}$) denotes the sum over $(m, \tau) \in \Gamma$ such that $(m, \tau) \leftrightarrow (n, \sigma)$ (resp. $(m, \tau) \not\leftrightarrow (n, \sigma)$). Since the relation $(m, \tau) \leftrightarrow (n, \sigma)$ holds only if $c(\sigma) \leq c(\tau)$ and $|m - n| < \mu$, it holds

$$\begin{aligned} \sum_{(n,\sigma) \in \Gamma} \left\| \sum_{(m,\tau) \leftrightarrow (n,\sigma)} 2^{(c(\sigma)-\epsilon)n} v_{n,\sigma}^{m,\tau} \right\|_{L^2}^2 &\leq C \sum_{(n,\sigma) \in \Gamma} \sum_{(m,\tau) \in \Gamma} 2^{2(c(\tau)-\epsilon)m} \|v_{n,\sigma}^{m,\tau}\|_{L^2}^2 \\ &\leq C \sum_{(m,\tau) \in \Gamma} 2^{2(c(\tau)-\epsilon)m} \|u_{\Theta,m,\tau}\|_{L^2}^2 \leq C|u|_{\Theta}^2 \end{aligned}$$

where the second inequality follows from (32). Also it follows from Lemma B.1 and Schwarz inequality that

$$\begin{aligned}
(34) \quad & \sum_{(n,\sigma) \in \Gamma} \left\| \sum_{(m,\tau) \not\leftrightarrow (n,\sigma)} 2^{c(\sigma)-\epsilon} n v_{n,\sigma}^{m,\tau} \right\|_{L^2}^2 \\
& \leq \sum_{(n,\sigma) \in \Gamma} \left\| \sum_{(m,\tau) \not\leftrightarrow (n,\sigma)} 2^{c(\sigma)n-c(\tau)m-(r-1-\epsilon)\max\{n,m\}} \cdot 2^{(r-1)\max\{n,m\}+(c(\tau)-\epsilon)m} v_{n,\sigma}^{m,\tau} \right\|_{L^2}^2 \\
& \leq \sum_{(n,\sigma) \in \Gamma} \left(\sum_{(m,\tau) \in \Gamma} 2^{2c(\sigma)n-2c(\tau)m-2(r-1-\epsilon)\max\{n,m\}} \right) \left(\sum_{(m,\tau) \in \Gamma} 2^{2(c(\tau)-\epsilon)m} \|u_{\Theta,m,\tau}\|_{L^2}^2 \right) \\
& \leq C|u|_{\Theta}^2.
\end{aligned}$$

Thus we obtain $|v|_{\Theta'} \leq C|u|_{\Theta}$ for $u \in C^r(R)$ and hence for $u \in W_{\dagger}(R; \Theta)$.

We next prove (10) and (11). The inequality (10) is easy to see:

$$(\|v\|_{\Theta'}^-)^2 \leq \|v\|_{L^2}^2 \leq \gamma(S)^{-1} \|h\|_{L^\infty}^2 \|u\|_{L^2}^2 \leq \gamma(S)^{-1} \|h\|_{L^\infty}^2 \|u\|_{\Theta}^2.$$

To prove (11), we begin with writing the left hand side as

$$(\|v\|_{\Theta'}^+)^2 = \|\psi_{\Theta',0,+}(D)v\|_{L^2}^2 + \sum_{n \geq 1} 2^{2n} \|\psi_{\Theta',n,+}(D)v\|_{L^2}^2.$$

The first term on the right hand side is bounded by $|v|_{\Theta'}^2$, and hence by $C|u|_{\Theta}^2$. The sum on the right hand side is bounded by

$$2 \cdot \left(\sum_{n \geq 1} \left\| \sum_{(m,\tau) \leftrightarrow (n,+)} 2^n v_{n,+}^{m,\tau} \right\|_{L^2}^2 + \sum_{n \geq 1} \left\| \sum_{(m,\tau) \not\leftrightarrow (n,+)} 2^n v_{n,+}^{m,\tau} \right\|_{L^2}^2 \right).$$

By Schwarz inequality, we have

$$\left\| \sum_{(m,+ \leftrightarrow (n,+)} 2^n v_{n,+}^{m,+} \right\|_{L^2}^2 \leq \left(\sum_{(m,+ \leftrightarrow (n,+)} 2^{2(n-m)} \right) \left(\sum_{(m,+ \leftrightarrow (n,+)} 2^{2m} \|v_{n,+}^{m,+}\|_{L^2}^2 \right)$$

where $\sum_{(m,+ \leftrightarrow (n,+)}$ denotes the sum over $m \geq 0$ such that $(m, +) \leftrightarrow (n, +)$. Note that we have $(m, \tau) \leftrightarrow (n, +)$ for $n \geq 1$ only if $\tau = +$ and $n \leq m + \nu + 6$. Thus we can see, by using (32), that

$$\sum_{n \geq 1} \left\| \sum_{(m,\tau) \leftrightarrow (n,+)} 2^n v_{n,+}^{m,\tau} \right\|_{L^2}^2 \leq C_0 \cdot 2^{2\nu} \gamma(S)^{-1} \|h\|_{L^\infty}^2 \|u\|_{\Theta}^2.$$

On the other hand, by using Lemma B.1 and Schwarz inequality, we can show

$$\sum_{n \geq 0} \left\| \sum_{(m,\tau) \not\leftrightarrow (n,+)} 2^n v_{n,+}^{m,\tau} \right\|_{L^2}^2 < C|u|_{\Theta}^2$$

in the similar manner as (34). Therefore we obtain (11). Obviously (10) and (11) imply (12) and hence L extends boundedly to $L : W_*(R; \Theta) \rightarrow W_*(R; \Theta')$. Finally we complete the proof by proving Lemma B.1.

Poof of Lemma B.1. Since K is compact, we can take closed cones $\tilde{\mathbf{C}}_+ \Subset \mathbf{C}_+$ and $\tilde{\mathbf{C}}_- \Subset \mathbf{C}_-$ such that

$$(DS_\zeta)^{tr}(\mathbb{R}^d \setminus \tilde{\mathbf{C}}_+) \Subset \mathbf{C}'_- \quad \text{for } \zeta \in K.$$

Let $\tilde{\varphi}_+, \tilde{\varphi}_- : S^1 \rightarrow [0, 1]$ be C^∞ functions satisfying

$$\tilde{\varphi}_+(\xi) = \begin{cases} 1, & \text{if } \xi \notin S^1 \cap \mathbf{C}_-; \\ 0, & \text{if } \xi \in S^1 \cap \tilde{\mathbf{C}}_-, \end{cases} \quad \tilde{\varphi}_-(\xi) = \begin{cases} 1, & \text{if } \xi \notin S^1 \cap \mathbf{C}_+; \\ 0, & \text{if } \xi \in S^1 \cap \tilde{\mathbf{C}}_+. \end{cases}$$

Recall the function χ and define $\tilde{\psi}_n(\xi) = \chi(2^{-n-1}|\xi|) - \chi(2^{-n+2}|\xi|)$ for $n \geq 1$ and

$$\tilde{\psi}_{\Theta, n, \sigma}(\xi) = \begin{cases} \tilde{\psi}_n(\xi)\tilde{\varphi}_\sigma(\xi/|\xi|), & \text{if } n \geq 1; \\ \chi(2^{-1}|\xi|), & \text{if } n = 0 \end{cases}$$

for $(n, \sigma) \in \Gamma$. Then we have $\tilde{\psi}_{\Theta, n, \sigma}(\xi) = 1$ if $\xi \in \text{supp}(\psi_{\Theta, n, \sigma})$. From the definition of the relation \hookrightarrow , there exists a constant $L > 1$, which may depend on S , such that, if $(m, \tau) \not\leftrightarrow (n, \sigma)$ and $\max\{m, n\} \geq L$, it holds

$$(35) \quad d(\text{supp}(\psi_{\Theta', n, \sigma}), (DS_\zeta)^{tr}(\text{supp}(\tilde{\psi}_{\Theta, m, \tau}))) \geq L^{-1} \cdot 2^{\max\{n, m\}} \quad \text{for } \zeta \in K.$$

In the case where $\max\{m, n\} < L$, it is easy to see that (33) holds with the constant C depending on L . Thus we assume $\max\{m, n\} \geq L$ in the following.

We consider the operator $S_{n, \sigma}^{m, \tau}$ defined by

$$S_{n, \sigma}^{m, \tau} = \psi_{\Theta', n, \sigma}(D) \circ L \circ \tilde{\psi}_{\Theta, m, \tau}(D).$$

Then we have $v_{n, \sigma}^{m, \tau} = S_{n, \sigma}^{m, \tau} u_{\Theta, m, \tau}$ because $\tilde{\psi}_{\Theta, m, \tau}(D)(u_{\Theta, m, \tau}) = u_{\Theta, m, \tau}$. We may rewrite this operator $S_{n, \sigma}^{m, \tau}$ as

$$(S_{n, \sigma}^{m, \tau} u)(x) = (2\pi)^{-4} \int V_{n, \sigma}^{m, \tau}(x, y) \cdot u \circ S(y) \cdot |\det DS(y)| dy,$$

where

$$(36) \quad V_{n, \sigma}^{m, \tau}(x, y) = \int e^{i(x-w)\xi + i(S(w)-S(y))\eta} h(w) \psi_{\Theta', n, \sigma}(\xi) \tilde{\psi}_{\Theta, m, \tau}(\eta) dw d\xi d\eta.$$

Since we have $\|u \circ S(y) \cdot |\det DS(y)|\|_{L^2} \leq C\|u\|_{L^2}$, the inequality (33) follows if the operator norm of the integral operator

$$H_{n, \sigma}^{m, \tau} : L^2(\mathbb{R}^2) \rightarrow L^2(\mathbb{R}^2), \quad H_{n, \sigma}^{m, \tau} v(x) = \int V_{n, \sigma}^{m, \tau}(x, y) v(y) dy$$

is bounded by $C \cdot 2^{-(r-1)\max\{n, m\}}$.

Apply the following formula of integration by parts for $(r-1)$ times in (36),

$$\int e^{if(w)} g(w) dw = i \cdot \int e^{if(w)} \cdot \sum_{k=1}^2 \partial_{w_k} \left(\frac{\partial_{w_k} f(w) \cdot g(w)}{\sum_{j=1}^2 (\partial_{w_j} f(w))^2} \right) dw$$

where $w = (w_k)_{k=1}^2 \in \mathbb{R}^2$. Then we obtain the expression

$$V_{n, \sigma}^{m, \tau}(x, y) = \int e^{i(x-w)\xi + i(S(w)-S(y))\eta} F(\xi, \eta, w) \psi_{\Theta', n, \sigma}(\xi) \tilde{\psi}_{\Theta, m, \tau}(\eta) dw d\xi d\eta$$

where $F(\xi, \eta, w)$ is continuous in w and C^∞ in ξ and η . Note that $F(\xi, \eta, w) = 0$ if $w \notin K$. From (35), there is a constant $C_{\alpha\beta}$ for multi-indices α and β , such that

$$(37) \quad |\partial_\xi^\alpha \partial_\eta^\beta F(\xi, \eta, w)| \leq C_{\alpha\beta} \cdot 2^{-n|\alpha| - m|\beta| - (r-1)\max\{n, m\}}$$

for $w \in \mathbb{R}^2$, $\xi \in \text{supp}(\psi_{\Theta', n, \sigma})$ and $\eta \in \text{supp}(\tilde{\psi}_{\Theta, m, \tau})$. For $n \geq 0$ and $m \geq 0$, we put

$$G_{nm}(\xi, \eta, w) = F(2^n \xi, 2^m \eta, w) \psi_{\Theta', n, \sigma}(2^n \xi) \tilde{\psi}_{\Theta, m, \tau}(2^m \eta).$$

By changes of variable, we can rewrite $V_{n, \sigma}^{m, \tau}(x, y)$ as

$$(38) \quad \int 2^{2n+2m} (\mathcal{F}_{\xi\eta}^{-1} G_{nm})(2^n(x-w), 2^m(S(w) - S(y)), w) dw$$

where $\mathcal{F}_{\xi\eta}^{-1}$ is the inverse Fourier transform with respect to the variables ξ and η . From (37), there exists a constant $C_{\alpha\beta}$ for any multi-indices α and β such that

$$|\partial_{\xi}^{\alpha} \partial_{\eta}^{\beta} G_{nm}|_{L^{\infty}} \leq C_{\alpha\beta} 2^{-(r-1) \max\{n, m\}}.$$

This implies that there exists a constant C such that

$$|\mathcal{F}_{\xi\eta}^{-1} G_{nm}(x, y, w)| \leq C 2^{-(r-1) \max\{n, m\}} (1 + |x|^2)^{-2} (1 + |y|^2)^{-2}.$$

Applying this inequality in the expression (38) of $V_{n, \sigma}^{m, \tau}(x, y)$, we obtain the required estimate for $H_{n, \sigma}^{m, \tau}$ from Young's inequality. \square

APPENDIX C. PROOF OF LEMMA 2.3

We prove the inequality (8). For the inequality (9), we refer [3, Lemma 7.1]. Recall the argument in the proof of Lemma 2.4 in Appendix B, setting $S = id$. Notice that the assumptions of Lemma 2.4 then hold since we assume $\Theta' < \Theta$. Put $v_i = g_i \cdot u$ for $1 \leq i \leq I$. Then we have

$$(39) \quad \sum_{i=1}^I (\|v_i\|_{\Theta'}^2)^2 \leq \sum_{i=1}^I \|v_i\|_{L^2}^2 \leq \|u\|_{L^2}^2 \leq C \|u\|_{\Theta}^2.$$

We have proved, in the proof of Lemma 2.4, that

$$\sum_{n \geq 0} 2^{2n} \left\| \sum_{(m, \tau) \not\leftrightarrow (n, +)} \psi_{\Theta', n, +}(D)(g_i u_{\Theta, m, \tau}) \right\|_{L^2}^2 \leq C |u|_{\Theta}^2$$

Since we can and do put $\mu = 6$ in the setting $S = id$, the relation $(m, \tau) \leftrightarrow (n, +)$ holds only if $|m - n| \leq 6$. Hence we have, by Schwarz inequality,

$$\begin{aligned} & \sum_{i=1}^I \sum_{n \geq 0} 2^{2n} \left\| \sum_{(m, \tau) \leftrightarrow (n, +)} \psi_{\Theta', n, +}(D)(g_i u_{\Theta, m, \tau}) \right\|_{L^2}^2 \\ & \leq 13 \cdot \sum_{i=1}^I \sum_{n \geq 0} \sum_{m: |m-n| \leq 6} 2^{2n} \|\psi_{\Theta', n, +}(D)(g_i u_{\Theta, m, +})\|_{L^2}^2 \\ & \leq 13 \cdot 2^{12} \cdot \sum_{m \geq 0} \sum_i 2^{2m} \|g_i u_{\Theta, m, +}\|_{L^2}^2 \leq C_0 \|u\|_{\Theta}^2. \end{aligned}$$

Therefore we obtain $\sum_{i=1}^I (\|v_i\|_{\Theta'}^2)^2 \leq C_0 \|u\|_{\Theta}^2 + C |u|_{\Theta}^2$. This and (39) yield (8).

REFERENCES

1. A. Avila, S. Gouëzel & M. Tsujii, *Smoothness of solenoidal attractors*, Discrete and Continuous Dynamical Systems **15** (2006), no. 1, 21–35.
2. V. Baladi, *Positive transfer operators and decay of correlations*. Advanced Series in Nonlinear Dynamics, 16. World Scientific, 2000
3. V. Baladi & M. Tsujii, *Anisotropic Hölder and Sobolev spaces for hyperbolic diffeomorphisms*, Ann. Inst. Fourier **57**(2007), no. 1, 127–154
4. V. Baladi & B. Vallée, *Exponential decay of correlations for surface semi-flows without finite Markov partitions*. Proc. Amer. Math. Soc. **133** (2005), no. 3, 865–874
5. R. Bowen, *Equilibrium states and the ergodic theory of Anosov diffeomorphisms*, Lecture note in Math. **470**, Springer, 1975
6. R. Bowen, *Periodic orbits for hyperbolic flows*. American J. Math. **94** (1972), 1-30
7. D. Dolgopyat, *On decay of correlations in Anosov flows*, Ann. Math. **147** (1998), 357–390,
8. D. Dolgopyat, *Prevalence of rapid mixing in hyperbolic flows*, Ergodic Theory Dynam. Systems **18** (1998), no. 5, 1097–1114.
9. D. Dolgopyat, *Prevalence of rapid mixing in hyperbolic flows II, topological prevalence*. Ergodic Theory Dynam. Systems **20** (2000), no. 4, 1045–1059.
10. S. Gouëzel and C. Liverani, *Banach spaces adapted to Anosov systems*, Ergodic Theory Dynam. Systems, **26**(2006), no. 1, 189–217
11. S. Gouëzel and C. Liverani, *Compact locally maximal hyperbolic sets for smooth maps: fine statistical properties*, preprint(2006)
12. B.R. Hunt, T. Sauer & J.A. Yorke, *Prevalence: a translation-invariant "almost every" on infinite-dimensional spaces*, Bull. Amer. Math. Soc., **27**, no. 2, 217–238(1992); Addendum *ibid.* **28**, no. 2, 306–307, (1993)
13. H. Hennion, *Sur un théorème spectral et son application aux noyaux lipschitziens*, Proc. Amer. Math. Soc. **118** (1993), 627–634.
14. C. T. Ionescu Tulcea & G. Marinescu, *Théorie ergodique pour des classes d'opérations non complètement continues*. Ann. of Math. (2) **52** (1950), 140–147.
15. C. Liverani, *On contact Anosov flows*, Ann. Math. **159** (2004), 1275–1312.
16. W. Parry & M. Pollicott, *Zeta functions and the periodic orbit structure of hyperbolic dynamics*, Astérisque **187-188** (1990)
17. M. Pollicott, *On the mixing of Axiom A attracting flows and a conjecture of Ruelle*. Ergodic Theory Dynam. Systems **19** (1999), no. 2, 535–548.
18. D. Ruelle, *A measure associated with Axiom A attractors*, Amer. J. Math. **98** (1976), 616–654
19. Ya. G. Sinai, *Gibbs measures in ergodic theory*, Russ. Math. Surveys **27** (1972), 21–70
20. M. Tsujii, *Fat solenoidal attractors*, Nonlinearity **14** (2001), no. 5, 1011–1027.
21. M. Tsujii, *A measure on the space of smooth mappings and dynamical system theory*, J. Math. Soc. Japan **44** (1992), no. 3, 415–425

DEPARTMENT OF MATHEMATICS, KYUSHU UNIVERSITY, FUKUOKA, 812-8581 JAPAN
E-mail address: tsujii@math.kyushu-u.ac.jp