

**CRITICAL TWO-POINT FUNCTION FOR LONG-RANGE
MODELS WITH POWER-LAW COUPLINGS: THE MARGINAL
CASE FOR $d \geq d_c$**

AKIRA SAKAI (HOKKAIDO UNIVERSITY)

Consider the long-range models on \mathbb{Z}^d of random walk, self-avoiding walk, percolation and the Ising model, whose translation-invariant 1-step distribution/coupling coefficient decays as $|x|^{-d-\alpha}$ for some $\alpha > 0$. In the previous work (Ann. Probab., 43, 639–681, 2015), we have shown in a unified fashion for all $\alpha \neq 2$ that, assuming a bound on the “derivative” of the n -step distribution (the compound-zeta distribution satisfies this assumed bound), the critical two-point function $G_{p_c}(x)$ decays as $|x|^{\alpha \wedge 2 - d}$ above the upper-critical dimension $d_c \equiv (\alpha \wedge 2)m$, where $m = 2$ for self-avoiding walk and the Ising model and $m = 3$ for percolation.

In this talk, I will show in a much simpler way, without assuming a bound on the derivative of the n -step distribution, that $G_{p_c}(x)$ for the marginal case $\alpha = 2$ decays as $|x|^{2-d}/\log|x|$ whenever $d \geq d_c$ (with a large spread-out parameter L). This solves the conjecture in the previous work, extended all the way down to $d = d_c$, and confirms a part of predictions in physics (Brezin, Parisi, Ricci-Tersenghi, J. Stat. Phys., 157, 855–868, 2014). The proof is based on the lace expansion and new convolution bounds on power functions with log corrections.